

The influence of black cumin seed oil therapy with dosage of 1.5 mL/day and 3 mL/day to interleukin21 (IL-21) expression of the patients with metabolic syndrome risk

*By Akrom*

**PAPER • OPEN ACCESS**

The influence of black cumin seed oil therapy with dosage of 1.5 mL/day and 3 mL/day to interleukin-21 (IL-21) expression of the patients with metabolic syndrome risk

2

To cite this article: D R Fajar *et al* 2017 *IOP Conf. Ser.: Mater. Sci. Eng.* **259** 012012

View the [article online](#) for updates and enhancements.

## Related content

34

- [The efficacy of black cumin seed \(\*Nigella sativa\*\) oil and hypoglycemic drug combination to reduce HbA1c level in patients with metabolic syndrome risk](#)  
P N R Rachman, Akrom and E Darmawan

- [Topical Review](#)  
Anil S Modak

12

- [The emerging role of reactive oxygen and nitrogen species in redox biology and some implications for plasma applications to medicine and biology](#)  
David B Graves

## The influence of black cumin seed oil therapy with dosage of 1.5 mL/day and 3 mL/day to interleukin-21 (IL-21) expression of the patients with metabolic syndrome risk

D R Fajar<sup>1</sup>, Akrom<sup>1,2</sup>, E Darmawan<sup>1</sup>

<sup>1</sup>Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

<sup>2</sup>Center of Drug Information and Study, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

E-mail: desi.rf1991@gmail.com

**Abstrak.** Metabolic syndrome (MS) is a metabolic disorder caused by obesity and insulin resistance. In the presence of obesity and insulin resistance occurs increased fat metabolism that causes production reactive oxygen species (ROS) and oxidative stress that makes dysregulation of adipose tissue that decreases antioxidant enzymes and immune system disorders. In diabetes mellitus there is pancreatic  $\beta$  cell damage caused by pro-inflammatory cytokines i.e interleukin-21 (IL-21). Black cumin seed oil (BCSO) contains antioxidants and immunomodulators but has not seen how it affects to IL-21. This study used cross over design to determine the effect of BCSO with dose of 1.5 mL/day and 3 mL/day dose of IL-21 in MS patients. The total subjects 66 of MS patients were divided into 2 groups. Group I was administered 1.5 mL/day and 3 mL/day of BCSO in dosage for 20 days and continued with washing out (BCSO discontinued) for 7 days. On the day 28, group I administrated 3 mL/day of BCSO and group II were given 1.5 mL/day of BCSO during 20 days. Statistical analysis showed that mean value expression of IL-21 were given by BCSO at dose 1.5 mL/day was  $5.06 \pm 5.48$  and BCSO at dose 3 mL/day  $4.66 \pm 3.63$  ( $p < 0.05$ ). The result showed that IL-21 expression in patient at risk of MS who received BCSO dose 3 mL/day were lower than those received adose of 1.5 mL/day for 20 days.

**Keywords:** black cumin seed oil, interleukin-21, metabolic syndrome risk

### 1. Introduction

Metabolic syndrome (MS) is a set of abnormalities involving various physiological, biochemical, clinical and metabolic factors that can lead to the development of cardiovascular disease and diabetes mellitus and all-cause mortality risk [1]. Some of the risks causing MS include lifestyle (diet, smoking, physical activity), genetics and socioeconomics [2]. MS in Indonesia is one of the main public health problems such as diabetes mellitus, central obesity, dyslipidemia, and hypertension, which is a component of MS and has an effect on the prevalence of metabolic syndrome. MS case rate in Indonesia reached 23% i.e 26.6% in women and 18.3% in men of productive age, this data is based on the result of basic health research [3].

The pathophysiology of metabolic syndrome is not known for certain, but some literature say that the primary reason is a person's obesity state. Excess fat in the body causes increased fat metabolism



Content from this work may be used under the terms of the Creative Commons Attribution 3.0 licence. Any further distribution of this work must maintain attribution to the author(s) and the title of the work, journal citation and DOI.

Published under licence by IOP Publishing Ltd

16

along with reactive oxygen species (ROS) production as well as increased oxidative stress characterized by a balance disorder of oxidation reduction reactions and decreased antioxidant enzymes in both adipose and tissue cells, this situation occurs in hypertension and atherosclerosis [4].

In addition, fat cell enlargement will secrete metabolic products such as pro-inflammatory cytokines, procoagulants, inflammatory peptides and angiotensinogen [5]. Insulin resistance affects type 2 of diabetes and hyperglycemia. A person with diabetes, oxidative stress will inhibit the taking of glucose in muscle cells and fat cells and decrease insulin secretion by pancreatic  $\beta$  cells (type 2 diabetes). This is due to autoimmune disorders mediated by lymphocytes T and activated dendritic cells involving inflammatory pathways Thelper1 (Th1), Thelper2 (Th2) and Thelper17 (Th17) and inhibition of T regulators. It is found that the Th17 pathway is involved in the inflammatory process and damage to pancreatic beta cells regulated by interleukin-21, interleukin-23 and interleukin-27. Interleukin-21 is a type 1 of cytokine [6, 7].

BSCO contains antioxidant and immunomodulatory substances which can be used as antidiabetes mellitus [8]. The main content of BSCO is essential oils such as timokuinon that have an effect on the immune system through inhibition of translocation of NF-kB into the nucleus so as to decrease pro-inflammatory cytokines, chemokines, adhesion factors and coagulation factor [33]. In addition, it is known that BSCO can decrease the oxidative stress condition that triggers the synthesis of pro-inflammatory cytokines through adhesion molecules. Interleukin-21 is a pro-inflammatory cytokine that plays a role in pancreatic B cell damage [35] it is not yet known how the effect of BSCO with a dose of 1.5 mL/day and 3 mL/day for 20 days in patients with metabolic syndrome risk.

## 2. Materials and Methods

This study used cross over design with a total of 66 patients with metabolic syndrome who state willing to follow the study by agreeing inform consent. The inclusion criteria were obesity, blood pressure  $\geq 140/90$  mmHg, triglyseride level  $\geq 150$  mg/dL or using the drug of methabolic syndrom and 18-70 years old. Exclusion criteria were pregnant and lactating women, tuberculosis and be thought immun disease. But during the course of the study there were 4 patients who were excluded due to side effects such as nausea, vomiting, gastritis, increased blood pressure and decreased kidney function. This research was conducted at Jetis I Public Health Centers, Bantul Regency, Yogyakarta in August 2016 until March 2017. This clinical trial protocol has been reviewed and approved by health research ethics committee of Faculty of Medicine and Health Sciences, Universitas Muhammadiyah Yogyakarta with approval number 279/EP-FKIK-UMY/VIII/2016.

Blood steering patients were taken from vena cubity by analysts in laboratory of the Jetis I Public Health Centers, Bantul Regency, Yogyakarta. Procedures undertaken in the implementation have been reviewed by research ethics committee of Universitas Muhammadiyah Yogyakarta. Blood that has been accommodated in a vacutainer containing anticoagulants is taken to the pharmacology laboratory of Universitas Gadjah Mada (UGM) and ready to be examined with flow cytometer. Here is the scheme of research implementation:




Figure 1. Schematic of research design

2

3. Results and Discussion

3.1. The characteristics data of patients at risk of metabolic syndrome at Jetis I Public Health Centers, Bantul Regency in 2016

From the previous studies, demographic data and characteristic data of metabolic syndrome as primary outcome and percentage of IL-21 expression as outcome secunder. From the data, statistical test using Chy-Square test and paired T-test sample with 95% confidence level.

This research is conducted at Jetis 1 Public Health Centers, Bantul Regency, Yogyakarta in August 2016 - March 2017. We involved doctors in getting patien 28 who will be joined in the research and obtained a total of 66 patients at risk of metabolic syndrome who have met the inclusion and exclusion criteria. However, during the study there were 4 patients who drop out because they complain the side effects such as nausea, vomiting, gastritis, increased blood 32 ssure and decreased kidney function. The 27 tal subjects who followed the study to the end were 62 patients at risk of metabolic syndrome.

Table 1 shows the characteristics of patients at risk of metabolic syndrome. Based on gender, it is known that the metabolic syndrome was mostly suffered by women (78.80%) compared to men (21.20%). Metabolic syndrome was also more prevalent in the age group 51-60 years (42.40%), married status (98.50%), high school education (78.80%) and employment status (57.60). This result is in line with research conducted before [9] which reported that there was a significant relationship among gender, age, and work with the incidence of metabolic syndrome. Metabolic syndrome is more risky in women than in men. In women there is an excessive accumulation of fat in the adipose tissue that causes hypertri 7 ceride. Metabolic syndrome in women is also caused by diet, genetics and lack of physical activity than men [9, 10]. Women are more at risk of dev 7 oping diabetes mellitus due to body mass index (BMI) that is more easily increased and the distribution of body fat easily accumulated due to hormonal processes in the body [11].

**Table 1.** The characteristics data of patients at risk of metabolic syndrome at Jetis I Public Health Centers in 2016

Characteristics of patients	n=66	Percentage (%)
Gender		
Men	14	21.20
Women	52	78.80
Age		
26 years old	1	1.50
41-50 years old	20	30.30
51-60 years old	28	42.40
>60 years old	17	25.80
Marital status		
Married	65	98.50
Single	1	1.50
Education		
No school	6	9.10
Elementary-High School	52	78.80
College-Master	8	12.10
Working		
Work	38	57.60
Not working	28	42.40

The incidence of metabolic syndrome is more experienced by age group > 40 years than age < 40 years [9]. Metabolic syndrome occurs as one gets older. This is due to a decrease in organ function and metabolic disorders that exist in the body. Diabetes mellitus occurs due to decreased insulin secretion because of the pancreatic  $\beta$  cells damage, insulin resistance, impaired metabolism of carbohydrates, proteins and fats and reduced elasticity of blood vessels may be at risk of hypertension and atherosclerosis leading to the incidence of metabolic syndrome [9, 12, 13]. Metabolic syndrome has a significant relationship with one's marital status [14, 15]. A married person will be faced with a social environment that allows problems and causes unhappiness. Each problem will create psychosocial stress potential that may be at risk of metabolic syndrome [15]. Education and work are two characteristics of metabolic syndrome which has interconnected each other. Education can determine the type of work of a person with different income to meet their daily needs. Lifestyle change can increase the prevalence of diseases such as diabetes, obesity and heart disease which is a factor risk of metabolic syndrome [16]. Job status relates to daily physical activity. In this study, patients with metabolic syndrome are more likely than workers, such as civil servants, entrepreneurs, farmers, fishermen and laborers with mild to severe activity. Physical activity can help lower cholesterol, triglycerides, increase HDL and insulin sensitivity and lower blood pressure which is a risk factor for the incidence of metabolic syndrome and cardiovascular disease complications [11].

Table 2 shows the characteristics data of patients with metabolic syndrome. Metabolic syndrome is more prevalent in patients at risk of metabolic syndrome with prolonged <5 years (59.10%) compared with 5 years of age (40.90%). Patients at risk for metabolic syndrome who participate in this study mostly suffered from diabetes mellitus with complications. Diabetes mellitus is a metabolic disease with characteristics of hyperglycemia due to abnormalities of insulin secretion and become a risk factor for organ damage and dysfunction especially in the eyes, kidneys, nerves, heart and blood vessels.

**Table 2.** The characteristics data of patients at risk of metabolic syndrome at Jetis I Bantul Public Health Centers in 2016

Characteristics of patients	n=66	Percentage(%)
Characteristics of metabolic syndrome		
DM	10	15.20
Hypertension	1	1.50
Hypertriglyceride	1	1.50
DM+ Hypertension	16	24.20
DM+ Hypercholesterolemia	2	3.00
DM+ Obesity	1	1.50
DM+ Hypertriglyceride	6	9.10
DM+ Dyslipidemia	3	4.50
DM+ Hypertension + Hypertriglyceride	12	18.20
DM+ Hypertension + Hypercholesterolemia	5	7.60
DM+ Hypertension + Obesity	2	3.00
DM+ Hypertension + Dyslipidemia	5	7.60
DM+ Hypertension + Hypertriglyceride + Obesity	2	3.00
Routine Medication		
DM Drug	25	37.90
Hypertension Drug	1	1.50
Dyslipidemia Drug	1	1.50
DM Drug+ Hypertension Drug	33	50.00
DM Drug +Dyslipidemia Drug	3	4.50
DM Drug + Hypertension Drug + Dyslipidemia Drug	3	4.50

3.2. Clinical examination result of patients at risk of metabolic syndrome before and after using BCSO Dose 1.5 mL/day and 3 mL/day

Patients at risk of metabolic syndrome before treatment are examined clinical conditions such as systolic and diastolic blood pressure, pulse rate, body mass index (BMI), using a tool that has been calibrated by the calibration laboratory of Universitas Ahmad Dahlan Yogyakarta. Examination of blood glucose level (BG), triglyceride level (TG) and SGOT and SGPT activity using 5010 spectrophotometer at the laboratory of Jetis I Public Health Centers, Bantul Regency, Yogyakarta. The data were analyzed using repeated measure ANOVA test and continued with Bonferroni test as shown in Table 3.

**Table 3.** The results of measurement of clinical condition of patients at risk of metabolic syndrome before and after using BCSO dose 1.5 mL/day and 3 mL/day at Jetis I Public Health Centers, Bantul Regency, Yogyakarta in 2016

Clinical Conditions of Patients	Average ± SD, n=62			p Values
	Pretreatment	Dose 1.5 mL/day	Dose 3 mL/day	
SBP(mmHg)	143.06±17.98 <sup>a</sup>	137.73±17.66	136.92±17.66	0.00
DBP(mmHg)	80.78±9.25	79.27±13.96	79.95±10.98	0.58
Pulse (x/min)	88.66±13.82 <sup>a</sup>	90.69±12.91	92.44±12.15	0.04
BMI (kg/m <sup>2</sup> )	24.16±4.13 <sup>a</sup>	24.79±4.27	24.35±5.10	0.00
BG (mg/dL)	246.69±109.30 <sup>ab</sup>	184.18±89.46	197.77±87.14	0.00
TG (mg/dL)	193.84±114.61 <sup>a</sup>	173.42±108.51	158.58±97.87	0.00
SGOT (U/L)	23.00±10.41	21.82±6.47	22.60±9.95	0.63
SGPT (U/L)	22.29±10.81	21.16±7.86	21.03±12.08	0.82

Description: a= there is a significant difference between the measurement before treatment and the dose measurement 3 mL/day; B= there is a significant difference between the measurement before treatment and the dose measurement of 1.5 mL/day.

The results of statistical analysis show that there are significant differences between measurement of systolic blood pressure (SBP), pulse, BMI, BG and TG before and after BCSO administration in patients at risk of metabolic syndrome ( $p < 0.05$ ). The results of statistical analysis show that there is no significant difference between diastolic blood pressure (DBP) measurement, SGOT and SGPT activities before and after BCSO administration in patients at risk of metabolic syndrome ( $p > 0.05$ ). These results indicate that giving BCSO dose of 1.5 mL/day and 3 mL/day for 20 days can decrease the mean blood pressure value in patients at risk of metabolic syndrome with targeted therapy  $< 140/90$  mmHg. This is in line with research conducted which states that the provision of black cumin seed powder can reduce blood pressure during 20 days of administration in patients with metabolic syndrome [18]. Another study mentioned that giving black seeds of cumin seeds for 8 weeks can reduce the blood pressure of systolic and diastolic without side effects [19]. This study shows that black seeds of cumin seeds for 20 days are able to increase pulse rate and BMI in patients at risk of metabolic syndrome but still within normal limits. According to the American Heart Association (AHA) the normal range of pulse rate is 60-100 x/min and according to the World Health Organization (WHO) the normal value of IMT in patients at risk of metabolic syndrome is  $< 30$  kg/m<sup>2</sup>. The results of this study stated that the administration of BCSO dose 1.5 mL/day and 3 mL/day for 20 days can reduce blood glucose levels in time and triglycerides in patients at risk of metabolic syndrome. BCSO contains active compounds such as thymoquinone, nigelon and thymol which play a role in lowering blood glucose. Black cumin in adipose cells can stimulate the secretion of insulin produced by pancreatic cells and is shown to have no effect on tissue sensitivity to glucose [20]. Criteria for metabolic syndrome according to International Education Diabetes Federation (IDF) in hypertension, triglycerides  $> 150$  mg/dL, HDL for women  $< 50$  mg/dL and men  $< 40$  mg/dL or in treatment of tubercidemia, diagnosed type 2 DM with GDP  $> 100$  mg/dL [21].

Statistical analysis shows that SGOT and SGPT activities in patients at risk of metabolic syndrome decrease after using BCSO dose 1.5 mL/day and 3 mL/day for 20 days but not statistically significant ( $p > 0.05$ ). This is because the patients who participate in this study had a mean value of SGOT and SGPT activity in normal limits. The incidence of metabolic syndrome is associated with fatty liver disease or NAFLD. In patients with insulin resistance will experience steatosis due to increased free fatty acids to the liver increased. Oxidative stress and cytokines can cause disruption to the function of cells and liver organ [22]. SGOT and SGPT can be an indicator of liver damage with SGOT values  $< 35$  U/L and SGPT  $< 41$  U/L.

3.3. Measurements of IL-21 expression in patients at risk of metabolic syndrome given BCSO dosage 1.5 mL/day and 3 mL/day

Measurements of IL-21 expression using a flow cytometer tool in the pharmacology laboratory of the Faculty of Medicine, Universitas Gajah Mada (UGM), Yogyakarta. Data on IL-21 expression measurements in patients at risk of metabolic syndrome given BCSO dose 1.5 mL/day and 3 mL/day for 20 days statistically analyzed using paired T-test as shown in Table 4.

**Table 4.** The results of IL-21 expression measurements in patients at risk of metabolic syndrome given BCSO dose 1.5 mL/day and 3 mL/day for 20 days

Clinical condition patient	Mean value ± SD		p Value
	Dose 1.5 mL/day (n=62)	Dose 3 mL/day (n=62)	
IL-21	5.06±5.48	4.66±3.63	0.69

The results of statistical analysis shows that the mean value of IL-21 expression in patients at risk of metabolic syndrome who received BCSO dose 3 mL/day was lower than those who received BCSO dose 1.5 mL/day but not statistically significant ( $p > 0.05$ ). Several factors that may influence the research results are the possibility of influence from previous treatment before cross over design on BCSO dose 1.5 mL/day and 3 mL/day for 20 days. Physiological conditions of the body and lifestyle



such as lack of exercise, smoking and taking drugs for long periods in patients at risk of metabolic syndrome can also affect the results of the study.

Metabolic syndrome occurs because of insulin resistance. This is due to the disruption of pancreatic  $\beta$  cell function and insulin hypersecretion resulting in insulin resistance, in addition to severe damage to pancreatic  $\beta$  cells that leads to decreased insulin production and hyperglycemia. The deficiency of IL-21 expression in diabetic nonobese mice (NOD) may lead to resistance to insulinitis, insulin autoantibody production and the onset of type I diabetes [7]. Excess of IL-21 expression in pancreatic  $\beta$  cells makes of inducing chemokines, inflammatory cytokines such as IL-17A, IL-17F, IFN- $\gamma$ , monocyte chemoattractant protein (MCP)-1 and MCP-2. IL-21 is a type I cytokine produced by T-cells and NK cells and is detected in CD4<sup>+</sup> and T-helper undergoing differentiation [23].

Giving *Nigella sativa* L. can cause demodulator in pro-inflammatory cytokines produced by T-helper-1 and cytokines. The active substance contained in the BCSO can evoke the immune system with ability to raise T-helper cell, Tsupressor cells, and NK cells are a product of lymphocytes [24, 25]. T-cells play an important role in inducing inflammation induces makrofaf and secrete inflammatory and pro-inflammatory cytokines in adipose tissue leads to insulin resistance and lead to diabetes mellitus. Inflammatory cytokines would improve insulin resistance through the role of inflammatory signals such as janus kinase (JK) and activation of transcription factors nuclearfactor kappaB (NF-kB). Signaling pathways such as JK and NF-kB may reduce the action of insulin in adipocyte tissue [26, 27].

Black cumin seed can be utilized as an antioxidant and immunomodulator. Hexane extracts of black cumin seeds (*Nigella sativa* L.) can improve macrophage phagocytic activity in DMBA-induced female rats in vitro [8]. Part of the most widely used black cumin seed plant is its oil. Black cumin seed oil contains essential oil content of non-ar thymoquinone that is 27.8% [28]. The working mechanism of thymoquinone is by inhibiting non-enzymatic lipid peroxidation in liposomes and has the potential to neutralize free radicals and inhibit the pro-inflammatory cytokines such as interleukin-1 $\beta$ , interleukin-8 and chemokines such as machrophage chemotactic protein (MCP)-1 with the mechanism of action via inhibition of TNF- $\alpha$  which induces activation of transcription factors nuclearfactor kappaB (NF-kB) signaling pathway is involved in IL-21 and inhibiting the translocation of NF-kB into the nucleus [29]. BCSO delivery in the metabolic syndrome will inhibit IL-21 activation by decreasing the deference of T-helper which also produces IL-21 (IL-21R) receptors as well as inhibition of the STAT 3 signaling pathway.

#### 4. Conclusion

IL-21 expressions in patients at risk of metabolic syndrome who received BCSO dose 3 mL/day were lower than those received a dose of 1.5 mL/day for 20 days.

#### References

- [1] Kaur J 2014 A comprehensive review on metabolic syndrome, *Cardiology Research and Practice*, ID 943162.
- [2] Sandra R 2015 Sindrom metabolik *Jurnal Majority* 4 (4) 88, Fakultas kedokteran Univeristas Lampung.
- [3] Ministry of Health 2014 Riset Kesehatan Dasar (RISKESDAS 2013), Jakarta.
- [4] Stocker R and Keaney JF 2004 Role of oxidative modification in atherosclerosis *J. Physiol.* 84 (5) 1381-1392.
- [5] Semiardji 2004 The significant of visceral fat in metabolic syndrome, Jakarta: *Diabetes meeting* 9-10 Oktober.
- [6] Dandona P, Aljada A and Chaudhuri A 2005 Metabolic syndrome: A comprehensive perspective based on interaction between obesity and diabetes *Circulation* 111(11):1448-54.
- [7] Andrew P, Tom, V B, Andrea L, Suto Akira, Michaud M, Zhang D, Grusby M and Herrath M V 2009 Interleukin-21 is required for the deelopment of type 1 diabetes in NOD Mice *Diabetes* 58(5) 1144-1155.

- [8] Akrom, 2013, Mekanisme kemoprefentif ekstrak heksan BJH pada tikus SD diinduksi DMBA: kajian antioksidan dan immunomodulator *Disertasi* Program Doktor Ilmu Kedokteran dan Kesehatan, FK UGM, Yogyakarta.
- [9] Suhaema and Masthalina H 2015 Pola Konsumsi dengan Terjadinya Sindrom metabolik di Indonesia *Jurnal Kesehatan Masyarakat Nasional* **9** 4.
- [10] Chen H 2006 Enhancing energy and glucose metabolism by disrupting triglyceride synthesis: lesson from Mice Lacking DGAT-1 *Nutrition & Metabolism* **3** 10.
- [11] Irawan D 2010 Prevalensi dan faktor resiko kejadian diabetes melitus tipe 2 di daerah Urban Indonesia *Tesis* Universitas Indonesia; Jakarta.
- [12] Jelantik I M and Haryati E 2014 Hubungan faktor resiko umur, jenis kelamin, kegemukan, hipertensi dengan kejadian diabetes mellitus tipe 2 di wilayah kerja Puskesmas Mataram *Media Bina Ilmiah* **8** 1 39-44.
- [13] Zahtamal, Chandra F, Suryanto and Restuastuti T 2007 Faktor pasien diabetes mellitus, *Berita Kedokteran Masyarakat* **23** (3) 142-147.
- [14] Mohammad A G, Abdulbari B, Asia N, Ismail, Mahmoud Z, Waleed K, Abdullatef, Abdullah OAA and Al Hamaq 2010 Gender and age related differences in patients with the metabolic syndrome in a highly endogamous population *J. Basic Med. Sci.* **10** (3) 210-217.
- [15] Troxel K A, Matthews L C, Gallo L H and Kuller 2005 Marital quality and occurrence of the metabolic syndrome in women *Archives of Int. Med.* **165**(9) 1022-1027.
- [16] Praditwi R 2008 Faktor risiko yang mempengaruhi terjadinya penyakit sindrom metabolik, *Skripsi*, Fakultas kesehatan masyarakat, Universitas Airlangga; Surabaya.
- [17] Bertrais S, Beyeme JP, Czernichow S, Galan P, Hercberg S and Oppert JM 2005 Sedentary behaviors, physical activity and metabolic syndrome in middle-aged French subject *Obesity Res.* **13** 936-944.
- [18] Najmi A, Haque S F, Naseeruddin M and Khan R A 2008 Effect of *Nigella sativa* oil on various clinical and biochemical parameters of metabolic syndrome *Int. J. Diabetes & Metab.* **16** 85-87.
- [19] Husaeni H F, Amini M, Mohtashami R, Ghamarchehre M E, Sadeqhi Z, Kianbakht S, Husein A F 2013 Blood pressure lowering effect of *Nigella sativa* L. seed oil in healthy volunteers: A randomized, double-blind, placebo-controlled clinical Trial *J. Phytother. Res.* **21** 12.
- [20] Andaloussi AB, Martineau LC, Spoor D, Vuong T, Leduc C, Joly E 2008 Antidiabetic activity of *Nigella sativa* seed extract in cultured pancreatic- $\beta$  cells, skeletal muscle cells and adipocytes *Pharmaceut. Biol. J.* **49** 96-104.
- [21] Mittal S 2008 *The Metabolic Syndrome in Clinical Practice* (London: Springer-Verlag).
- [22] Medina J, Fernandez S, Garcia B, and Moneno O 2004 Approach the pathogenesis and treatment of nonalcoholic steatohepatitis *Diabetes care* **27** 2057-2066.
- [23] Monteleone G, Pallone F and Macdonald TT 2009 Interleukin-21 as a new therapeutic target for immune-mediated diseases, *Trends in Pharmacological Sciences* **30** 8.
- [24] Islam and Muhammad T 2016 Biological activities and therapeutic promises of *Nigella sativa* L *Int. J. of Pharma Sci. and Scientific Res.* **2** 6 242-243
- [25] Gilani AH, Jabeen Q and Khan MAU, 2004 Review of medical uses and pharmacological activities of *Nigella sativa* L seeds *Pak. J. Biol. Sci.* **4** 441-451.
- [26] Arkan M C, Hevener A L, Greten F R, Maeda S, Li Z W and Long J M 2005 IKK beta links inflammation to obesity-induced insulin resistance *Nat. Med.* **11**:191.
- [27] Hirosumi J, Tuneman G, Chang L, Gorgun CZ, Uysal KT and Maeda K 2002 A central role for JNK in obesity and insulin resistance *Nature J.* **6**:333.
- [28] Claudia C, Maria G, Hanganu D, Olah N, Maria F, Hammam C and Hammam M 2010 Chemical composition of the tunisian *Nigella sativa* Note I. Profile on essential oil *Farmacia* **58** 4.

- [29] Ricky TN 2011 Pengaruh pemberian terapi tambahan adjuvan minyak jinten hitam (*Nigella sativa*) terhadap hitungLimfosit mencit Balb/C model sepsis *Skripsi*, Fakultas Kedokteran Universitas Sebelas Maret: Surakarta.

# The influence of black cumin seed oil therapy with dosage of 1.5 mL/day and 3 mL/day to interleukin21 (IL-21) expression of the patients with metabolic syndrome risk

ORIGINALITY REPORT

# 13%

SIMILARITY INDEX

PRIMARY SOURCES

- 1** [openaccess.city.ac.uk](https://openaccess.city.ac.uk) Internet 109 words — 3%
- 2** D A Perwitasari, S Urbayatun, I N Faridah, N Masyithah. "Relationship of therapeutic outcome with quality of life on type 2 diabetes mellitus patients in Abdul Aziz Singkawang hospital", IOP Conference Series: Materials Science and Engineering, 2017 Crossref 48 words — 1%
- 3** Tetsuji Shimizu, Yuzuru Ikehara. "Benefits of applying low-temperature plasma treatment to wound care and hemostasis from the viewpoints of physics and pathology", Journal of Physics D: Applied Physics, 2017 Crossref 27 words — 1%
- 4** [www.cambridge.org](https://www.cambridge.org) Internet 25 words — 1%
- 5** Encyclopedia of Behavioral Medicine, 2013. Crossref 24 words — 1%
- 6** [tomf.org](https://tomf.org) Internet 22 words — 1%
- 7** D A Perwitasari, I N Faridah, Y Kulle, M Yulistika. "Relationship between perception with the quality of life of T2DM patients in Dok II Jayapura Hospital", IOP Conference Series: Materials Science and Engineering, 2017 Crossref 20 words — < 1%

8	"European Obesity Summit (EOS) - Joint Congress of EASO and IFSO-EC, Gothenburg, Sweden, June 1 - 4, 2016: Abstracts", Obesity Facts, 2016	18 words — < 1%
Crossref		
9	TH17 Cells in Health and Disease, 2011.	15 words — < 1%
Crossref		
10	www.medical-genetics.com	13 words — < 1%
Internet		
11	"Posters", Diabetic Medicine, 12/2006	13 words — < 1%
Crossref		
12	www.citeulike.org	13 words — < 1%
Internet		
13	B. V. Iversen. "IN SITU, ON-SITE AND LABORATORY MEASUREMENTS OF SOIL AIR PERMEABILITY: BOUNDARY CONDITIONS AND MEASUREMENT SCALE :", Soil Science, 02/2001	12 words — < 1%
Crossref		
14	repository.ubaya.ac.id	12 words — < 1%
Internet		
15	www.chansonalkalinewater.com	10 words — < 1%
Internet		
16	dro.deakin.edu.au	10 words — < 1%
Internet		
17	ai.jsaweb.jp	9 words — < 1%
Internet		
18	Amin, Faridah, Malik Hassan Mehmood, Bina S. Siddiqui, Nasima Khatoon, and Anwarul-Hassan Gilani. "Co-administration of Black seeds and Turmeric shows enhanced efficacy in preventing metabolic syndrome in fructose-fed rats :", Journal of Cardiovascular Pharmacology, 2014.	9 words — < 1%
Crossref		

19	<a href="http://bioclima.ro">bioclima.ro</a> Internet	9 words — < 1%
20	<a href="http://www.wjgnet.com">www.wjgnet.com</a> Internet	9 words — < 1%
21	Jefferson C. Frisbee. "Obesity, Insulin Resistance, and Microvessel Density", <i>Microcirculation</i> , 6/2007 Crossref	9 words — < 1%
22	Nurkhasanah, R D Santoso, R Fauziah. " The immunomodulatory effect of ethanolic extract on phagocytic activity, nitrit oxide and reaxtive oxygen intermediate secretions of macrophage in mice ", <i>IOP Conference Series: Materials Science and Engineering</i> , 2017 Crossref	9 words — < 1%
23	<a href="http://jbiomedsci.biomedcentral.com">jbiomedsci.biomedcentral.com</a> Internet	8 words — < 1%
24	ElKoshairy, Nehal, Rania Hassan, and Ahmed Halawa. "The Effect of Lead Toxicity on Albino Rats\ Filiform and Fungiform Papillae and the Possible Protective Role of Honey and Black Seed", <i>Journal of Environmental and Occupational Science</i> , 2014. Crossref	8 words — < 1%
25	<a href="http://www.fasebj.org">www.fasebj.org</a> Internet	8 words — < 1%
26	Izabela Cichocka, Jan Krupa. "The preferences in selecting dietary supplements used in slimming", <i>Polish Journal of Public Health</i> , 2016 Crossref	8 words — < 1%
27	Bener, A., M. Abdulmalik, M. Al-Kazaz, A.-G. Mohammed, R. Sanya, S. Buhmaid, M. Al-Harthy, and M. Zirie. "Medical Audit of the Quality of Diabetes Care: Is Primary Care More Successful Than Hospitals?", <i>Journal of Primary Care &amp; Community Health</i> , 2012. Crossref	8 words — < 1%

---

28 N Harun, E Darmawan, L H Nurani. " Effect of calyx capsule-ethanol extract L. on renal function of healthy volunteers ", IOP Conference Series: Materials Science and Engineering, 2017 8 words — < 1%  
Crossref

---

29 [www.ncbi.nlm.nih.gov](http://www.ncbi.nlm.nih.gov) 8 words — < 1%  
Internet

---

30 "EFFECTS OF REPLACING ANTIBIOTIC GROWTH PROMOTERS (AGPS) WITH BOTANICAL EXTRACTS AND OILS IN FEED OF ", Journal of Animal and Plant Sciences, August 31 2016 Issue 6 words — < 1%  
Publications

---

31 Sutherland, Andrew P.R. van Belle, Tom W. "Interleukin-21 is required for the development of type 1 diabetes in NOD mice.(ORIGINAL ARTICLE)(non", Diabetes, May 2009 Issue 6 words — < 1%  
Publications

---

32 Wakana Ohashi, Yuichi Hattori. "Drugs that Affect Lipid Metabolism", Elsevier BV, 2014 6 words — < 1%  
Crossref

---

33 Tilg, Herbert, and Alexander Moschen. "Inflammatory and Anti-Inflammatory Mediators Secreted by Adipose Tissue", Oxidative Stress and Disease, 2009. 6 words — < 1%  
Crossref

---

34 Sakine Yalçın. "Effects of dietary black cumin seed (*Nigella sativa* L.) on performance, egg traits, egg cholesterol content and egg yolk fatty acid composition in laying hens", Journal of the Science of Food and Agriculture, 2009 6 words — < 1%  
Crossref

---

35 Esposito, Katherine Ciotola, Miryam Carl. "Effect of rosiglitazone on endothelial function and inflammatory markers in patients with the metabo", Diabetes Care, May 2006 Issue 5 words — < 1%

## Publications

---

EXCLUDE QUOTES      OFF  
EXCLUDE BIBLIOGRAPHY   ON

EXCLUDE MATCHES      OFF