Response Letter to Editor and Reviewer

Dear editor and reviewer

Thank you so much for reviewing our manuscript and raising many valuable comments towards our paper. We have made some corrections and promoted our manuscript according to these comments, which are shown as follows:

1. Comment on abstract:

with aged 17 years and above \rightarrow Must mention the upper limit

Author's reply:

Thank you for the suggestion. We have added the upper limit as suggested. Therefore, we have revised in our abstract. The change is highlighted in green within the manuscript.

Revision

An analytical cross-sectional study was conducted among 227 respondents living in Indonesia with aged 17 to 70 years old.

2. Comments on introduction:

a. These behaviours are significantly influenced by how much danger they perceive the event to be, how likely to occur, how effective their current coping behaviours are, and what they believe they can do to solve the problem (McCaffrey, 2004) \rightarrow Kindly adjust the citation format

Author's reply:

Thank you for the suggestion. We have incorporated these suggestion by adjust the citation format in our article

Revision :

These behaviours are significantly influenced by how much danger they perceive the event to be, how likely it is to occur, how effective their current coping behaviours are, and what they believe they can do to solve the problem¹⁴.

The COVID-19 risk perception is considered an essential aspect of health and risk communication as its goal is to understand what risks of COVID-19 to the public are and how the public addresses them ¹⁴

b. These influencing factor refer to sex, gender, region, education level, occupation, marital status, monthly personal income, income condition, direct cash assistance, health status, quarantine conditions, history of chronic illness, smoking history and supplement use \rightarrow Kindly mention the reference source for each factor. If it doesn't exist, the authors have to justify it

Author's reply:

Thank you for the suggestion. The aims of our study was to explore illness risk perception and efficacy belief of COVID-19 and its related individual characteristic factors. Therefore, we have revised the statement in the manuscript.

Revision :

Therefore, our study investigated the individual characteristic factors influencing COVID-19 risk perception and efficacy belief in different outbreak stages, when the number of cases increased significantly. Those factors include sex, gender, region, education level, occupation, marital status, monthly personal income, income condition, direct cash assistance, health status, quarantine conditions, chronic illness, smoking history and supplement use.

3. Comments on method:

a. The target population was Indonesian active social media user \rightarrow By social media, is that meant by a certain conversation platform (Twitter, Instagram, etc)? Or are one-way social media users like blogs also included?

Author reply:

Thank you for pointing this out. We launched an online survey through certain social networking sites such Facebook, Twitter, Whatapp, and Instagram. We do not include one-way social media platform like blogs. The revision statement is :

The target population was Indonesian active social media user who used specific platforms such as Facebook, Twitter, Whatapp and Instagram.

2. The eligibility criteria of participants were the Indonesian people aged 17 years above \rightarrow This requirement is too loose, the authors should include an age limit requirement and target a certain generation, considering that several generations use social media differently.

Author reply:

We agree with your suggestion. We have revised the eligibility criteria by including the age limit requirement, and target a certain generation. The revision statement is :

The eligibility criteria of participants were Indonesian people aged 17 to 70 years old. We classified them into several age groups such adolescence (17 to 25 years old), adult (26 to 45 years old), elderly (46 to 65 years old) and geriatric (above 65 years old).

3. The online questionnaire was distributed via Google Form \rightarrow Mention the link for the survey conducted

Author reply :

We accept the reviewer suggestion. The link for survey have added in the article. The revision statement is :

The online questionnaire was distributed via a link of Google Form as follows: http://bit.ly/WHOQOLID.

4. Why is each dimension rated on a different Likert scale? (10,5,4)

Author reply :

Thank you for raising this question.

The measurement of risk perception based on the constructs of the protection motivation theory (PMT). The perceived severity assessed the severity of COVID-19 using a 10-point Likert

scale, from 1 (not severe) to 10 (very severe). Meanwhile, the perceived vulnerability assessed the likelihood of acquiring this disease using a 5-point Likert scale, from 1 (very unlikely) to 5 (very likely). The questionnaire used in this study was adapted from a previous study, whereas each perceived dimension was rated on a different Likert scale. Furthermore, we calculated the perceived threat as the overall risk perception measure, which was determined by the formula as follows (the square root of the multiplication of severity/2 and vulnerability). In order to achieve a level of comparability between the scores, the severity score was initially divided by two. A square root transformation was performed to normalize the skewed distribution of the new variable, resulting in a scale ranging from 1 (low) to 5 (high) for measuring perceived threat.

We have also added those explanations in the method section

4. Comments on results and discussion.

a. The study sampled 232 eligible subjects filled the questionnaire with a response rate of $94.8\% \rightarrow$ Authors should also state if there are certain groups that dominate the survey (such as certain college students).

Author reply :

Thank you for pointing this out. We agree with the suggestion. Therefore, we have stated the certain groups that dominate the survey. The revision statement is :

The majority of participants which dominated the survey were female (56.8%), adult (60.4%), living in the western region (74.4%), holding higher degrees in education (63.9%) and married (67.4%).

b. In Table I, what is the age limit for each age category?

Author reply :

We have revised the Table 1 and included the age limit for each age category. The revision of Table 1 is :

Variables	n	%
Sex		
Male	98	43.2
Female	129	56.8
Age		
Adolescent (17 to 25		
years old)	50	22.0
Adult (26 to 45 years		
old)	137	60.4
Elderly (46 to 65 years		
old)	37	16.3
Geriatric (above 65		
years old)	3	1.3

Table 1. Demographic characteristic of respondents

Region

Western region	169	74.4
Middle region	56	24.7
Eastern region	2	0.9
Education		
Primary Education	12	5.3
Middle Education	70	30.8
Higher Education	145	63.9
Occupation		
Student	33	14.5
Private sector employee	48	21.1
Government worker	37	16.3
Entrepreneur	32	14.1
Others	77	33.9
Marital Status		
Married	153	67.4
Single	59	26.0
Widow/Widower	15	6.6
Monthly Personal		
Income (IDR)		
Low income	9	4.0
Lower middle income	56	247
Upper middle income	92	40.5
High income	70	30.8
Income Conditions		
Decreased income	118	52.0
Increased revenue	2	0.9
No changes	101	44.5
No income	6	2.6
Direct Cash Assistance		
Yes	14	6.2
No	213	93.8
Health Status		
Healthy	205	90.3
Do not know	22	9.7
Quarantine Conditions		
Full time activities at		
home	32	14.1
Still leaving the house 2-		
3x a week is not for		
work	51	22.5
Work outside the home		
every day	82	36.1
Work outside the home		
2-3x a week	46	20.3
Others	16	7.0
History of Chronic		
Illness		
Yes	11	4.8
No	216	95.2
Smoking History		

No 180 79.3 Supplements Use 134 59.0	Yes	47	20.7
Supplements UseYes13459.0	No	180	79.3
Yes 134 59.0	Supplements Use		
	Yes	134	59.0
No 93 41.0	NT-	03	41.0

c. Still in Table I, are there certain provisions in the division of the region? Like based on the Indonesian time zone for example?

Author reply :

Thank you so much. We agree with your suggestion and have added this statement in the manuscript. The revision was in Table 1 :

Table 1. Demographic characteristic of respo	ondents

Variables	n	%
Sex		
Male	98	43.2
Female	129	56.8
Age		
Adolescent (17 to 25		
years old)	50	22.0
Adult (26 to 45 years		
old)	137	60.4
Elderly (46 to 65 years		
<mark>old)</mark>	37	16.3
Geriatric (above 65		
years old)	3	1.3
<mark>Region (Indonesian time</mark>		
<mark>zone)</mark>		
Western Region	<mark>169</mark>	<mark>74.4</mark>
Middle Region	<mark>56</mark>	<mark>24.7</mark>
Eastern Region	<mark>2</mark>	<mark>0.9</mark>
Education		
Primary Education	12	5.3
Middle Education	70	30.8
Higher Education	145	63.9
Occupation		
Student	33	14.5
Private sector employee	48	21.1
Government worker	37	16.3
Entrepreneur	32	14.1
Others	77	33.9
Marital Status		
Married	153	67.4
Single	59	26.0
Widow/Widower	15	6.6

Monthly Personal		
Income (IDR)		
Low income	9	4.0
Lower middle income	56	247
Upper middle income	92	40.5
High income	70	30.8
Income Conditions		
Decreased income	118	52.0
Increased revenue	2	0.9
No changes	101	44.5
No income	6	2.6
Direct Cash Assistance		
Yes	14	6.2
No	213	93.8
Health Status		
Healthy	205	90.3
Do not know	22	9.7
Quarantine Conditions		
Full time activities at		
home	32	14.1
Still leaving the house 2-		
3x a week is not for		
work	51	22.5
Work outside the home		
every day	82	36.1
Work outside the home		
2-3x a week	46	20.3
Others	16	7.0
History of Chronic		
Illness		
Yes	11	4.8
No	216	95.2
Smoking History		
Yes	47	20.7
No	180	79.3
Supplements Use		
Yes	134	59.0
No	93	41.0

d. Regardless, a majority of respondents believed the COVID-19 pandemic was severe and fatal \rightarrow Where did this conclusion come from?

Author reply :

Thank you so much for raising this question. In our study reported that a majority of respondents believed the COVID-19 pandemic was severe and fatal. This conclusion come from the Table II which reported the mean score of perceived severity of respondents was male (8.38+2.04) and female (8.59+2.06). These high score indicated that the perceived severity of COVID-19 among male and female was severe and fatal.

The revision statement is :

Table II reported that the mean score of perceived severity of respondents was male (8.38+2.04) and female (8.59+2.06). This high score indicated that the perceived severity of COVID-19 among males and females was severe and fatal.

5. Comments on conclusions.

a. Most of the respondents believed that they were susceptible to COVID-19 disease \rightarrow What research data support this statement? Meanwhile in Table I, the majority of respondents are in a favorable condition (Health status, chronic illness, smoking, supplements)

Author reply :

Thank you so much for pointing out this question. In our study reported that most of the respondents believed they were susceptible to COVID-19 diseases. This conclusion come from the Table II which reported the mean score of perceived threat of respondents was male (3.30+0.91) and female (3.25+0.82). The high score indicated that the respondent believed the threat of a COVID-19 risk to be serious and susceptible.

Considering the main purpose of study, we decided to remove the sentences of "Most of the respondents believed that they were susceptible to COVID-19 disease" from the conclusions

6. Also, add Funding detail, Data availability, and Authors contributions (according to the guidelines and templates)

Funding detail

The authors received no financial support for the research, authorship, and/or publication of this article.

Data availability None

Authors contribution

Lolita, Lolita : study design, methodology, data collection, validation, and writing -original draft. Azis Ikhsanudin : data management, data collection, visualization, statistical analysis, editing.

We sincerely hope that our revised manuscript could be suitable for publication in your journal. Best regards!

Lolita, M.Sc., Apt Assistant Professor, Department of Clinical Pharmacy, Ahmad Dahlan University Prof Dr Soepomo Warungboto Umbulharjo, Yogyakarta, Indonesia Email: lolita_ur@yahoo.com

[bjop] Editor Decision

2022-11-11 08:39 AM

Lolita Lolita, Azis Ikhsanudin:

The editing of your submission, "Illness Risk Perceptions and Efficacy Beliefs Among Indonesian In The Course of COVID-19 Pandemic," is complete. We are now sending it to production.

Submission URL: <u>https://journal.umpr.ac.id/index.php/bjop/authorDashboard/submission/3287</u>

Mohammad Rizki Fadhil Pratama

Borneo Journal of Pharmacy

Manuscript Assessment Form Borneo Journal of Pharmacy

Manuscript title

Illness Risk Perceptions and Efficacy Beliefs Among Indonesian In The Course of COVID-19 Pandemic

Comments on **abstracts**.

with aged 17 years and above \rightarrow Must mention the upper limit

Comments on the **introduction**.

- 1. These behaviours are significantly influenced by how much danger they perceive the event to be, how likely to occur, how effective their current coping behaviours are, and what they believe they can do to solve the problem (McCaffrey, 2004) → Kindly adjust the citation format
- 2. These influencing factor refer to sex, gender, region, education level, occupation, marital status, monthly personal income, income condition, direct cash assistance, health status, quarantine conditions, history of chronic illness, smoking history and supplement use \rightarrow Kindly mention the reference source for each factor. If it doesn't exist, the authors have to justify it

Comments on **methods**.

- 1. The target population was Indonesian active social media user → By social media, is that meant by a certain conversation platform (Twitter, Instagram, etc)? Or are one-way social media users like blogs also included?
- 2. The eligibility criteria of participants were the Indonesian people aged 17 years above → This requirement is too loose, the authors should include an age limit requirement and target a certain generation, considering that several generations use social media differently.
- 3. The online questionnaire was distributed via Google Form → Mention the link for the survey conducted
- 4. Why is each dimension rated on a different Likert scale? (10,5,4)

Comments on **results and discussion**.

- 1. The study sampled 232 eligible subjects filled the questionnaire with a response rate of 94.8% \rightarrow Authors should also state if there are certain groups that dominate the survey (such as certain college students).
- 2. In Table I, what is the age limit for each age category?
- 3. Still in Table I, are there certain provisions in the division of the region? Like based on the Indonesian time zone for example?
- 4. Regardless, a majority of respondents believed the COVID-19 pandemic was severe and fatal → Where did this conclusion come from?

Comments on **conclusions**.

Most of the respondents believed that they were susceptible to COVID-19 disease \rightarrow What research data support this statement? Meanwhile in Table I, the majority of respondents are in a favorable condition (Health status, chronic illness, smoking, supplements)

Comments against references.

Please write another review that has not been covered above. Also, add Funding detail, Data availability, and Authors contributions (according to the guidelines and templates)

Overall score (scale 0 to 9): ___7

Recommendation

□ Accept submission/No revision

☑ Revisions required/Minor revision (please write an outline of the revisions to the text, and the revised text please return to us)

 \Box Resubmit for review/Major revision (please write if there are specifics below)

□ Decline submission/Manuscript rejected (please write the reason)



Illness Risk Perception and Efficacy Belief Among Indonesian In The Course of COVID-19 Pandemic Lolita Lolita^{*1,2}, Azis Ikhsanudin^{*1,3}

¹Undergraduate programme, Faculty of Pharmacy, Universitas Ahmad Dahlan, Special District of Yogyakarta Province, Indonesia

²Research Division of Clinical Pharmacology, The First Affiliated Hospital of Nanjing Medical University, Jiangsu Province, P.R. China

³Doctoral Programme, Faculty of Pharmacy, Universitas Gadjah Mada, Special District of Yogyakarta Province, Indonesia

Author hidden^{*1,2}

*email: hidden

Keywords: COVID-19 Perceived Risk Efficacy Beliefs Indonesia

Abstract

COVID-19, a worldwide pandemic, has posed a significant challenge for public health systems worldwide. Health risk perception and efficacy belief are primary constructs in influencing individuals' protective behavior due to the outbreak. Our study investigated each item of illness risk perception, efficacy belief, and its related factors concerning the COVID-19 pandemic. An analytical cross-sectional study was conducted among 227 respondents with aged 17 to 70 years old. Data collection was conducted using convenience sampling by distributing the web questionnaire between April and July 2020. Mann-Whitney or Kruskal-Wallis bivariate analysis was performed using SPSS version 21.0 to assess the relationship between individual characteristic factors among the illness risk perception and efficacy belief. The study established that respondents had a medium to a high level of illness risk perception and a reasonable efficacy belief in dealing with the COVID-19 pandemic. Region (p=0.027) and occupation (p=0.036) differences were significantly associated with the threat and severity perception, respectively. Smoking history (p=0.037), supplement use (p=0.029), and occupation (p=0.018) differences were significantly associated with self-efficacy. Meanwhile, gender (p=0.045) differences were significantly associated with response efficacy. Therefore, the public's illness risk perception and efficacy belief could be substantial in planning, modifying and implementing a coordinated response for risk communication in current and future epidemics.

Received: Month Year

Accepted: Month Year

Published: Month Year



© year The Authors. Published by Institute for Research and Community Services Universitas Muhammadiyah Palangkaraya. This is Open Access article under the CC-BY-SA License (http://creativecommons.org/licenses/by-sa/4.0/). DOI: https://doi.org/10.33084/bjop.vxix.xxx.

INTRODUCTION

The novel coronavirus, SARS-CoV-2, was identified as a cluster cause of atypical cases of pneumonia in Wuhan, China¹. WHO has declared this coronavirus disease (COVID-19) a global health emergency due to many confirmed cases in more than 70 countries globally². Indonesia, the world's fourth most populous country, was reported as having confirmed two cases of COVID-19 infection on March 2, 2020. To date, the number of COVID-19 case keeps increasing rapidly in this country ³. Over the study period, the latest data regarding COVID-19 reported an increase significantly with an average of over 1790 confirmed cases with 113 new cases, 170 dead cases, and 112 recovered cases ⁴.

The Indonesian government has issued several restrictive measures to curtail the spread of the virus across the nations. However, those policies affect an individual psychologically, which causes frustration, anxiety, and even the need to change their daily life behavior ^{5 6 7}. A systematic review reported that these pandemic has led to high mental disorder rates among the general population⁸. Separate inline, another study has also stated that quarantine measures could worsen a person's psychological condition, such as depression, anxiety, stress disorder, and health risk perception ⁹.

Illness risk perception and efficacy beliefs are reliable predictors of preventive health behavior¹⁰ ¹¹. Illness risk perception is a subjective assessment to respond to fearful communications about a health threat. It could relate with the efficacy beliefs as individual capabilities in taking protective action behavior towards a potential threat ¹². Health behavior theories suggested that perceptions of illness risks related to perceptions of vulnerability, severity, and threat¹³. Individuals perceiving significant risks were more likely to implement protective behaviors. These behaviors are significantly influenced by how much danger they perceive the event to be, how likely it is to occur, how effective their current coping behaviors are, and what they believe they can do to solve the problem ¹⁴. Therefore, monitoring risk perceptions and efficacy beliefs is an integral part of public health emergency management.

The COVID-19 risk perception is considered an essential aspect of health and risk communication as its goal is to understand what risks of COVID-19 to the public and how the public addresses them ¹⁴. During the COVID-19 pandemic, the public will have different efficacy beliefs that will influence how people react to risk ¹³. Our previous study found that the perceived risk of acquiring COVID-19 was low when there was no confirmed case among Indonesian ¹⁵. Meanwhile, the COVID-19 perceived threat was high at the beginning of outbreaks from March 3rd to 27th, 2020 16. People are more considered that COVID-19 is a life-threatening danger to them at that point. Therefore, our study investigated the individual characteristic factors influencing COVID-19 risk perception and efficacy beliefs in different outbreak stages when the number of cases increased significantly. Those factors include sex, gender, region, education level, occupation, marital status, monthly personal income, income condition, direct cash assistance, health status, quarantine conditions, chronic illness, smoking history and supplement use. In collaborating with the private sector, the Indonesian government has pursued comprehensive policies such as large-scale social distancing, work from home, region quarantine, self-isolation, face mask use,

and social distancing to prevent the transmission of COVID-19¹⁷. Hence, understanding risk perception and efficacy belief will give public health authorities a vital reference about protective behavior among Indonesian. Furthermore, these results will determine the willingness of the Indonesian's efforts and contribution to handling COVID-19.

MATERIALS AND METHODS

Study Design and Data Collection:

The study has been reviewed for ethical considerations and obtained approval from the University of Aisyiyah Ethics Committee 1305/KEP-Research (No. UNISA/IV/2020). This cross-sectional online-survey was conducted from April to July 2020. The target population was Indonesian active social media users who used specific platforms such as Facebook, Twitter, Whatapp and Instagram. According to Global Digital, reported by Hootsuite and We are Social, active social media user in Indonesia has reached 150 million people per January 2020¹⁸. The eligibility criteria of participants were Indonesian people aged 17 to 70 years old, active social media users who resided in Indonesia and could give informed consent. We classified the participants into several age groups such as adolescence (17 to 25 years old), adult (26 to 45 years old), elderly (46 to 65 years old) and geriatric (above 65 years old). Exclusion criteria were those non-Indonesian residents who did not complete to respond to one or more online survey items. The minimum sample sizes of 220 participants were selected using the survey system sample size calculator, an online survey software package, with 95% confidence and a 5% significance level. Quantitative data was generated from a questionnaire containing closed-ended questions. This

study was voluntary and anonymous. The individuals' consent was obtained before data collection.

Research Instrument and Study Variable:

The instrument was designed based on previous SARS research¹⁹, translated and modified to Indonesian¹⁶. Prior to the distribution of the questionnaires, reliability tests were carried out. The pilot test was conducted on a total of 30 study participants. The assessing instrument for risk perception and efficary belief were reliable. The Cronbach's alpha and the validity test for risk perceptions were 0.806 and 0.782, while efficacy beliefs were 0.703 and 0.612. The online questionnaire was distributed via link of Google Form as follows: a http://bit.Ly/WHOQOLID. The questionnaire comprised two sections: sociodemographic characteristics and risk perception with efficacy beliefs. The first section comprised questions on respondent sociodemographic characteristics, namely age, sex, region, education level, occupation, marital status, personal income, income condition, direct cash assistance, health status, quarantine conditions, history of chronic illness, smoking history and the use of supplements. The second section consisted of a question about perceived risk and efficacy beliefs. There are three dimensions of risk perception : perceived threat, vulnerability, severity. In comparison, efficacy beliefs are associated with response efficacy and self-efficacy. The measurement of risk perception is based on the construct of the protection motivation theory (PMT). The perceived severity assessed the severity of COVID-19 using a 10point Likert scale, from 1 (not severe) to 10 (very severe). Meanwhile, the perceived vulnerability assessed the likelihood of acquiring this disease using a 5-point Likert scale, from 1 (very unlikely) to 5 (very likely). The questionnaire used in this study was adapted from a previous study, whereas each perceived dimension was rated on a different Likert scale. Furthermore, we calculated the perceived threat as the overall risk perception measure, which was determined by the formula as follows (the square root of the multiplication of severity/2 and vulnerability). In order to achieve a level of comparability between the scores, the severity score was initially divided by two. A square root transformation was performed to normalize the skewed distribution of the new variable, resulting in a scale ranging from 1 (low) to 5 (high) for measuring perceived threat ¹⁹. The perceived threat rating was on a scale from 1 to 5, with 1 being "low" and 5 being "high". The response-efficacy was assessed by asking participants to respond to how confident they believe others around them would be in taking practical actions to prevent contracting COVID-19 using a 4-point Likert scale from 1 (not at all) to 4 (very much). Additionally, self-efficacy was determined by asking how confident people felt that they could prevent contracting the disease. The respondents were asked each question on a rating scale from 1 ("not confident") to 4 (very confident). Respondents completed a survey concerning these categories.

Data Analysis:

A descriptive statistical analysis was used to examine the frequency of data on socio-demographic characteristics, risk perception, and efficacy belief toward COVID-19. All the variables were tested for normality using the Kolmogorov-Smirnov test, and none of the variables was normally distributed. Therefore Kruskal-Wallis test and

Mann-Whitney test were employed to determine significant differences in the categorical independent variable (socio-demographics) on the dependent variable of risk perception (perceived vulnerability, perceived severity, perceived severity) and efficacy beliefs (response efficacy, self-efficacy). We have analyzed the data using SPSS version 21.0. Values of p less than 0.05 were considered statistically significant.

RESULTS AND DISCUSSION

The study sampled 232 eligible subjects who filled the questionnaire with a response rate of 94.8%. After excluding 5 participants with incomplete data, a final sample of 227 subjects were required in the current study. The majority of participants who dominated the survey were female (56.8%), adult (60.4%), living in the western region (74.4%), holding higher degrees in education (63.9%) and married (67.4%). Overall, 89% of the participants had good health, 59% used supplements, and 4.8% had a prior history of chronic illness. Regarding income conditions, they still work outside the home daily (36.1%), whereas 52.0% have decreased income during the pandemic. Only 6.2% of participants provided direct financial aid from the government. The sociodemographics of the participant are listed in Table 1.

Tab	le 1.	Demograp	hic cl	haracteri	istic	of res	pondents
-----	-------	----------	--------	-----------	-------	--------	----------

Variables	n	%
Sex		
Male	98	43.2
Female	129	56.8
Age		
Adolescent (17 to 25 years old)	50	22.0
Adult (26 to 45 years old)	137	60.4
Elderly (46 to 65 years old)	37	16.3
Geriatric (above 65 years old)	3	1.3
Region (Indonesian time zone) Western Region	169	74.4

Middle Region	56	24.7
Eastern Region	2	0.9
Education		
Primary Education	12	5.3
Middle Education	70	30.8
Higher Education	145	63.9
Occupation		
Student	33	14.5
Private sector employee	48	21.1
Government worker	37	16.3
Entrepreneur	32	14.1
Others	77	33.9
Marital Status		
Married	153	67.4
Single	59	26.0
Widow/Widower	15	6.6
Monthly Personal Income (IDR)		
Low income	9	4.0
Lower middle income	56	247
Upper middle income	92	40.5
High income	70	30.8
Income Conditions		
Decreased income	118	52.0
Increased revenue	2	0.9
No changes	101	44.5
No income	6	2.6
Direct Cash Assistance		
Yes	14	6.2
No	213	93.8
Health Status		
Healthy	205	90.3
Do not know	22	9.7
Quarantine Conditions		
Full time activities at home	32	14.1
Still leaving the house 2-3x a week		
is not for work	51	22.5
Work outside the home every day	82	36.1
Work outside the home 2-3x a		
week	46	20.3
Others	16	7.0
History of Chronic Illness		
Yes	11	4.8
No	216	95.2
Smoking History		
Yes	47	20.7
No	180	79.3
Supplements Use		
Yes	134	59.0
No	93	41.0

Table II revealed a statistically significant difference between the efficacy responses between men and women (p=0.045). Men participants had a significantly higher mean of response efficacy than women. Therefore, they are more confident in being able to take action in trying to prevent COVID-19. Moreoever, men are physically stronger and emotionally more stable than women. Thus they are more willing to take precautions to reduce their risk of COVID-19²⁰. It is also likely because men have a lower immune system, which can be attributed to their differences in innate and adaptive immune responses. Sex-specific responses result from X chromosome inheritance which contains genes associated with high immunity ²¹. Therefore, men perceived a higher efficacy response to prevent them from contracting COVID-19 during a pandemic.

This study also showed a significant difference in perceived threat between regions (p=0.027). Participants in western region had a significantly lower mean of a perceived threat than those in middle or eastern regions. People living in the western region perceive that they are less likely to be exposed to the COVID-19 threat. The highest number of cases in Indonesia is in the western regions. World Health Organization estimates that as of February 3, 65.8% of Indonesia's cumulative confirmed cases have been reported on Java Island.

In contrast, Jakarta has the highest number of confirmed cases per one million, followed by East Kalimantan, North Kalimantan, Special District of Yogyakarta and Central Java²². The low perception of threat among people in the western region could affect adherence to health protocols. These regions also have high mobility and population density, where many business industrial centers are still operating continuously. It will be a potential cause of the increasing number of confirmed cases in this area.

Occupation differences also have a statistically significant relationship with perceived severity (p=0.036) and selfefficacy (p=0.018). Those who work in government have a significantly higher perceived seriousness than those who work in the private sector or entrepreneurship. It means that if government workers suffer from COVID, it will severely threaten them. The potentially higher risk of severe outcomes for COVID-19 depends on the worker's characteristics in various occupations ²³. Previous research has also demonstrated that government employees have the highest risk of serious adverse outcomes due to COVID-19 ²⁴. Furthermore, our study found that those in the private sector have a greater sense of self-efficacy than others. Private companies have stringent rules in issuing their employees' policies regarding work regulations and health protection due to the COVID-19 pandemic ²⁵. Therefore, the private sector employees have more ability to defend themselves from the pandemic.

Our findings also revealed that smoking history and use of supplements were significantly correlated with selfefficacy (p= 0.037; p= 0.029, respectively). Non-smokers have a stronger belief in their capability to counteract the pandemic threat. Smoking can increase the likelihood of hand-to-mouth transmission of COVID-19. It can pose a significant threat to the COVID-19 spread since contaminated fingers and cigarette sticks will contact the smoker's lips ²⁶. A clinical study suggested that ACE2 may be the receptor being used by SARS-CoV-2 to gain entry into cells ²⁷.

Meanwhile, cigarette smoke could induce mucosa, the primary source of ACE2 in the lungs. Smoking also increases ACE2 in the lungs, thus enhancing the individual's susceptibility to COVID-19²⁸. This statement aligns with a study about tobacco smokers at high risk of developing severe co-infections due to impaired lung function, cross-infection, and vulnerable hygiene habits ²⁶. Furthermore, the mortality rate among smokers with COVID-19 infection is higher at 38.5% than non-smokers ²⁹.

Our study stated that people who consume nutritional supplements have significantly greater self-efficacy than those who do not. It indicates that they have a lower sense of risk associated with the pandemic threat, as they take supplements regularly. Regular diet supplementation with vitamins and micronutrients can enhance the immune system. It is a different approach to preventing the transmission of COVID-19. Sahebnasagh et al. demonstrated that specific vitamins have a vital role in innate and adaptive immune responses. Vitamins A, D, E, C and B have antioxidant and immunomodulatory properties which benefit the immune system ²⁷. A study has shown that taking probiotics, omega-3 fatty acids, multivitamins, or vitamin D supplements can reduce the risk of positive COVID-19 test results ³⁰.

According to our findings, participants' mean perceived threat and severity score was (3.28±0.86) and (8.50±2.05), respectively. Furthermore, we identified that most respondents had moderate to high levels of concern regarding the risks related to COVID-19. The majority of participants revealed that they were susceptible to COVID-19. As COVID-19 cases increase significantly in the field, public concern in Indonesia regarding the severity of the disease and population vulnerability is also growing³¹

6

No	Independent Dependent Variables										
	Variables	Perceiv	ved	Perceived S	Severity	Perceived	Threat Response Efficacy		fficacy	Self Eff	icacy
		Vulneral	oility								
		$\bar{x} \pm SD$	р	$\bar{x} \pm SD$	р	$\bar{x} \pm SD$	р	$\bar{x} \pm SD$	р	$\bar{x} \pm SD$	р
1	Sex									11	
	Male	2.76 <u>+</u> 1.09	0.070	8.38 <u>+</u> 2.04	0.045	3.30 <u>+</u> 0.91	0.675	3.49 <u>+</u> 1.03	0.045*	4.13 <u>+</u> 0.74	0.000
	Female	2.61 <u>+</u> 0.99	0.370	8.59 <u>+</u> 2.06	0.245	3.25 <u>+</u> 0.82	0.675	3.22 <u>+</u> 1.01	0.045*	4.13 <u>+</u> 0.73	0.980
2	Age										
	Adolescent	2.60±1.09		8.76 <u>+</u> 1.51		3.29 <u>+</u> 0.80		3.20 <u>+</u> 1.16		4.16 <u>+</u> 0.62	
	Adult	2.74±0.98	0.228	8.42 <u>+</u> 2.27	0.605	3.30 <u>+</u> 0.87	0.203	3.36 <u>+</u> 0.95	0.848	4.08 <u>+</u> 0.79	0.283
	Elderly	2.49±1.17	0.220	8.38 <u>+</u> 1.85	0.000	3.13 <u>+</u> 0.93	0.205	3.46 <u>+</u> 1.12	0.040	4.30 <u>+</u> 0.62	0.205
	Geriatric	3.00±1.00		9.00 <u>+</u> 1.73		3.62 <u>+</u> 0.73		3.00 <u>+</u> 1.00		4.00 <u>+</u> 1.00	
3	Region				1	-			-	1	
	Western region	2.60 <u>+</u> 1.00		8.38 <u>+</u> 2.16		3.20 <u>+</u> 0.85		3.26 <u>+</u> 0.02		0.41 <u>+</u> 0.72	
	Middle region	2.86 <u>+</u> 1.10	0.063	8.79 <u>+</u> 1.67	0.328	3.59 <u>+</u> 1.00	0.027*	3.59 <u>+</u> 1.00	0.092	0.42 <u>+</u> 0.70	0.697
	Eastern region	4.00 <u>+</u> 0.00		10.0 <u>+</u> 0.00		4.47 <u>+</u> 0.00		3.00 <u>+</u> 1.41		0.35 <u>+</u> 2.12	
4	Education	0.00.1.01		0.00.1.05	1	0.0000		2 22 - 2 22		4.00.0 74	
	Primary	3.00 <u>+</u> 1.21		9.00 <u>+</u> 1.35		3.60 <u>+</u> 0.92		3.33 <u>+</u> 0.99		4.00+0.74	
	Education	2 (4) 1 04		0.0(11.44	-	0.05+0.74		2 44+1 07		4 10 10 74	
	Middle	2.64 <u>+</u> 1.04	0.504	8.96 <u>+</u> 1.44	0.121	3.35 <u>+</u> 0.74	0.411	3.44 <u>+</u> 1.06	0.855	4.13 <u>+</u> 0.74	0.09
	Highor	2 66+1 02		8 22+2 20	-	2.21 ± 0.00		2 20+1 01		4 14+0 72	
	Education	2.00 <u>+</u> 1.02		0.23 <u>+</u> 2.29		5.21 <u>+</u> 0.90		5.29 <u>+</u> 1.01		4.14+0.75	
5	Occupation									<u> </u>	
0	Student	2 58+1 00		8 85+1 66		3 29+0 74		3 24+1 17		4 15+0 57	
	Private sector	2.73+1.09		8.52+2.12		3.31+0.94		3.52+1.03		4.29+0.74	
	employee			<u>-</u>		<u>-</u>		<u>-</u>			
	Government	2.84+1.01	0.771	7.68+2.40	0.036*	3.19+0.87	0.952	3.22+1.00	0.161	3.78+0.82	0.018*
	worker	—		—		_		—		—	
	Entrepreneur	2.53 <u>+</u> 1.05		9.19 <u>+</u> 1.23		3.31 <u>+</u> 0.75		3.63 <u>+</u> 1.16		4.28 <u>+</u> 0.77	
6	Marital Status										
	Married	2.73 <u>+</u> 1.01		8.46 <u>+</u> 2.19		3.31 <u>+</u> 0.88		3.39 <u>+</u> 0.99		4.12 <u>+</u> 0.74	
	Single	2.63 <u>+</u> 1.07	0 297	8.66 <u>+</u> 1.66	0.881	3.29 <u>+</u> 0.81	0135	3.15 <u>+</u> 1.06	0 1 3 9	4.14 <u>+</u> 0.68	0 734
	Widow/	2.27 <u>+</u> 1.10	0.277	8.66 <u>+</u> 1.66	0.001	3.29 <u>+</u> 0.81	0.100	3.53 <u>+</u> 1.12	0.135	4.27 <u>+</u> 0.80	0.754
	Widower										
7	Monthly Persona	l Income (IDR	.)		1				1		
	Low income	3.00 <u>+</u> 1.41		8.56 <u>+</u> 2.24	-	<u>3.48+1.08</u>		<u>3.11+</u> 1.36		4.33 <u>+</u> 0.71	
	Lower middle	2.70 <u>+</u> 1.04		8.52 <u>+</u> 2.05		3.31 <u>+</u> 0.90		3.50 <u>+</u> 1.03		4.09 <u>+</u> 0.64	
	Income	2 54+0.02	0.547	8 EQ 12 02	0.215	2 22 + 0 78	0.690	2 20+1 04	0.691	4 21+0 72	0.539
	income	2.34 <u>+</u> 0.92		8.39 <u>+</u> 2.03		5.22 <u>+</u> 0.78		5.50 ± 1.04		4.21 ± 0.75	
	High income	2 79+1 11		8 36+2 09		3 29+0 90		3 29+0 97		4 04+0 79	
8	Income Condition	ns		0.00-2.09		0.29 - 0.90		0.29 0.97		1.01_0.79	
	Decreased	2.67+1.01		8.62+2.01		3.30+0.84		3.31+1.04		4.16+0.74	
	income			<u>-</u>		<u>-</u>		<u>-</u>			
	Increased	3.50+0.71	0.010	9.00+1.41	0.400	3.96+0.71	0.074	4.00+0.00	0.422	4.00+0.00	0.415
	revenue	_	0.319	_	0.490	_	0.064	_	0.622	_	0.415
	No changes	2.69 <u>+</u> 1.03		8.45 <u>+</u> 1.94		3.28 <u>+</u> 0.83		3.35 <u>+</u> 1.00		4.08 <u>+</u> 0.72	
	No income	2.17 <u>+</u> 1.60		6.83 <u>+</u> 3.97		2.38 <u>+</u> 1.34		3.67 <u>+</u> 1.21		4.50+0.84	
9	Direct Cash Assis	stance									
	Yes	2.71+1.27	0.884	8.57+1.83	0.966	3.29+0.95	0.988	2.93 <u>+</u> 0.92	0.101	4.00+0.68	0.416
	No	2.67 <u>+</u> 1.02	0.004	8.50 <u>+</u> 2.07	0.700	3.28 <u>+</u> 0.86	0.700	3.36 <u>+</u> 1.03	0.101	4.14 <u>+</u> 0.73	0.110
10	Health Status				1	1			1		
	Healthy	2.62 <u>+</u> 1.02	0.028	8.47 <u>+</u> 2.09	0.955	3.23 <u>+</u> 0.85	0.050	3.37 <u>+</u> 1.03	0.269	4.17 <u>+</u> 0.74	0.029
4.5	Do not know	<u>3.19+1.08</u>		8.67 <u>+</u> 1.71		3.68 <u>+</u> 0.91		3.10 <u>+</u> 0.99		3.81 <u>+</u> 0.60	
11	Quarantine condi	itions									

Table II. Illness Risk Perceptions and Efficacy Beliefs toward COVID-19

	Full time activities at home	2.56 <u>+</u> 1.19		8.38 <u>+</u> 2.03		3.19 <u>+</u> 0.97		3.44 <u>+</u> 1.16		4.28 <u>+</u> 0.68	
	Leaving the house 2-3x per week not for work	2.65 <u>+</u> 0.87	0.052	8.86 <u>+</u> 1.71	0.559	3.35 <u>+</u> 0.68	0.087	3.16 <u>+</u> 0.93	0.464	4.16 <u>+</u> 0.67	0.567
	Work outside every day	2.94 <u>+</u> 1.13		8.44 <u>+</u> 2.14		3.43 <u>+</u> 0.95		3.39 <u>+</u> 1.03		4.12 <u>+</u> 0.79	
	Work outside 2-3x per week	2.35 <u>+</u> 0.85		8.54 <u>+</u> 1.92		3.08 <u>+</u> 0.71		3.43 <u>+</u> 1.00		4.09 <u>+</u> 0.73	
	Others	2.56 <u>+</u> 0.97		7.75 <u>+</u> 2.84		3.00 <u>+</u> 0.96		3.19 <u>+</u> 1.11		3.94 <u>+</u> 0.68	
12	History of Chron	ic Illness									
	Yes	2.55 <u>+</u> 0.82	0.770	9.00 <u>+</u> 1.55	0.470	3.35 <u>+</u> 0.74	0 595	2.91 <u>+</u> 0.83	0.204	4.18 <u>+</u> 0.75	0.825
	No	2.68 <u>+</u> 1.05	0.779	8.47 <u>+</u> 2.07	0.479	3.27 <u>+</u> 0.87	0.385	3.36 <u>+</u> 1.03	0.204	4.13 <u>+</u> 0.73	0.655
13	Smoking History	÷									
	Yes	2.89 <u>+</u> 1.05	0.114	8.45 <u>+</u> 1.82	0.402	3.43 <u>+</u> 0.87	0 1 7 2	3.47 <u>+</u> 1.04	0.259	3.94 <u>+</u> 0.73	0.027*
	No	2.62 <u>+</u> 1.03	0.114	8.51 <u>+</u> 2.11	0.405	3.23 <u>+</u> 0.85	0.175	3.31 <u>+</u> 1.02	0.358	4.18 <u>+</u> 0.72	0.037
14	Supplement Use										
	Yes	2.69 <u>+</u> 1.07	0.800	8.50 <u>+</u> 2.25	0.182	3.26 <u>+</u> 0.90	0.828	3.28 <u>+</u> 1.07	0 224	4.22 <u>+</u> 0.74	0.070*
	No	2.66 <u>+</u> 0.98	0.800	8.49 <u>+</u> 1.74	0.165	3.29 <u>+</u> 0.80	0.020	3.42 <u>+</u> 0.96	0.334	4.01 <u>+</u> 0.70	0.029

Table II reported that the mean score of perceived severity of respondents was male (8.38+2.04) and female (8.59+2.06). This high score indicated that the perceived severity of COVID-19 among males and females was severe and fatal. The general population's severity perception in Indonesia is higher than in the Myanmar-based study³². Similar results were found in a study in Hongkong, where all participants agreed that the COVID-19 disease was very severe ³³. Regarding the pandemic, the internet and other information sources can better influence people's thinking in applying protective measures ³⁴. A study reported that respondents in Indonesia had taken more protective behavior. People who often get information related to COVID will have higher self-efficacy beliefs 35. Mya et al. have reported that individuals would engage in more protective behavior due to the easy access to mass media and social media³².

A person perceiving the high risk of COVID-19 is likely to feel stress, panic, depression, and try to adapt to others' behavior ³⁶. It is because strong negative emotions could encourage one to think about protective behavior in the face of this pandemic³⁷. Nevertheless, the higher threat perceived by vulnerable groups may increase their self-protective behavior, which is beneficial in pandemic control. However, those with a low-risk perception of COVID-19 are less likely to engage in protective behavior. Thus, public health education is targeted for this group ³⁸.

Understanding risk perception is a complex phenomenon created from various psychological, social and cultural factors in different places and times ³⁹. This phenomenon can be interpreted as a form of pandemic preparedness. Based on previous studies, risk perception can assess and evaluate an individual's response to a pandemic ⁴⁰. Though perceived risk acts as a trigger for preventive actions, it is also determined by a person's social networks, community beliefs and the source of information about the health behavior 41,42. Social networks may amplify the spread of beneficial or dangerous behavior during this COVID-19 epidemi43. As a non-medical measure, personal protective practices are needed control the COVID-19 pandemic to by implementing health protocols, wearing masks, avoiding crowds and maintaining social distancing44. The community's willingness could play a vital role in successfully implementing government policies 45.

CONCLUSION

We concluded a moderate to a high level of risk perceptions associated with COVID-19 in Indonesia's general population. Additionally, they had a relatively good efficacy response in adopting self-protection measures in the COVID-19 pandemic. The public's risk perception of a pandemic contribute to increasing public participation in preventing the COVID-19 pandemic. Furthermore, these findings will contribute to the health authorities regarding COVID-19 pandemic risk communication management.

ACKNOWLEDGMENT

We are grateful to all participants who gave their time to this research.

AUTHORS' CONTRIBUTION

Lolita,Lolita : study design, methodology, data collection, validation, and writing – original draft. Azis Ikhsanudin: data management, data collection, visualization, statistical analysis, editing.

FUNDING DETAILS

The authors received no financial support for this study, authorship, and publication.

DATA AVAILABILITY

None

CONFLICT OF INTEREST:

The authors declare no conflict of interest.

REFERENCES

- Kunjana Rahardi R. Building Critical Awareness of Corona Virus-Related News: Cyber-Pragmatic Study of COVID-19 Hoaxes on Social Media. Int J Adv Sci Technol. 2020;29(6):5398–409.
- de Groot RJ, Baker SC, Baric RS, Brown CS, Drosten C, Enjuanes L, et al. Middle East respiratory syndrome coronavirus (MERS-CoV): announcement of the Coronavirus Study Group. J Virol. 2013/05/15. 2013 Jul;87(14):7790–2.
- 3. Pradana M, Rubiyanti N, S W, Hasbi I, Utami DG. Indonesia's fight against COVID-19: the roles of local government units and community organisations. Local Environ. 2020;25(9):741–3.
- 4. WHO Indonesia, "Coronavirus Disease 2019 (COVID-19) Situation Report-18. World Heal. Organ. 2020. p. 1–20.
- Vahia I V, Blazer DG, Smith GS, Karp JF, Steffens DC, Forester BP, et al. COVID-19, Mental Health and Aging: A Need for New Knowledge to Bridge Science and Service. Vol. 28, The American journal of geriatric

psychiatry : official journal of the American Association for Geriatric Psychiatry. 2020. p. 695–7.

- 6. Petzold MB, Bendau A, Plag J, Pyrkosch L, Mascarell Maricic L, Betzler F, et al. Risk, resilience, psychological distress, and anxiety at the beginning of the COVID-19 pandemic in Germany. Brain Behav. 2020 Sep;10(9):e01745.
- Djalante R, Lassa J, Setiamarga D, Sudjatma A, Indrawan M, Haryanto B, et al. Review and analysis of current responses to COVID-19 in Indonesia: Period of January to March 2020. Prog Disaster Sci. 2020;6.
- Xiong J, Lipsitz O, Nasri F, Lui LMW, Gill H, Phan L, et al. Impact of COVID-19 pandemic on mental health in the general population: A systematic review. J Affect Disord. 2020 Dec;277:55–64.
- 9. Giallonardo V, Sampogna G, Del Vecchio V, Luciano M, Albert U, Carmassi C, et al. The Impact of Quarantine and Physical Distancing Following COVID-19 on Mental Health: Study Protocol of a Multicentric Italian Population Trial. Front psychiatry. 2020 Jun;11:533.
- 10. Ferrer R, Klein WM. Risk perceptions and health behavior. Curr Opin Psychol. 2015 Oct;5:85–9.
- 11. Rimal RN, Real K. Perceived Risk and Efficacy Beliefs as Motivators of Change. Hum Commun Res. 2003 Jul;29(3):370–99.
- Samadipour E, Ghardashi F. Factors Influencing Iranians' Risk Perception of Covid-19 TT - از ایرانیان خطر درک بر موثر عوامل 19. Journal-Mil-Med. 2020 Apr;22(2):122-9.
- He S, Chen S, Kong L, Liu W. Analysis of Risk Perceptions and Related Factors Concerning COVID-19 Epidemic in Chongqing, China. J Community Health. 2021 Apr;46(2):278–85.
- 14. MCCAFFREY S. Thinking of Wildfire as a

Natural Hazard. Soc Nat Resour. 2004 Jul;17(6):509–16.

- Nanda RO, Lolita L, Indayati W, Rusdiyanti I, Ikhsanudin A, Mareti S. Knowledge, precautionary actions, and perceived risk of COVID-19 among indonesian people. Int J Public Heal Sci. 2021;10(1):8–15.
- Nanda RO, Lolita L, Indayati W, Rusdiyanti I, Nurjannah, Ikhsanudin A, et al. Covid-19 risk perception among Indonesians in early stage of the outbreak. Int J Public Heal Sci. 2021;10(2):249–57.
- Aulawi A. Penerbitan Peraturan Pemerintah Pengganti Undang-Undang Nomor 1 Tahun 2020 Sebagai Strategi Kebijakan Pajak Pemerintah Indonesia Dalam Menghadapi Dampak Pandemi Covid-19 Terhadap Keuangan Negara. Prog J Pendidikan, Akunt dan Keuang. 2020;3(2):110–32.
- 18. Kemp S. DIGITAL 2020: INDONESIA. Data reportal. 2020.
- De Zwart O, Veldhuijzen IK, Elam G, Aro AR, Abraham T, Bishop GD, et al. Perceived threat, risk perception, and efficacy beliefs related to SARS and other (emerging) infectious diseases: Results of an international survey. Int J Behav Med. 2009;16(1):30–40.
- 20. Umamaheswar J, Tan C. "Dad, Wash Your Hands": Gender, Care Work, and Attitudes toward Risk during the COVID-19 Pandemic. Socius. 2020;6.
- 21. Griffith DM, Sharma G, Holliday CS, Enyia OK, Valliere M, Semlow AR, et al. Men and COVID-19: A Biopsychosocial Approach to Understanding Sex Differences in Mortality and Recommendations for Practice and Policy Interventions. Prev Chronic Dis. 2020 Jul;17:E63.
- WHO Indonesia, "Coronavirus Disease 2019 (COVID-19) Situation Report-41,. World Heal. Organ., vol. 19, no. May. 2020. p. 1–20.

- 23. Koh D. Occupational risks for COVID-19 infection. Vol. 70, Occupational medicine (Oxford, England). 2020. p. 3–5.
- 24. Walsh B, Redmond P, Roantree B, Redmond P. Differences in Risk of Severe Outcomes From Covid-19 Across Occupations in Differences in Risk of Severe Outcomes From Covid-19 Across Occupations in. 2020;(93):1–21.
- 25. Lan F-Y, Wei C-F, Hsu Y-T, Christiani DC, Kales SN. Work-related COVID-19 transmission in six Asian countries/areas: A follow-up study. PLoS One. 2020 May;15(5):e0233588.
- 26. Ahmed N, Maqsood A, Abduljabbar T, Vohra F. Tobacco Smoking a Potential Risk Factor in Transmission of COVID-19 Infection. Pakistan J Med Sci. 2020 May;36(COVID19-S4):S104–7.
- Sahebnasagh A, Saghafi F, Avan R, Khoshi A, Khataminia M, Safdari M, et al. The prophylaxis and treatment potential of supplements for COVID-19. Eur J Pharmacol. 2020 Nov;887:173530.
- 28. Polverino F. Cigarette Smoking and COVID-19: A Complex Interaction. Am J Respir Crit Care Med. 2020 Jun;202(3):471– 2.
- 29. Alqahtani JS, Oyelade T, Aldhahir AM, Alghamdi SM, Almehmadi M, Alqahtani AS, et al. Prevalence, Severity and Mortality associated with COPD and Smoking in patients with COVID-19: A Rapid Systematic Review and Meta-Analysis. PLoS One. 2020;15(5):e0233147.
- Mrityunjaya M, Pavithra V, Neelam R, Janhavi P, Halami PM, Ravindra P V. Immune-Boosting, Antioxidant and Antiinflammatory Food Supplements Targeting Pathogenesis of COVID-19. Front Immunol. 2020;11:570122.
- 31. WHO. "Media Statement Knowing the risks for COVID-19," Media Statement: Knowing the risks for COVID-19.

https://www.who.int/indonesia/news/d etail/08-03-2020-knowing-the-risk-forcovid-19#. 2020.

- 32. Mya KS, S. M. A, Hlaing WA, Hlaing SS, Aung T, Lwin SMM, et al. Awareness, perceived risk and protective behaviours of Myanmar adults on COVID-19. Int J Community Med Public Heal. 2020;7(5):1627.
- Kwok KO, Li KK, Chan HHH, Yi YY, Tang A, Wei WI, et al. Community Responses during Early Phase of COVID-19 Epidemic, Hong Kong. Emerg Infect Dis. 2020;26(7):1575–9.
- 34. Yıldırım M, Güler A. COVID-19 severity, self-efficacy, knowledge, preventive behaviors, and mental health in Turkey. Death Stud. 2020 Jul;1–8.
- 35. Xiong H, Yi S, Lin Y. The Psychological Status and Self-Efficacy of Nurses During COVID-19 Outbreak: A Cross-Sectional Survey. Inquiry. 2020;57:46958020957114.
- Kuang J, Ashraf S, Das U, Bicchieri C. Awareness, Risk Perception, and Stress during the COVID-19 Pandemic in Communities of Tamil Nadu, India. Int J Environ Res Public Health. 2020 Sep;17(19).
- 37. Wise T, Zbozinek TD, Michelini G, Hagan CC, Mobbs D. Changes in risk perception and self-reported protective behaviour during the first week of the COVID-19 pandemic in the United States. R Soc open Sci. 2020 Sep;7(9):200742.
- 38. Abdelrahman M. Personality Traits, Risk Perception, and Protective Behaviors of Arab Residents of Qatar During the COVID-19 Pandemic. Int J Ment Health Addict. 2022;20(1):237–48.
- Lohiniva A-L, Sane J, Sibenberg K, Puumalainen T, Salminen M. Understanding coronavirus disease (COVID-19) risk perceptions among the public to enhance risk communication efforts: a practical approach for outbreaks,

Finland, February 2020. Euro Surveill Bull Eur sur les Mal Transm = Eur Commun Dis Bull. 2020 Apr;25(13).

- 40. Cori L, Bianchi F, Cadum E, Anthonj C. Risk Perception and COVID-19. Vol. 17, International journal of environmental research and public health. 2020.
- 41. Chan EYY, Huang Z, Lo ESK, Hung KKC, Wong ELY, Wong SYS. Sociodemographic Predictors of Health Risk Perception, Attitude and Behavior Practices Associated with Health-Emergency Disaster Risk Management for Biological Hazards: The Case of COVID-19 Pandemic in Hong Kong, SAR China. Int J Environ Res Public Health. 2020 May;17(11).
- 42. Lin C-Y, Imani V, Majd NR, Ghasemi Z, Griffiths MD, Hamilton K, et al. Using an integrated social cognition model to predict COVID-19 preventive behaviours. Br J Health Psychol. 2020 Nov;25(4):981–1005.
- 43. Abrams EM, Greenhawt M. Risk Communication During COVID-19. J allergy Clin Immunol Pract. 2020 Jun;8(6):1791–4.
- 44. Khosravi M. Perceived risk of COVID-19 pandemic: The role of public worry and trust. Electron J Gen Med. 2020;17(4):1–2.
- 45. Serwaa D, Lamptey E, Appiah AB, Senkyire EK, Ameyaw JK. Knowledge, risk perception and preparedness towards coronavirus disease-2019 (COVID-19) outbreak among Ghanaians: a quick online cross-sectional survey. Pan Afr Med J. 2020;35(Suppl 2):44.