

LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH: JURNAL ILMIAH*

Judul karya ilmiah (artikel) : Black Cumin Seed Oil Increases Phagocytic Activity and Secretion Of IL-12 by Macropages
 Jumlah Penulis : 2 Orang
 Nama Penulis : Akrom, Mustofa
 Status Pengusul : Penulis pertama/penulis ke.../penulis korespondensi **
 Identitas Jurnal Ilmiah : a. Nama Jurnal : Biomedical Research
 b. No ISSN : 5241-5246
 c. Vol, No, Bulan, Tahun : Vol. 28, No.2, tahun 2017
 d. Penerbit : Biomedical Research
 e. DOI artikel (jika ada) :
 f. Alamat web jurnal : www.biomedres.info
 g. Terindeks Scimagojr/Thomson Reuter ISI Knowledge atau di...**

Kategori Publikasi Jurnal Ilmiah : Jurnal Ilmiah Internasional/Int. Bereputasi**
 (beri V pada kategori yang tepat) Jurnal Ilmiah Nasional Terakreditasi
 Hasil Penilaian Peer Review : Jurnal Ilmiah Nasional/ Nasional Terindeks di DOAJ, CABI, COPERNICUS**

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi	Internasional	Nasional Terakreditasi	Nasional Tidak Terakreditasi	Nasional Terindeks DOAJ dll	
a. Kelengkapan unsur isi artikel (10%)	4					4
b. Ruang lingkup dan kedalaman pembahasan (30%)	12					10
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)	12					10
d. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)	12					12
Total = (100%)	40					36
Nilai Pengusul	24					21,6

<p>Komentar Peer Review</p>	<p>1. Tentang kelengkapan dan kesesuaian unsur :</p> <p><i>baik</i></p> <p>2. Tentang ruang lingkup & kedalaman pembahasan :</p> <p><i>Akademik hanya minimalis</i></p> <p>3. Kecukupan dan kemutakhiran data serta metodologi :</p> <p><i>Sudah banyak penelitian sejenis dg bahan berbeda, selaman tidak signifikan</i></p> <p>4. Kelengkapan unsur kualitas penerbit :</p> <p><i>baik</i></p> <p>5. Indikasi plagiasi :</p> <p><i>Tidak ada</i></p> <p>6. Kesesuaian bidang ilmu :</p> <p><i>Seni</i></p>
-----------------------------	--

2017

Reviewer *1/2*

Nama *Prof. Sugianto*
 NIP/NIY. _____
 Bidang Ilmu *Farmakol & Toksikol*
 Jabatan Akademik *Guru Besar*
 Unit Kerja *Farmasi UAS*

*dinilai oleh dua Reviewer secara terpisah
 ** coret yang tidak perlu
 *** nasional/ terindeks di DOAJ, CABI, Copernicus

**LEMBAR HASIL PENILAIAN SEJAWAT SEBIDANG ATAU PEER REVIEW
KARYA ILMIAH: JURNAL ILMIAH***

Judul karya ilmiah (artikel) : Black Cumin Seed Oil Increases Phagocytic Activity and Secretion Of IL-12 by Macropages
 Jumlah Penulis : 2 Orang
 Nama Penulis : Akrom, Mustofa
 Status Pengusul : Penulis pertama/penulis ke.../penulis korespondensi **
 Identitas Jurnal Ilmiah : a. Nama Jurnal : Biomedical Research
 b. No ISSN : 5241-5246
 c. Vol, No, Bulan, Tahun : Vol. 28, No.2, tahun 2017
 d. Penerbit : Biomedical Research
 e. DOI artikel (jika ada) :
 f. Alamat web jurnal : www.biomedres.info
 g. Terindeks Scimagojr/Thomson Reuter ISI Knowledge atau di...**

Kategori Publikasi Jurnal Ilmiah :
(beri v pada kategori yang tepat)

v

Jurnal Ilmiah Internasional/Int. Bereputasi**
 Jurnal Ilmiah Nasional Terakreditasi
 Jurnal Ilmiah Nasional/ Nasional Terindeks di DOAJ, CABI, COPERNICUS**

Hasil Penilaian Peer Review :

Komponen yang dinilai	Nilai Maksimal Jurnal Ilmiah (isikan di kolom yang sesuai)					Nilai Akhir Yang Diperoleh
	Internasional Bereputasi	Internasional	Nasional Terakreditasi	Nasional Tidak Terakreditasi	Nasional Terindeks DOAJ dll	
a. Kelengkapan unsur isi artikel (10%)		3				3
b. Ruang lingkup dan kedalaman pembahasan (30%)		9				8,8
c. Kecukupan dan kemutakhiran data/informasi dan metodologi (30%)		9				8,9
d. Kelengkapan unsur dan kualitas terbitan/jurnal (30%)		9				9
Total = (100%)		30				29,7
Nilai Pengusul		18				17,82

<p>Komentar Peer Review</p>	<p>1. Tentang kelengkapan dan kesesuaian unsur :</p> <p style="text-align: center;">Jesuis</p> <p>2. Tentang ruang lingkup & kedalaman pembahasan :</p> <p style="text-align: center;">Baik</p> <p>3. Kecukupan dan kemitakhiran data serta metodologi :</p> <p style="text-align: center;">Baik</p> <p>4. Kelengkapan unsur kualitas penerbit :</p> <p style="text-align: center;">Jesuis</p> <p>5. Indikasi plagiasi :</p> <p style="text-align: center;">Tidak ada</p> <p>6. Kesesuaian bidang ilmu :</p> <p style="text-align: center;">Jesuis</p>
-----------------------------	--

2017

Reviewer 1/2 *

Nama Prof. Dr. Sunjaya Kusno
 NIP/NIY. 1952.11.20.1977031004
 Bidang Ilmu Fisiologi Fisiologi
 Jabatan Akademik Guru Besar
 Unit Kerja Fakultas Farmasi UBT

*dinilai oleh dua Reviewer secara terpisah

** coret yang tidak perlu

*** nasional/ terindeks di DOAJ, CABI, Copernicus

Black cumin seed oil increases phagocytic activity and secretion of IL-12 by macrophages.

By Akrom Akrom

1

See discussions, stats, and author profiles for this publication at: <https://www.researchgate.net/publication/318835885>

Black cumin seed oil increases phagocytic activity and secretion of IL-12 by macrophages.

Article · January 2017

CITATIONS

0

READS

12

2 authors, including:



Akrom Akrom

Ahmad Dahlan University

29 PUBLICATIONS 7 CITATIONS

SEE PROFILE

Some of the authors of this publication are also working on these related projects:



Development of Nigella sativa oil as immunomodulator and antioxidant for metabolic syndroms patients [View project](#)



Nigella sativa as chemopreventive and immunomodulator [View project](#)

All content following this page was uploaded by Akrom Akrom on 16 August 2017.

The user has requested enhancement of the downloaded file.

Black cumin seed oil increases phagocytic activity and secretion of IL-12 by macrophages.

Akrom^{1,2*}, Mustofa³

¹Department of Pharmacology and Clinical Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia

²Drug information and Crisis Center, Ahmad Dahlan University, Indonesia

³Department of Pharmacology, Faculty of Medicine, Gadjah Madah University, Yogyakarta, Indonesia

Abstract

Interleukin-12 (IL-12) is one of cytokines that regulate the innate and adaptive immune responses produced by macrophages. This study was to investigate the effect of Black Cumin Seed Oil (BCSO) to the structure of peripheral blood, phagocytic activity, and secretion of IL-12 by macrophage peritoneal Sprague-Dawley (SD) rat. The 36 female Sprague Dawley (SD) rats divided into 6 groups. Normal group was only given standards food and drink. Treatments group were given BCSO 0.25, 2.5 and 5 ml/kg BW/day for 14 d. Positive control group was given thymoquinone 50 mg/kg BW/day for 14 d. Solvent control group was given DMSO. Blood test, phagocytic activity and secretion of IL-12 by macrophage *in vitro* performed on the 15th day. Latex and NBT assay method were conducted to measure the phagocytic activity of macrophages. The expression level of IL-12 and TLR4 were analysed by ELISA. Data were statistically tested by one-way ANOVA with significance 95%. BCSO was increase phagocytic activity, secretion of IL-12 by macrophage, and the expression of TLR4. The highest phagocytosis percent ($56.83 \pm 6.37\%$) and phagocytosis index (5.18 ± 0.39) was performed by 0.25 BCSO group and did not significantly different with thymoquinone group. The highest level of IL-12 was in 2.5 BCSO group ($66.33 \pm 2.11 \mu\text{M}$).

Keywords: Black cumin seed oil, IL-12, Macrophage, Phagocytosis, Thymoquinone, TLR-4.

Accepted on April 10, 2017

Introduction

Interleukin-12 (IL-12) is one of the main cytokines in immune response regulation. It is produced by activated lymphocytes and accessory cells such as macrophage, dendritic cell, neutrophil, and monocyte. *In vivo* study proved that IL-12 increase the activity of CD4 and CD8 [1,2], lead the formation of CD4 T cell, induce the secretion of IFN γ by T and NK cell, anti-angiogenesis, and DNA repair [3,4]. The production of IL-12 by macrophage is increased by immunogen such as phytoimmunogen [5].

In Indonesia, black cumin seed is widely used as herbal medicine with the efficacy as a tonic, anti-cancer, anti-pain, and anti-asthma and reinforcing the body's defenses and antioxidant, but many people use the BCSO to maintain of fitness and health status. BCSO as guardians of fitness and health status generally is consumed every day in the time varying between 1-2 weeks with $2-3 \times 1.5-2$ ml/day. Black cumin seeds contain oil evaporate, fatty acids, sterols mainly β -sitosterol, thymoquinone, dithymoquinone and saponin [6]. Black Cumin Seed Oil (BCSO) inhibits inflammation of

bronchi and decrease asthma by inhibiting mRNA expression of IL-4, IL-5, IL-6 and TGF- β in the allergens-induced experimental animals [7]. Thymoquinone, as the main active compound of BCSO, decreases inflammatory reactions in mice bronchi, IgE and IgG specific-OVA, IL-5, IL-4, IL-13 and increases IFN- γ in ovalbumin-induced mice [8,9]. Thymoquinone as a benzoquinone compound has a chemical structure identical to the active compounds as anti-inflammatory, so it is estimated that thymoquinone also has anti-inflammatory activity. However, the results of Finley et al. study [10,11] showed that through the Toll-Like Receptor 4 (TLR-4) thymoquinone is able to increase the activity of macrophages. The immunomodulatory effects of daily consumption of BCSO in 1-2 weeks in healthy people have not been clear, particularly against components cells of blood and macrophage activity.

In this study, we want to clarify the activity of BCSO as an immunomodulator by analyzing the phagocytic activity and the secretion of IL-12 by the macrophage.

Method

BCSO treatment

Black cumin seed oil has been prepared by the department of Phytochemistry and Pharmacognosy, Faculty of Pharmacy, University of Ahmad Dahlan. Thymoquinone levels in oil has been established which is 2.7% volume/volume (=21.6 mg thymoquinone per ml BCSO). Thirty six one-month-old female SD rats were divided into 6 groups randomly. Group I as a normal group was only given standards food and drink. Groups II-IV as the treatment group were given BCSO 0.25, 2.5 and 5 ml/kg BW/day (similar to 5.4 mg, 54 mg and 108 mg/kgbw thymoquinone) for 14 d. Group V as the positive control group was given thymoquinone 50 mg/kg BW/day. Group VI as a solvent control group was given DMSO7. Research procedures and the research protocol had been reviewed and approved by the ethics committee of the Gadjah Mada University (no: 222/KEC-LPPT/III/2015).

Blood test

A peripheral blood test conducted in the laboratory of Gadjah Mada University (GMU). It was done by hematology analyser Sysmeix kx. The blood of rats was taken through orbital sinus as much as ± 1.5 ml and collected in labeled eppendorf tubes that were already containing anticoagulants [12]. As suggested by the ethics committee, the blood sampling and isolation of macrophage cells is conducted under the animals test were decapitated.

Macrophage cell preparation

On day 15, all of the rats were euthanized using chloroform. They were sprayed with 70% ethanol and mounted on the styrofoam block on its back. The abdomen skin was opened and 10 ml cold Roswell Park Memorial Institute medium (RPMI) 1640 (Sigma-Aldrich) was injected into the peritoneal cavity using a 27 g needle. The peritoneum was gently massaged to dislodge any attached cells into the RPMI solution. The fluid was collected by inserting a 25 g needle into the peritoneal cavity and saved on the tube. It was centrifuged at 1200 rpm, 4°C for 10 min. Supernatant was discarded while pellet cell was resuspended by RPMI contained Fetal Bovine Serum (FBS) 10% (Sigma-Aldrich). The number of the cell was counted by hemocytometer. 5×10^5 cell/ml macrophage cells were cultured in 24 well microculture with a coverslip and incubated in CO₂ incubator 37°C for 30 min. Each well was added by 1 ml complete medium and phytoimmunogen, Phytohaemagglutinin (PHA) and incubated 2 h. Cells then were washed twice with 1 mL RPMI and incubated in complete medium for 24 h [13].

Macrophage phagocytosis activity test

Macrophage cells were washed twice with RPMI then added by 200 μ L of 2.5×10^6 /ml latex suspension/well. It was incubated in CO₂ incubator 5%, 37°C for 60 min. Cells were washed three times with PBS, air dried, fixed with absolute methanol for 30 s and stained with 20% Giemsa. The percentage of phagocytic activity and the phagocytic index was measured by light microscope with 400X magnification.

ROI and NO secretion activity test

The ability of macrophages cells in secreting ROI was measured by NBT reduction assay [13,14]. Determination of nitrite (NO₂) in the supernatant of macrophage culture was performed by Grees assay [15].

IL-12p40 secretion activity test and expression of TLR4

The level of IL-12p40 from the supernatant of splenocytes cell culture was measured by Enzyme-Linked Immunosorbent Assay (ELISA) sandwich. The kit used was rat IL-12 (IL-12p40, Biosource, USA, catalog KRC 4022). The expression of TLR-4 was also determined by ELISA according to Finley [14].

Statistical analysis

The number of blood cells, phagocytic activity, phagocytic index, ROI secretion activity, the levels of NO and IL12 and TLR-4 expression were analysed by normality test, homogeneity and mean difference with 95% confidence interval. Normality test was used Kolmogorov-Smirnov test and homogeneity test was used Levene test then followed by LSD.

Result and Discussion

The overview of peripheral blood cell

The blood tests showed that administration of BCSO for 14 d did not affect the number of erythrocytes, platelets, and hemoglobin and hematocrit levels, as shown in Table 1. The level of hemoglobin, hematocrit, MCV, MCH, MCHC, and the number of erythrocytes and platelet with the oral administration of BCSO were still in normal values. However, the lowest of platelet number was in 5 ml/kg BW BCSO and significantly different from the normal and the solvent group ($p < 0.05$). The condition of the thirty-six SD rats were used for this study until the sampling are all in good health.

Table 1. The number (\pm Sd) of erythrocytes, hemoglobin, hematocrit, MCV, MCH, MCHC and platelet of SD rats after 14 d BCSO treatment.

Group	RBC (10^9 /ml) (mean \pm Sd)	Hb (g/dl) (mean \pm Sd)	Hematocrit (mean \pm Sd)	(%) MCV(μ m ³) (mean \pm Sd)	MCH(pg) (mean \pm Sd)	MCHC(g/dl) (mean \pm Sd)	Platelet (10^3 /ml) (mean \pm Sd)
-------	--------------------------------------	------------------------------	-------------------------------	--	----------------------------	-------------------------------	---

Black cumin seed oil increases phagocytic activity and secretion of IL-12 by macrophages

Normal	6.6 ± 0.04	13.0 ± 0.51	37.8 ± 1.78	56.9 ± 2.52	19.6 ± 0.76	34.5 ± 0.27	1011 ± 106
BCSO 0.25 ml/kgBW	6.5 ± 0.05	12.7 ± 0.15	36.7 ± 0.99	55.8 ± 1.14	19.4 ± 0.15	34.7 ± 0.67	801 ± 139
BCSO 2.5 ml/kgBW	6.6 ± 0.53	11.5 ± 2.38	37.0 ± 3.36	57.3 ± 1.44	17.8 ± 3.41	30.9 ± 4.73	850 ± 121
BCSO 5 ml/kgBW	6.9 ± 0.43	13.2 ± 1.11	37.7 ± 2.35	55.8 ± 1.85	20.0 ± 0.70	35.8 ± 0.25	640 ± 33
Thymoquinone	6.5 ± 0.40	13.1 ± 1.29	36.7 ± 3.35	56.5 ± 2.65	20.2 ± 0.98	35.8 ± 0.38	878 ± 40
Solvent	6.5 ± 0.26	11.9 ± 0.39	35.6 ± 1.23	54.8 ± 0.83	18.4 ± 0.47	33.5 ± 0.33	837 ± 91

Note: BCSO: Black Cumin Seed Oil.

The results of the number of leukocytes measurement (thousand/mm³) after oral administration of BCSO for 14 d were presented in Table 2. BCSO did not affect the number of leukocytes but affects leukocyte counts especially for a dose of

5 ml/kg BW. 5 ml/kg BW decreased the number of neutrophils and increased the number of lymphocytes compared to the normal group (p<0.05). It did not affect the number of monocytes, eosinophil, and basophil.

Table 2. The number (± Sd) of total leukocyte and composition of leukocyte counts of SD rats after 14 d treatment.

Group	Leukocyte (10 ³ /ml)	Neutrophil (%)	Lymphocyte (%)	Monocyte (%)	Eosinophil (%)	Basophil (%)
Normal	6.3 ± 0.70	28.0 ± 2.00	65.0 ± 1.00	5.7 ± 1.15	1.3 ± 0.58	0.0
BCSO 0.25 ml/kgBW	5.7 ± 0.96	27.3 ± 2.08	64.3 ± 0.58	5.3 ± 0.58	2.7 ± 2.08	0.0
BCSO 2.5 ml/kgBW	7.8 ± 0.83	35.7 ± 3.06	58.0 ± 2.00	6.7 ± 1.15	1.7 ± 1.15	0.0
BCSO 5 ml/kgBW	5.8 ± 1.05	23.7 ± 1.53a	71.0 ± 5.29a	6.0 ± 1.00	2.7 ± 2.08	0.0
Thymoquinone	7.4 ± 0.95	31.7 ± 3.76	61.0 ± 3.61	6.3 ± 0.58	1.3 ± 0.58	0.0
Solvent	7.6 ± 1.06	30.3 ± 3.95	62.3 ± 3.50	6.3 ± 0.50	1.5 ± 0.58	0.0

Note: ^asignificantly different (p<0.05) from normal group; BCSO: Black Cumin Seed Oil.

The number of macrophage cell, phagocytic activity, secretion of ROI and NO

As shown in Table 3, BCSO and thymoquinone influence the number of macrophage cell, phagocytic activity and phagocytic index of macrophage culture cells. The number of macrophage cell, phagocytic activity and phagocytic index in BCSO and thymoquinone group were significantly increase compared to

normal and solvent group. BCSO and thymoquinone were also increase the ROI secretion activity (Table 4). ROI secretion activity of macrophages culture cells from SD rat peritoneal which have been exposed by 0.25, 2.5 and 5 ml/kg BW BCSO and 50 mg/kg thymoquinone higher than control (p<0.05). The increasing of NO secretion activity only occurred in 5 ml/kg BW BCSO.

Table 3. The number of peritoneal macrophage, phagocytic activity and phagocytic index of peritoneal macrophage of SD rats after 14 d BCSO treatment.

Group	Macrophage (1 × 10 ⁶ cell/ml) (mean ± Sd)	Phagocytic activity (%) (mean ± Sd)	Phagocytic index (mean ± Sd)
Normal	1.65 ± 0.20	49.67 ± 1.97	2.99 ± 0.54
BCSO 0.25 ml/kgBW	2.56 ± 0.68 ^a	56.83 ± 6.37 ^a	5.18 ± 0.39 ^a
BCSO 2.5 ml/kgBW	2.57 ± 0.50 ^a	52.50 ± 5.24 ^a	4.92 ± 0.61 ^a
BCSO 5 ml/kgBW	2.02 ± 0.35 ^a	56.17 ± 6.52 ^a	4.35 ± 0.43 ^a
Thymoquinone	2.18 ± 0.47 ^a	55.83 ± 4.88 ^a	5.55 ± 0.58 ^a
Solvent	1.53 ± 0.13	41.33 ± 1.51	2.97 ± 0.14

Note: ^asignificantly different (p<0.05) from normal group; BCSO: Black Cumin Seed Oil

Table 4. The activity of ROI secretion (%) and NO levels in supernatant of peritoneal macrophage cells culture of SD rats after 14 d BCSO treatment.

Group	ROI secretion activity (%) (mean± Sd)	NO level (µM) (mean± Sd)
Normal	31.33 ± 5.32	2.86 ± 0.55
BCSO 0.25 ml/kgBW	52.33 ± 1.63 ^{ab}	3.54 ± 0.39
BCSO 2.5 ml/kgBW	52.83 ± 5.88 ^{ab}	3.58 ± 0.34
BCSO 5 ml/kgBW	65.17 ± 14.19 ^{ab}	4.17 ± 0.36
Thymoquinone	44.67 ± 7.55	3.85 ± 0.12
Solvent	28.00 ± 1.55	2.86 ± 0.82

Note: ^asignificantly different (p<0.05) from normal group; ^bsignificantly different (p<0.05) from thymoquinone group; BCSO: Black Cumin Seed Oil

Secretion of IL-12p40 and expression of TLR-4

BCSO or thymoquinone exposure for 14 d increased the activity of macrophage cells in IL-12 secretion and TLR-4

expression. Level of IL-12 the increasing of IL-12 secreted by macrophages and the expression of TLR-4 in BCSO groups did not differ with thymoquinone group (Table 5).

Table 5. The levels of IL-12 and the ratio of TLR-4 expression in peritoneal macrophage cells of SD rats after 14 d BCSO treatment.

Group	IL-12 levels (µL/ml) (mean ± Sd)	The ratio of TLR-4 expression (×10 ²) (mean ± Sd)
Normal	18.83 ± 1.37	1.00 ± 0.00
BCSO 0.25 ml/kgBW	58.33 ± 7.95 ^a	1.22 ± 0.17 ^a
BCSO 25 ml/kgBW	66.33 ± 2.11 ^a	1.22 ± 0.17 ^a
BCSO 5 ml/kgBW	51.17 ± 2.25 ^a	1.28 ± 0.25 ^a
Thymoquinone (50 mg/kgBW)	52.83 ± 10.49 ^a	1.33 ± 0.21 ^a
Solvent	18.75 ± 1.25	1.00 ± 0.00

Note: ^asignificantly different (p<0.05) from normal group; BCSO: Black Cumin Seed Oil

Discussion

Macrophages are professional phagocytes that act as APC and the main effectors in cellular innate and adaptive immune response [16]. This study proved that administration of BCSO and thymoquinone increased the activity of SD rat peritoneal macrophages. The number of peritoneal macrophages, phagocytic activity, macrophage phagocytosis index, secretory activity of ROI, NO and IL-12 in BCSO and thymoquinone groups higher than normal and solvent groups. One of the cytokines produced by activated macrophages is IL-12 [16]. The secretion of IL-12 by macrophages can be stimulated by the presence of parasitic intracellular microorganisms, endotoxins, and glycosides of the medicinal plant such as skrosafisida A of *Pikroriza skrofulariflora*. Secretion of IL-12 is inhibited by the presence of anti-inflammatory agents, immunosuppressants including anti-inflammatory cytokines, such as IL-10 or TGF-β [17].

Macrophages have TLR4 receptor which serves to identify the presence of natural antigen from herbal or pathogen. Black

cumin (*Nigella sativa* L.) contain evaporated oil, fatty acids, rich in sterols especially β-sitosterol, thymoquinone, dithymoquinone and saponins [18] that can activate macrophages through TLRs, especially TLR-4. Thymoquinone proved increases the activity of TLR-4 [19,20].

IL-12 cytokine is produced by APC, has various functions, and plays an important role in enhancing immune response mediated by CD4Th1 cells [21]. IL-12 is a regulatory cytokine that mediates a natural immune response by the adaptive immune response. The main roles of IL-12 in regulating the immune response are: (i). activate macrophages and dendritic cells [22], (ii). Activate CD4Th0 cell and increase the proliferation and differentiation of CD4Th0 into CD4Th1, (iii). Increase the secretion of IFN-γ by T cells, macrophages and NK cells, (iv). Increase the proliferation and activity of macrophage and NK cells, (v). Increase the activity of CTL, and (vi). Activate B lymphocytes to produce antibodies. The biological effects of IL-12 are regulating the proliferation and differentiation of lymphoid, regulate the function of

macrophages and dendritic cells, regulate the tolerance, memory and lymphocyte homeostasis [21,23].

Conclusion

BCSO treatment (0.25, 2.5 and 5 ml/kg BW) for 14 d in SD rats did not affect the number of erythrocytes, hemoglobin concentration, hematocrit, MCV, MCH, MCHC and the number of leukocytes. Administration BCSO of 5 ml/kg BW decreased the number of platelet and neutrophils but increased the number of lymphocytes. All of BCSO dose increased phagocytic activity, secretion of ROI, NO and IL-12p40 and expression of TLR-4.

Acknowledgment

Researchers gave a thank you to the staff and director of the Institute for Testing and Research of UGM who have given permission and facilities for the implementation of this study.

References

- Colombo MP, Trinchieri G. Interleukin-12 in anti-tumor immunity and immunotherapy. *Cytokine Growth Factor Rev* 2002; 13: 155-168.
- Curtsinger JM, Schmidt CS, Mondino A, Lins DC. Inflammatory cytokines provide a third signal for activation of naïve CD4 and CD8 T cells. *J Immunol* 1999; 162: 3256-3262
- Del Vecchio M, Bajetta E, Canova S, Lotze MT, Wesa A. Interleukin-12: biological properties and clinical application. *Clin Cancer Res* 2007; 13: 4677-4685.
- Watford WT, Moriguchi M, Morinobu A, OShea JJ. The biology of IL-12: coordinating innate and adaptive immune responses. *Cytokine Growth Factor Rev* 2003; 14: 361-368.
- Meeran SM, Akhtar S, Katiyar SK. Inhibition of UVB-induced skin tumor development by drinking green tea polyphenols is mediated through DNA repair and subsequent inhibition of inflammation. *J Invest Dermatol* 2009; 129: 1258-1270.
- Nickavar B, Mojab F, Javidnia K, Amoli MA. Chemical composition of the fixed and volatile oils of *Nigella sativa* L. from Iran. *J Naturforsch* 2003; 58: 629-631.
- Shahzad M, Yang X, Raza Asim MB, Sun Q, Han Y. Black seed oil ameliorates allergic airway inflammation by inhibiting T-cell proliferation in rats. *Pulm Pharmacol Ther* 2009; 22: 37-43.
- El Gazzar MAEL, Mezayen R, Nicolls MR, Dreskin SC. Thymoquinone attenuates proinflammatory responses in lipopolysaccharide-activated mast cells by modulating NF- κ B nuclear transactivation. *Biochim Biophys Acta* 2007; 1770: 556-564.
- El Gazzar M, El Mezayen R, Nicolls MR, Marecki JC, Dreskin SC. Downregulation of leukotriene biosynthesis by thymoquinone attenuates airway inflammation in a mouse model of allergic asthma. *Biochim Biophys Acta* 2006; 1760: 1088-1095.
- Finlay TM, Abdulkhalek S, Gilmour A, Guzzo C, Jayanth P, Amith SR, Gee K, Beyaert R, Szewczuk MR. Thymoquinone-induced Neu4 sialidase activates NF κ B in macrophage cells and pro-inflammatory cytokines in vivo. *Glycoconj J* 2010; 27: 583-600
- Finlay TM, Jayanth P, Amith SR, Gilmour A, Guzzo C, Gee K, Beyaert R, Szewczuk MR. Thymoquinone from nutraceutikankerl black cumin oil activates Neu4 sialidase in live macrophage, dendritic, and normal and type I sialidosis human fibroblast cells via GPCR Galphai proteins and matrix metalloproteinase-9. *Glycoconj J* 2010; 27: 329-348.
- Zaoui A, Cherrah Y, Alaoui K, Mahassine N, Amarouch H. Effects of *Nigella sativa* fixed oil on blood homeostasis in rat. *J Ethnopharmacol* 2002; 79: 23-26.
- Akrom WA, Djannah SN. Effect of the ethanolic extract of *Nigella sativa* on phagocytosis activity of macrophage male swiss mice during infection of *Listeria monocytogenes*, The 4th International Eijkman Conference, Bali 2007.
- Finlay TM. Thymoquinone is A novel ligand which activities neu4 sialidase to promote a pro-inflammatory response. Thesis Queens University Kingston Ontario Kankernada 2009.
- El-Mahmoudy A, Shimizu Y, Shiina T, Matsuyama H, Nikami H, Takewaki T. Macrophage-derived cytokine and nitric oxide profiles in type I and type II diabetes mellitus: effect of thymoquinone. *Acta Diabetol* 2005; 42: 23-30.
- Murphy KP. *Janeways Immunobiology*, Garland Science New York USA 2012.
- Wang D, Zhu T, Zeng S, Kankero Y, Cui J, Deng X, Song Y. Effects of scrocaffeside A from *Picrorhiza Scrophulariiflora* on immunocyte function in vitro. *Immunopharmacol Immunotoxicol* 2009; 31: 451-458.
- Cheikh-Rouhou S, Besbes S, Hentati B, Blecker C, Deroanne C, Attia H. *Nigella sativa* L. Chemical composition and physicochemical characteristics of lipid fraction. *Food Chemistry* 2007; 101: 673-681.
- Swann JB, Vesely MD, Silva A, Sharkey J, Akira S, Schreiber R, Smyth M. Demonstration of inflammation-induced cancer and cancer immunoediting during primary tumorigenesis. *PNAS* 2008; 105: 652-656.
- Sica A, Bronte V. Altered macrophage differentiation and immune dysfunction in tumor development. *J Clin Invest* 2007; 117: 1155-1166.
- Bastos KR, Marinho CR, Barboza R, Russo M, Alvarez JM. What kind of message does IL-12/IL-23 bring to macrophages and dendritic cells? *Microbes Infect* 2004; 6: 630-636.
- Jiang H, Chess L. An integrated view of suppressor T cell subsets in immunoregulation. *J Clin Invest* 2004; 114: 1198-1208.
- Thiery J, Dorothee G, Haddada H, Echchakir H, Richon C. Potentiation of a tumor cell susceptibility to autologous CTL killing by restoration of wild-type p53 function. *J Immunol* 2003; 170: 5919-5926.

***Correspondence to**

Akrom

Department of Pharmacology and Clinical Pharmacy

Ahmad Dahlan University

Indonesia

Black cumin seed oil increases phagocytic activity and secretion of IL-12 by macrophages.

ORIGINALITY REPORT

6%

SIMILARITY INDEX

PRIMARY SOURCES

- | | | |
|---|---|-----------------|
| 1 | phylodiversity.net
Internet | 36 words — 1% |
| 2 | api.uinjkt.ac.id
Internet | 19 words — 1% |
| 3 | Reena Mukherjee. "Selenium and vitamin E increases polymorphonuclear cell phagocytosis and antioxidant levels during acute mastitis in riverine buffaloes", <i>Veterinary Research Communications</i> , 04/2008
Crossref | 15 words — < 1% |
| 4 | www.freepatentsonline.com
Internet | 15 words — < 1% |
| 5 | Titiek Hidayati, Akrom, Indrayanti, Sagiran. "Evaluation of the Black Cumin Seed Oil Role (BCSO) on a Decline in eNOS Expression and Plasma NO Levels: Initial Studies Kemopreventive BCSO for Lung Cancer", <i>International Journal of Bioscience, Biochemistry and Bioinformatics</i> , 2017
Crossref | 13 words — < 1% |
| 6 | S Cadenas. "Oxidative DNA damage estimated by oxo8dG in the liver of guinea-pigs supplemented with graded dietary doses of ascorbic acid and alpha- tocopherol", <i>Carcinogenesis</i> , 12/01/1997
Crossref | 12 words — < 1% |
| 7 | discovery.ucl.ac.uk
Internet | 11 words — < 1% |
-

- 8 Lee, K.Y.. "Subchronic toxicity study of dietary N-acetylglucosamine in F344 rats", Food and Chemical Toxicology, 200404
Crossref 10 words — < 1%
-
- 9 www.plosone.org
Internet 9 words — < 1%
-
- 10 Bastos, K.R.B.. "What kind of message does IL-12/IL-23 bring to macrophages and dendritic cells?", Microbes and Infection, 200405
Crossref 9 words — < 1%
-
- 11 Tien-Fu Chuang. "Electroporation-mediated IL-12 gene therapy in a transplantable canine cancer model", International Journal of Cancer, 2009
Crossref 9 words — < 1%
-
- 12 pubs.sciepub.com
Internet 9 words — < 1%
-
- 13 Al-Gayyar, Mohammed M. H., Hanan M. Hassan, Abdullah Alyoussef, Ahmed Abbas, Mohamed M. Darweish, and Amany A. El-Hawwary. "Nigella sativa oil attenuates chronic nephrotoxicity induced by oral sodium nitrite: Effects on tissue fibrosis and apoptosis", Redox Report, 2016.
Crossref 8 words — < 1%
-
- 14 www.ijbbb.org
Internet 8 words — < 1%
-
- 15 archneur.highwire.org
Internet 8 words — < 1%
-
- 16 Systems Biology of Free Radicals and Antioxidants, 2014.
Crossref 7 words — < 1%

