

Lymphocyte Proliferation Activity Mtt -Test Of Ethanolic Extract Of Pasak Bumi Root (Eurycoma Longifolia Jack) On Induced 7,12 Dimethylbenz[A] Anthracene (Dmba) Female Sprague Dawley Rat

By SAPTO YULIANI

Lymphocyte Proliferation Activity MTT -Test of Ethanolic Extract of Pasak Bumi Root (*Eurycoma longifolia* Jack) on Induced 7,12-Dimethylbenz[A] Anthracene (DMBA) Female Sprague Dawley Rat

Lily Noor Falah and Sapto Yuliani

Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta
Jl. Prof. Soepomo SH Janturan, Telp: (+62274) 379418, Yogyakarta, Indonesia

Abstract

Pasak bumi root (*Eurycoma longifolia* Jack) contains quasinoid. Quasinoid potentially has antitumor/anticancer, anti-parasitic, and immunostimulatory activity. The aim of this research was to study the lymphocyte proliferation activity MTT-test of ethanol extract of *Eurycoma longifolia* Jack root on induced by 7,12-dimethylbenz[a] anthracene (DMBA) female Sprague Dawley rat. The test was done at 6 groups of 10 SD rat each. Each groups was administered orally with DMBA (Group I) dose of 20 mg/kg bw, corn oil (Group II), baseline (Group III), and ethanol extract of pasak bumi root dose 12.6; 25.2; and 50.4 mg/kg bw (Group IV,V, and VI). Ethanol extract is given for 5 weeks. At the second week five rats were sacrificed to isolate the lymphocytes of spleen then rats are injected orally with DMBA 2 times a week for 5 weeks. At the eight until twenty fourth week all groups were just administered by aquades. Measurement of spleen relative weight and isolation of the lymphocyte spleen were done at second week and the last of experiment. Lymphocyte proliferation activity were measured by MTT-reduction method. Data were analyzed statistically by analyzes of variance (anova) continued by LSD test and Kruskal Wallis continued by Mann Whitney test. The result showed that lymphocyte proliferation activity MTT of ethanol extract of *E. longifolia* root doses of 12.6; 25.2; and 50.4 mg/kg bw for 14 days prior to DMBA-induced are 339.35 ± 8.43 , 61.20 ± 6.27 , 310.28 ± 6.81 and dose 50.4 mg/kg bw after DMBA-induced is 122.108 ± 11.426

Key word: *E. longifolia* root, spleen, lymphocyte cell, 7,12-Dimethylbenz (a) anthracene (DMBA)

6

INTRODUCTION

Eurycoma longifolia Jack, known in Indonesia as Pasak bumi, has been widely used as traditional medicine for aphrodisiac, antiplasmodium and anticancer (Ang et al., 2001). Pasak bumi herbal contains chemical compound such as eurycomalacton, lauricomalacton A, dehidroeuricomalacton, euricomanon, euricomanol, benzoqui-non, sterol, saponin, ester sterol fatty acid (Supriyadi et al., 2001), quasinoid, xanthin-6-on alkaloid, β -carbolin alkaloids, tiruccallan tipe triterpens, derivatif squallen, dan biphenilneolignans (Kuo et al., 2004). Activity of quasinoid as antitumor have been researched by Jiwanjinda et al (2003).

Breast cancer is a serious problem worldwide, increases up to 1-2% annually in many countries. Since year 2000 approximately there are additional 1 million women worldwide suffered from this disease. The incidence of breast cancer in Indonesia women are ranked second after cervical cancer (Tjindarbumi and Mangunkusumo, 2002). The therapy currently used including surgery, radiotherapy and chemotherapy were still unsatisfied as the side effect and non selectivity of radiotherapy and chemotherapy.

*Corresponding author email : syuliani@yahoo.com

3 Development of tumor immunotherapies focuses on inducing autoimmune responses against tumor. One of parameters system immune activity is lymphocyte proliferation activity in spleen. Splenic lymphocyte proliferation is usually followed to evaluate a general effect on immune cells.

METHODS

Materials

E. longifolia Jack root were obtained from Kalimantan, DMBA (7,12-dimethylbenz(a)antracen) and RPMI medium were purchased from Sigma Chem. Other chemicals such as Phospat Buffered Saline (PBS), Sodium Dodecyl Sulphat (SDS), MTT(3-(4,5-dimethylethiazol-2-yl)-2,5-diphenyle tetrazolium bromide), 1N NaOH, 1N HCl, Tris Buffered Ammonium Chloride (TBAH) were obtained from Integrated Research and Testing Laboratory, Gadjah Mada University.

Extraction of *E. longifolia* root with ethanol

Identification of the samples was done using standard botanical monographs. The ethanol extract was prepared by cold maceration of 250 g of *E.longifolia* root powder in 500 ml of ethanol 70% allowed to stand overnight. The solution was then filtered, concentrated, dried in rotary evaporator and 63.5 g of thick dark brown extract residue stored in a refrigerator at 2-8°C for subsequent experiments.

Animal treatment

Female albino Sprague Dawley rats, weighing 100-180 g obtained from Gadjah Mada University breeding colony were used for the present investigations. The animals were maintained on standard rat feed supplied Comfeed. The test was done at 6 groups of 10 SD rat each. Group I (cancer control) was given aquadest, group II was given corn oil,

Group III was as baseline, Group IV,V, VI were given ethanol extract of pasak bumi root dose 12.6; 25.2; and 50.4 mg/kg bw. After 2 weeks, all groups except of group II and III, were administered orally by DMBA dose of 20 mg/kg bw 2 times a week for 5 weeks. Observation of tumor multiplicity and tumor incidence were done for sixteen weeks after the last DMBA administration. Measurement of spleen relative weight and isolation of the lymphocyte were done at second week and the end of experiment.

Spleen Lymphocyte Cell Isolation

Rats were weighed and then sacrificed by cervical dislocation. Spleens were removed by aseptic technique, weighed then mechanically dissociated into a single cell suspension in RPMI-1640 medium and centrifuged at 1200 rpm, temperature 4°C for 10 min. Centrifugation eliminates polymorphonuclear cells, erythrocytes and dead cells. The band containing the mononuclear cells was washed, pelleted and resuspended in Tris Buffered Ammonium Chloride (TBAH) and Phospat Buffered Saline (PBS). This suspension of splenic mononuclear cells was defined as the splenocytes. Cells were counted in a haemocytometer. Lymphocytes cell were cultured in CO₂ incubator at 37 °C.

MTT-reduction method

MTT (3-(4,5-dimethylethiazol-2-yl)-2,5-diphenyle tetrazolium bromide (Sigma) was dissolved in PBS at 5 mg/ml and filtered to sterilize and remove a small amount of insoluble residue present in some batches of MTT. Stock MTT solution (10 µl) medium was added to all wells of an assay, and plates were incubated at 37°C for 4 h. Sodium Dodecyl Sulfat (SDS) was added to all wells and thoroughly to dissolve the dark blue crystals. After a few minutes at room temperature to ensure that all crystals were dissolved, the plates were read on Elisa reader.

Data Analyzes

Calculation of relative spleen weigh:

$$\text{Relative spleen weigh} = \frac{\text{Spleen weigh(g)}}{\text{The last body weigh(g)}}$$

Calculation of % lymphocyte proliferation activity:

$$\% \text{ lymphocyte proliferation activity} = \frac{\text{Absorbance}_{\text{treatment}} - \text{Absorbance}_{\text{medium}}}{\text{Average of Absorbance}_{\text{dmba group}} - \text{Absorbance}_{\text{medium}}} \times 100 \%$$

All data are expressed as mean and standard deviation. Relative spleen weight prior and after DMBA- induced were analyzed statistically using one way ANOVA continued by LSD test. Percentage of lymphocyte proliferation activity prior to and after DMBA-induced was analyzed using Kruskal Wallis test continued by Mann Whitney test.

RESULTS

Spleen weight

Relative weight of the spleen was measured by analytical balance and expressed

in gram. A significant increase in the weight of spleen was observed in ethanol extract of *E. longifolia* (group IV, V, and VI) when compared to other group (group I, II, III) prior to DMBA-induced. Administration of *E. longifolia* dose 12.6mg/Kg bw/day(group IV), dose 25.2mg/Kg bw/day(group V), dose 50.4mg/Kg bw/day(group VI) for 2 week prior to DMBA-induced can significant increase relative spleen weight. There was not differ significantly among the *E.longifolia* groups. On the other hand the relative spleen weight of all groups after DMBA-induced were not differ significantly when compared each other (Table 1).

Table 1. Relative spleen weigh (g) prior to and after DMBA-induced

Group	Before DMBA- induced	After DMBA-induced
DMBA (Group I)	0.0080 ± 0.0014 ^b	0.0026 ± 0.0005 ^d
Corn Oil (group II)	0.0074 ± 0.0008 ^b	0.0030 ± 0.0003 ^d
Baseline (group III)	0.0083 ± 0.0012 ^b	0.0021 ± 0.0008 ^d
dose 12.6mg/Kg bw/day(group IV)	0.0124 ± 0.0015 ^c	0.0036 ± 0.0007 ^d
dose 25.2mg/Kg bw/day(group V)	0.0106 ± 0.0022 ^c	0.0049 ± 0.0037 ^d
dose 50.4mg/Kg bw/day(group VI)	0.0108 ± 0.0025 ^c	0.0034 ± 0.0007 ^d

Data are expressed as mean ± SD (n =5 rats per group). Groups with different superscript letters are significantly different at $P < 0.05$ by analysis of variance continued by LSD test.

Lymphocyte Proliferation Activity

Ethanol extract of *E. longifolia* root administration dose of 12.6 mg/Kgbw/day, 25.2 mg/kgbw/day and 50.4 mg/Kgbw/day for 2 weeks (prior to DMBA-induced) can

significant increase in the percentage of proliferation lymphocyte activity. Ethanol extract dose 12.6 mg/Kgbw/day was not differ significantly with dose 50.4 mg/Kgbw/day. Ethanol extract *E.longifolia* root dose 25.2

mg/kgbw/day showed the lowest percentage of proliferation lymphocyte activity among the ethanol extract groups prior to DMBA-induced. Dose ethanol extract 50.4 mg/Kgbw/day after DMBA – induced showed the highest percentage of lymphocyte

proliferation activity among ethanol extract groups. Increasing of percentage of lymphocyte proliferation activity dose 12.6 mg/Kgbw/day, 25.2 mg/Kgbw/day and corn oil group were not differ significantly when compared with baseline group (Table 2).

Table 2. Percentage of lymphocyte proliferation activity in all groups prior to and after DMBA-induced

Groups	% of lymphocyte proliferation activity prior to DMBA- induced	% of lymphocyte proliferation activity after DMBA- induced
DMBA	100 ± 0 ^a	100 ± 0 ^a
Corn Oil	15.30 ± 0.61 ^b	51.928 ± 19.265 ^c
Base Line	17.44 ± 10.98 ^b	38.689 ± 10.885 ^c
dose 12.6 mg/Kgbw/day	339.35 ± 8.43 ^c	61.825 ± 10.554 ^c
dose 25.2 mg/Kgbw/day	61.20 ± 6.27 ^d	49.357 ± 14.628 ^{de}
dose 50.4 mg/Kgbw/day	310.28 ± 6.81b ^c	122.108 ± 11.426 ^f

1 Data are expressed as mean ± SD (n =5 rats per group). Groups with different superscript letters are significantly different at $P < 0.05$ by Kruskal Wallis continued by Mann Whytney test

DISCUSSION

In this study, DMBA is used to induce the occurrence of cancer. According to Dandekar et al., (1986) Dimetilbenz (a) anthracene (DMBA), known as specific carcinogenic compounds. Carcinogenic activity of DMBA occurs because of the ability of DMBA metabolites (ultimate carcinogens) binds to DNA and cause somatic mutations that encourages rapid cell division (Miyata et al., 1999). The spleen is an organ found in virtually all vertebrate animals with important roles in regard to red blood cells and the immune system. Underwood (1999) explained that the white pulp is the biggest place that produce antibodies and to form the centrum germinativum when stimulated. Forming spleen white pulp lymphocytes is play a role in the immune system. Increased immune system will provide the protective effects of the body against foreign agents (antigens) that can damage cells.

Based on macroscopic observations show that not all rats edge taper-shaped organ. According to Brake (1997) exposure to the spleen by imunogen will evoke an immune activity of the spleen. In the event of increased immunity, spleen lymphocyte proliferation activity will increase so that the morphology of

spleen size became larger. Normal spleen organ has the shape of the edge of the taper or crescent-like shape. Organs which have an enlarged spleen or swelling, the shape tends to blunt or rounded edges.

As shown at Table 1 that group IV, V, VI with ethanol extract of *E. longifolia* root administration prior to DMBA- induced differ significantly with groups non-ethanol extract of *E. longifolia* root administration (I, II, and III). Relative spleen weight was significantly higher in all doses of ethanol extract of *E. longifolia* root groups when compared with the control group. The result suggest that all of dose ethanol extract of *E. longifolia* root administration can increase relative spleen weigh of rats prior to DMBA-induced.

Lymphocyte proliferation is activation phase of immune system. Increasing of spleen activity may indicate an increase of lymphocyte proliferation causing an increase spleen in size. However, ethanol extract of *E. longifolia* can not affect to the relative spleen weight after DMBA-induced. This result was supported by Nurhaini (2009) that ethanol extract of *E. longifolia* can not affect histopathologic of spleen on rat induced by DMBA.

In this research determination of lymphocyte proliferation activity using MTT

method. MTT-test as alternative method to measurement of the cellular activity and proliferation. Indirectly counting using MTT assay is based on cell capability in order to reduce MTT to insoluble water formazan (Fig. 1). Grey salt is correlation with a number of cell that can be detected by colorimetry (Shalini and Harriom, 2009). In this study determination of lymphocyte proliferation activity was done at second week and end of the treatment. Percentage of the increasing lymphocyte proliferation activity were presented in Table 2. Extract treatment group had a higher percentage proliferation activity compared with the CMC Na group (solvent control). These results indicate that administration of ethanol extract of *E. longifolia* Jack for 14 days can increase the lymphocyte proliferation activity significantly getting CMC Na.

The results of lymphocyte proliferation activity in the 16th week after the last DMBA administration showed that DMBA can improve lymphocyte proliferation activity significantly when compared with solvent control (corn oil). This is probably caused that DMBA is a carcinogen which can be recognized by the immune system so that increased lymphocyte proliferation (Kresno, 2002). The lymphocyte highest proliferation activity after administration of DMBA is shown in the dose of ethanol extract of 50.4 mg / kg bw / day. The biochemistry of the immunostimulatory activity by ethanol extract of *E. longifolia* roots may be caused that it serves as imunogen which kuasinoid able to be recognized by T cell receptor. Components of the ethanol extract of the roots can bind to T cell surface receptor through hydrogen bonding. Antigen binding receptor on the surface of T cells with interleukin 1 (IL-1) of the APC (Antigen Presenting Cell) may activate G-proteins which then produce phospholipase C. This enzyme hydrolyzes fosfatidil inositol bisphosphate (PIP2) into a reactive product diacylglycerol (DAG) and inositol triphosphate (IP3). The reaction took place in the plasma membrane. IP3 then stimulates the release of Ca^{2+} into the cytoplasm so that the concentration of Ca^{2+}

and have been able to act as imunogen. The results of this study shows compatibility with previous studies that exposure to the spleen by imunogen will evoke an immune activity so that the size of the spleen and spleen lymphocyte proliferation activity will increase. (Hargono, 2000; Akrom, 2004). From this study demonstrated that administration of ethanol extract of *E. longifolia* Jack for 14 days on animal Sprague Dawley female rats prior to DMBA-induced turns out to as immunostimulatory activity with a mechanism for increased lymphocyte proliferation. The results are consistent with data supporting the relative weights of the spleen that the extract dose group 12.6, 25.2, 50.4 mg / kg bw / day can increase the relative spleen weight significantly compared with the group that just

increase. Increased Ca^{2+} plays an important role in stimulating action of the enzyme protein kinase C and 5-lipoxygenase. Protein kinase C production of interleukin 2 (IL-2), IL-2 was then activates B cells and T cells to proliferate (Ikawati, 2006). Solving advanced through the DAG into arachidonic 5-lipoxygenase enhances the formation of cGMP. Increased cGMP resulted in increased activity of cGMP-dependent protein kinase that functions in the activation of DNA-dependent RNA polymerase, and synthesis of ribosomal (rRNA) and other RNA. RNA and protein synthesis causes B cells and T lymphocytes entering the phase of cleavage (Ikawati, 2006).

CONCLUSION

Based on the results of this study can be concluded that lymphocyte proliferation activity MTT of ethanol extract of *E. longifolia* root doses of 12.6; 25.2; and 50.4 mg/kg bw for 14 days prior to DMBA-induced are 339.35 ± 8.43 , 61.20 ± 6.27 , 310.28 ± 6.81 and dose 50.4 mg/kg bw after DMBA-induced is 122.108 ± 11.426 . Administration of ethanol extract of *E. longifolia* roots of dose 50.4 mg / kg / day can increase the activity of rat spleen lymphocyte proliferation in female SD rat that has been induced DMBA.

REFERENCES

- Akrom, 2004, Cellular immune activity of Male Swiss Mice Infected Plasmodium berghei with The Ethanol Extract Phyllanthus niruri, Research Report, Fac. of Pharmacy, Ahmad Dahlan University, Yogyakarta
- Ang, H.H., Ngai, T.H., 2001, Aphrodisiac evaluation in non-copulator male rats after chronic administration of Eurycoma longifolia, *Fundamental & Clinical Pharmacology*, **15**; 265-268
- Dandekar, S., Sukumar, S., Zaeld, H., Yuong, L., J.T., Cardiff, R.D., 1986, Spesific Activation of The Cellular Harvey-ras Oncogen in Dimethylbenzanthracene-Induced Mouse Mammary Tumors, *Molecular and Cellular Biology*, vol **6** no.11, p. 4104-4108.
- Hargono, D, Winarno, M. W., and Werawati, A., 2000, Effect of Gynura Ngokilo Leaf Juice (Procumbers Lour Meer) on Mice againts Immune System Activity, *Cermin Dunia Kedokteran*, No. 127
- Ikawati, Z., 2006, *Pengantar Farmakologi Molekuler*, Yogyakarta, Gadjah Mada University Press
- Jiwajinda, S., Santisopasari, V., Murakami, A., Kawanaka, M., Kawanaka, H., Gasquet, M., Elias, R., Balansard, G., Ohigashi, H., 2002, Quassinoid from Eurycoma longifolia as Plant growth inhibitors, *Phytochemistry*, **58**:959-962
- Kresno, S. B., 2001, *Imunologi: Diagnosis dan prosedur Laboratorium*, Ed IV, Medi , Jakarta
- Kuo, P.C., Damu A.G., Lee K.H., Wu TS., 2003, Cytotoxic and Antimalarial beta-carboline alkaloids from the roots of Eurycoma longifolia, *J. Nat Product*, Oct; **66** (10); 1324
- Miyata, M., Kudo, G., Lee, Y., Yang, T.J., Gelboin, H.V., Gonzalez, F.J., 1999, Targeted Disruption of the Microsomal Epoxide Hydrolase Gene, *J. Biol. Chem.*, vol **274**, hal 23963-23968.
- Shalini, J and Hariom Y., 2009, Cell Proliferation Assay, Protocol Online.
- Supriyadi, H,M & Taryono., 2001, Pasak bumi (Eurycoma longifolia Jack) Plant Drug Indonesia, Use and Medicinal Properties, Pustaka Populer Obor, Jakarta
- Tjindarbumi, D. dan Mangunkusumo, R., 2002. Cancer in Indonesia, Present and Future, *Jpn J. Clin Oncol* : **32** (Supplement 1),17-21
- Underwood, J. C. E. 1993. General and Systemic Pathology, 2nd d., Vol II. Editor : Sarjadi. Jakarta : EGC.

Lymphocyte Proliferation Activity Mtt -Test Of Ethanolic Extract Of Pasak Bumi Root (Eurycoma Longifolia Jack) On Induced 7,12 Dimethylbenz[A] Anthracene (Dmba) Female Sprague Dawley Rat

ORIGINALITY REPORT

3%

SIMILARITY INDEX

PRIMARY SOURCES

1	Yang, Ruei-Cheng, Ming-Hong Chen, Pei-Yu Chen, Ching-Yun Chen, Shih-Feng Tsai, Cheng-Kung Cheng, and Jui-Sheng Sun. "A mutation of the Col2a1 gene (G1170S) alters the transgenic murine phenotype and cartilage matrix homeostasis", Journal of the Formosan Medical Association, 2014.	14 words — 1%
2	"4. Comments on residues of specific veterinary drugs.(p. 25-59)", WHO Technical Report Series, Sept 2015 Issue	13 words — < 1%
3	Wolfgang Dummer. "T cell homeostatic proliferation elicits effective antitumor autoimmunity", Journal of Clinical Investigation, 07/15/2002	11 words — < 1%
4	mdpi.com	8 words — < 1%
5	Bilal Ahmad, Muneeb U Rehman, Insha Amin, Manzoor ur Rahman Mir et al. "Zingerone (4-(4-hydroxy-3-methylphenyl) butan-2-one) protects against alloxan-induced diabetes via alleviation of oxidative stress and inflammation: Probable role of NF-kB activation", Saudi Pharmaceutical Journal, 2018	8 words — < 1%

6 L H Nurani, A Mursyidi, S Widyarini, A Rohman. " 8 words — < 1%
The effects of combination of Jack ethanolic extract
and doxorubicine on hematological profile in rats
given by 7,12-dimethylbenz(a)anthracene ", IOP Conference
Series: Materials Science and Engineering, 2017
Crossref

7 Pezzuto, John M.. "Resveratrol as an Inhibitor of 8 words — < 1%
Carcinogenesis", Pharmaceutical Biology, 2008.
Crossref

8 www.webmd.com 8 words — < 1%
Internet

EXCLUDE QUOTES ON
EXCLUDE BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF