

9. 933-2593-1-PB

by Irip123 Mpfis

Submission date: 22-Jan-2020 04:41PM (UTC-0600)

Submission ID: 1245116340

File name: 9._933-2593-1-PB.pdf (118.76K)

Word count: 3636

Character count: 19241

Research Article

3

THE CORRELATION BETWEEN WORM INFECTIONS AND NUTRITIONAL STATUS AMONG ELEMENTARY SCHOOL STUDENTS

Liena Sofiana*, Erni Gustina, Yuniar Wardani, Suci Musvita Ayu, Aniq Diya Nata Maula

2

Department of Public Health, Public Health Faculty, Universitas Ahmad Dahlan, Yogyakarta, Indonesia

8

*Correspondence: liena.sofiana@ikm.uad.ac.id. Telp.: +628587555584

Received 07 August 2019; Accepted 09 September 2019; Published 09 September 2019

ABSTRACT

Background: Worm¹ disease is an infectious disease that still becomes a public health problem in Indonesia. Worm can cause a decrease in health, nutrition, intelligence and productivity of the sufferers. Worms as parasitic animals do not only take nutrients in the intestines of children, but they also damage the intestinal wall so that they interfere the absorption of these nutrients, consequently it will impact² on the decreasing of one's nutritional status and cause other complex nutritional problems. Moyudan Health Center is a Community Health Center in Moyudan² Sub-District Region which has the highest cases of *helminthiasis* (worm infection). Therefore, this study aims to determine the¹⁴ relationship between *helminthiasis* infection and nutritional status.

Methods: This was an observational analytic study² by using a cross sectional approach. The sample of this study were elementary school students in the work area² of Moyudan Health Center with samples of 311 respondents. The sampling technique used by multistage random sampling is the selection of samples consisting of the smallest areas of an area. Faecal examination was carried out by using the direct method and nutritional status assessment was carried out² by anthropometric method based on the anthropometric index Body Mass Index per Age (BMI / U). Data were analyzed by Fisher's test.

Results: The results showed that there were 8 respondents (2.57%) who were positively infected by worms, 28 respondents had underweight nutritional status (9.01%), and 54 respondents had a nutritional status of fat (17.36%). The results of the study showed no correlation between *helminthiasis* infection and¹³ nutritional status (p value = 0.534).

Conclusions: There is no relationship between *helminthiasis* and nutritional status⁸ in elementary school students in the Moyudan Community Health Center Working Area in Sleman Regency with a p value of 0.534 so that it is concluded that *helminthiasis* infection does not provide any significant contribution to nutritional status even though cases of *helminthiasis* still occur in elementary school students. Therefore, it needs more attention from parents, schools and health facilities in dealing with worm problems.

Keywords: Worm Infection, Nutritional Status, Elementary School Students

INTRODUCTION

Worm disease is an infectious disease that is still a public health problem in Indonesia because it spreads in most regions of Indonesia. Worm can cause health, nutrition, intelligence and productivity problems. Similarly, it can also cause loss of fat and protein and blood loss, thus reducing the quality of human resources (1). Worms are part of 17 Neglected Disease (NTD) (2). The most common types of worm that infect humans are *Ascaris lumbricoides*, *Necator americanus*, *Ancylostoma duodenale*, *Trichuris trichiura*, *Strongyloides stercoralis*, *Trichinella spiralis* and *Enterobius vermicularis* (3).

More than 1.5 billion people or 24% of the world's population are infected by earthworms that spread throughout the world. Infection is widespread in tropical and subtropical regions with the largest number occurring in Sub-Saharan, Africa, America, China and East Asia. More than 267 million pre-schoolers and more than 568 million school-aged children live in the area while these parasites are intensively transmitted and intervened. Therefore, treatment is immediately needed (4). The ones who often get infected are pre-schoolers, school-age children, women of childbearing age (including pregnant women in the second and third trimesters and breastfeeding), and adults with high-risk jobs such as tea pickers or miners are also vulnerable to worm infection (4). This disease generally attacks children because their resistance is still low. The factors that influence it are tropical condition, low awareness of cleanliness, poor sanitation, low socio-economic conditions, and population density (5).

Worms as parasitic animals do not only take nutrients in the child's intestine, but they also damage the intestinal wall so that they release these nutrients (6). Worms get food from tissues including proteins that will cause substances of protein and iron. Worms will also increase malabsorption nutrition. Some earthworms can also cause reduced appetite and therefore they can reduce nutritional intake and physical fitness (4).

The prevalence data of worm infection in 2016 in Sleman suggested that there were 559 cases of infection and the highest of all was found in Moyudan Community Health centre with 182 cases (7). Meanwhile, the screening data of students' weight and height from two elementary schools in Moyudan sub-district showed that 6 out of 170 students were categorized fat and 136 of them were normal. In addition, 12 students were thin and 16 were very thin. After studying the data, the researcher thinks that it is necessary to research this case further as it is important to know how big this case is and what efforts can be done to control and prevent this case.

METHODS

Study design

This research belongs to analytic observational research with cross sectional approach.

Setting

This research was conducted among the elementary students in the coverage area of Moyudan community health centre, Sleman, Yogyakarta. There were 21 elementary schools from 4 villages.

Data source and data collection

The data collection was done by measuring body weight and height to determine BMI (Body Mass Index) of students in the elementary schools. Meanwhile, the nutritional status was measured by using scales and microtoise. Worm examination was carried out with a direct method using a microscope conducted at the Moyudan Community Health Center laboratory. In addition, examination of the number of worm species in faecal samples was also carried out by counting the number of eggs, larvae or adult worms in the preparation.

Sample population

The research sample were elementary school students in the work area of Moyudan Health Center with a total of 311 samples. The school sampling technique was carried out by using probability sampling by means of multiple stage random sampling. 21 elementary schools from 4 villages were randomly selected and each of them had 3 primary schools which were sampled. Thus, the number of primary schools used was 12 schools which consists of grade1 to grade 6. The sample consists of

286 people so that the total sample taken from each primary school were 24 people. Respondent samples were taken randomly by using simple random sampling.

Variables

The independent variable in this research was worm infection while the dependent variable was nutritional status of elementary school students.

Analysis

The analysis was done using chi square test with trust interval of 95%. The value of α was 0.05 which also used fisher as its alternative. The requirement for chi square test is when there is no value of expected count below 5 or maximum 20% in cell.

Ethical approval

Number: 011804053

The ethical approval is based on ethical research committee of Universitas Ahmad Dahlan Yogyakarta.

RESULTS

Table 1. Respondent Distribution based on Gender and Age

Variable	Frequency	Percentage (%)
Age (Year)		
5	1	0,3
6	8	2,6
7	50	16,1
8	64	20,6
9	64	20,6
10	53	17,0
11	55	17,7
12	15	4,8
13	1	0,3
Gender		
Male	150	48,0
Female	161	52,0
Total	311	100,0

Source: Primary Data of 2018

Table 1 shows that the genders of the respondents are different. Male respondents consisted of 150 students (48.23%) and female respondents are 161 students (51.77%). In terms of age, the table shows that most of respondents are 8- and 9-years old totalling 64 students (20.58%) while the least number is those aged between 5 to 13 years old accounting only 1 person (0.32%).

Table 2. Worm Infections Distribution based on Gender

Variable		Gender				Total Number	Percentage (%)
		Male		Female			
		Total	Percentage (%)	Total	Percentage (%)		
Worm Infection	Positive	6	1.93	2	0.64	8	2.57
	Negative	144	46.30	159	51.13	303	97.43
	Total	150	48.23	161	51.77	311	100.0
Nutritional Status	Thin	11	3.54	17	5.47	28	9.01
	Normal	111	35.69	118	37.94	229	73.63
	Fat	28	9.00	26	8.36	54	17.36
	Total	150	48.23	161	51.77	311	100.0

Source: Primary Data of 2018

Table 2 shows that most of respondents infected with helminthiasis were male with a total of 6 respondents, while female who were infected with helminthiasis were only 2. Based on the nutritional status and sex of respondents, those with the lowest nutritional status were mostly female by 5.47%. Respondents with the most normal nutritional status were also female by 37.94%, while respondents with surplus nutritional status were more dominated by male, by 9%.

Table 3. The Distribution of Worm Infection Based on Worm Type

Worm Type	Gender		Total	Percentage (%)
	Male	Female		
<i>Ascaris lumbricoides</i>	0	1	1	12.5
<i>Necator americanus</i>	1	0	1	12.5
<i>Ancylostoma duodenale</i>	2	0	2	25.0
<i>Enterobius vermicularis</i>	2	1	3	37.5
<i>Trichuris trichiura</i>	1	0	1	12.5
Total	6	2	8	100.0

Source: Primary Data of 2018

Table 3 shows that of the 8 respondents who were positively infected with helminthiasis, in which most of them were infected with *Enterobius vermicularis* by 37.5% and *Ancylostoma duodenale* by 25%. Other types of infecting worms were *Ascaris lumbricoides*, *Necator americanus* and *Trichuris trichiura*, by 12.5% consecutively.

Table 4. Fisher's test results between helminthiasis infection and nutritional status in elementary school students in the work area of Moyudan Public Health Center, Sleman Regency

Variable	Nutritional Status				Total	RP	95% CI	P Value
	Thin		Normal + Fat					
	n	%	n	%				
Positive	1	0.3	7	2.3	8	2.6	1.403	0.217-9.087
Negative	27	8.7	276	88.7	303	97.4		
Sum	28	9.0	283	91.0	311	100		

Table 4 shows that after cell merging and fisher test for a 2x2 table, it is found that there is 1 cell (25%) that has an expected count value of less than 5, so that the chi-square test requirements are not met. Therefore, fisher test was used as an alternative. From the results of the fisher test it was known that the P value was 0.534, which means that there was no relationship between helminthiasis infection and nutritional status of elementary school students in the work area of Moyudan Health

Center, Sleman Regency. Fisher's test results also showed a value of Rp 1,403 with Confidence Interval of 95% (CI) 0.217-9,087 which means that from the existing data it could not be concluded that helminthiasis could really become a risk factor that affects nutritional status.

DISCUSSIONS

The description of the respondents of the elementary school students in the work area of Moyudan Public Health Center Sleman Regency were mostly 8 and 9 years old with a percentage of 20.58% while the least were students aged 5 and 13 years with a percentage of 0.32%. The percentage of female students was 51.77% which was greater than male students (48.23%). The percentage of students infected with helminthiasis was 2.57%, students with malnutrition status was 9.01%, normal was 73.63% and fat was 17.36%.

Based on table 5, it shows that out of 311 respondents there were 9.01% of respondents with malnutrition status, 17.36% of respondents were categorized fat while 73.63% of respondents were normal. From the results of this study it is known that there was more normal nutritional status than respondents who were thin and fat. Based on table 14, it is known that respondents with more or less nutritional status were experienced by respondents who were female and that is equal to 5.47%. Meanwhile respondents with surplus nutritional status were more experienced by male respondents with a percentage of 9%.

Worm disease is an endemic and chronic disease caused by worms with a high prevalence. It is however not lethal, but it risks the health of the human body resulting in a decrease in nutritional conditions and mostly attacks toddlers and children of primary school age (8). From the results of the study in table 3 it is known that out of 311 respondents there were 8 respondents who experienced helminthiasis or by 2.57%. The results of the study shown in Table 4 indicate that helminthiasis experienced by respondents occurred due to infection from several types of worms, namely *Ascaris lumbricoides* by 12.5%, *Americanus* by 12.5%, *Ancylostoma duodenale* by 25%, *Trichuris trichiura* by 12.5% and *Enterobius vermicularis* by 37.5%.

Based on observations the low prevalence of helminthiasis could be caused by good behaviours that they had such as hand washing habits, defecation in the toilet, nails cutting once a week, fingernails biting prevention, and footwear using. This can also be a cause of low prevalence of worms. Behaviour had a relationship with infection with Soil Transmitted Helminth in students at Pertiwi Lamgarot Elementary School in Ingin Jaya District, Aceh Besar (9). The difference in the number of helminthiasis infections transmitted through soil can be influenced by differences in risk factors including geography and environment, community behaviour towards environmental sanitation and personal hygiene (10).

¹⁸ The results of this study also showed that of the 8 respondents who were positively infected with helminthiasis, there were 6 respondents who were male and 2 females. This is in line with the previous studies which said that male sex can be a risk factor for infection with helminthiasis (OR = 1.27). This can be due to the fact that a boy more often plays outside the house, so he interacts more frequently with the soil. Boys usually play soccer, play marbles or make toys from the ground (10).

Based on table 6, it is known that out of 8 respondents infected with worms, there were 1 respondent who had underweight nutritional status, 6 respondents had good nutritional status and 1 respondent had fat nutritional status. This is in line with previous studies which showed that respondents who were positively infected with helminthiasis with a status of malnutrition were less compared to students who experienced helminthiasis with a good nutritional status (8). Student ⁴ with malnutrition status were 36.4% and 63.6% were nutritionally good. Previous research showed that there was no relationship between sex and nutritional status (p value = 0.550).

The results of statistical tests using Fisher's test for 2x2 tables or ⁷ cell merging that can be seen in table 16 which shows that the relationship between helminthiasis infection and nutritional status ⁴ in elementary school students in the Moyudan Health Center work area shows that statistically ¹¹ there was no significant relationship between infection with nutritional status (p value = 0.534). The results

of this study's test can be concluded that helminthiasis was a risk factor that affects nutritional status but was not statistically significant (RP = 1,403, 95% CI = 0.217-9,087).

The absence of a relationship between helminthiasis infection and nutritional status in elementary school students in the working area of Moyudan Health Center Sleman Regency could be due to the low number of children infected with helminthiasis, this is supported by the finding of the previous studies that the low intensity of helminthiasis was a factor that caused the absence in the relationship between helminthiasis and nutritional status (11). Besides, a good environmental condition around might also contribute to this case where most of the respondents' homes had already have the availability of latrines in the category of fulfilling the conditions. In addition, the distance of the source of water from the disposal of faeces was ≥ 10 m. moreover, most of the respondents' houses had used the type of floor that meets the requirements (12). Other studies also shown that many underweight children lived in slum areas with poor water sources, unhealthy latrines and poor sources of drinking water (13). The results of this study suggest that children who had poor nutritional status were not always due to helminthiasis infection, but there were other factors that affect the nutritional status. Previous research also shown the same thing that there was no relationship between helminthiasis infection and nutritional status based on BMI / U in children in Barabai darat 2 Primary School with a value of p value = 1,000. The results of the study showed that there was no significant influence between helminthiasis infection and nutritional status. One reason was because it was not only an infection that could affect the state of human nutrition, but it could also be caused by many other factors. In addition, the process from helminthiasis to a decrease in nutritional status requires a certain period of time and with the severity of certain infections (14).

Theoretically, helminthiasis, which is one of the infectious diseases, is one of the direct causes that will affect one's nutritional status. This worm can lead to a decrease in the condition of health, nutrition, intelligence and productivity of the sufferer so that it causes many economic losses. Worms can cause loss of carbohydrates and proteins and blood, thus reducing the quality of human resources. Worms as parasitic animals do not only take nutrients in the child's intestine, but also damage the intestinal wall so that they disturb the absorption of these nutrients (6). Nutritional loss is calculated based on the behaviour of each species of worm which harms the sufferer (15).

Calculation of nutrient loss and economic loss in 3 worm species, namely *Ascaris lumbricoides*, *Trichuris trichiura* and Hookworm was done because these three species provide the greatest loss to sufferers. Food intake and infectious diseases are the direct causes of malnutrition (15). Nutritional and blood loss is likely to increase as the worm's intensity increases in the patient's body, the rapid development of worms requires immediate treatment to prevent new infections from occurring (2).

CONCLUSIONS

Based on the results of the research and discussion, it could be concluded that there is no correlation between helminthiasis infection and nutritional status in elementary school students in the Moyudan Health Center Working Area in Sleman Regency.

AUTHORS' CONTRIBUTION

All researchers contributed to conducting research ranging from surveys, data collection, data analysis to the preparation of research reports.

FUNDING

Institute for Research and Community Service.

Acknowledgements

The researcher would like to thank Universitas Ahmad Dahlan Yogyakarta for providing support and funding for this research.

Conflict of interest

There are no conflicts of interest.

REFERENCES

1. Kementerian Kesehatan Republik Indonesia. Peraturan Menteri Kesehatan Republik Indonesia Nomor 15 Tahun 2017 Tentang Penanggulangan Kecacingan [Internet]. Jakarta; 2017 [cited 2018 Dec 28]. Available from: www.peraturan.go.id
2. Kesehatan BP dan P. Penyakit Menular Neglected Kajian Program dan Penelitian. Buku 1. Tjitra, Emiliana., Sudomo, Mohammad., Karyana M, editor. Jakarta: Kementerian Kesehatan Republik Indonesia; 2013. 43 p.
3. Sutanto I, Ismid I., Sjarifuddin P., Sungkar S. Buku Ajar Parasitologi Kedokteran. Jakarta: Balai Penerbit Fakultas Kedokteran Universitas Indonesia; 2009.
4. Organization WH. Soil-Transmitted Helminthiasis: Eliminating Soil-Transmitted Helminthiasis as a Public Health Problem in Children [Internet]. Progress Report. France: World Health Organization; 2012. 1-90 p. Available from: http://apps.who.int/iris/bitstream/handle/10665/44804/9789241503129_eng.pdf;jsessionid=08A9C47002D754E4C5CD704D24C26420?sequence=1
5. Kementerian Kesehatan Republik Indonesia. Penyakit Kecacingan Masih Dianggap Sepele [Internet]. Kementerian Kesehatan Republik Indonesia. 2010 [cited 2018 Dec 28]. p. 1. Available from: <http://www.depkes.go.id/article/print/1135/penyakit-kecacingan-masih-dianggap-sepele.html>
6. Kementerian Kesehatan Republik Indonesia. Pedoman Pengendalian Kecacingan. Jakarta: Kementerian Kesehatan Republik Indonesia; 2012. p. 1–48.
7. Dinas Kesehatan Kabupaten Sleman. Data Sekunder Dinas Kesehatan Kabupaten Sleman. Yogyakarta: Dinas Kesehatan Kabupaten Sleman; 2016.
8. M AR, Nelly M, Franly O. Hubungan Kecacingan Dengan Status Gizi Pada Murid Sekolah Dasar di Kabupaten Bolaang Mongondow Utara. *Ejournal Keperawatan*. 2013;1(1):1–6.
9. Rahmayanti, Razali, Mudatsir. Hubungan Pengetahuan, Sikap Dan Tindakan Dengan Infeksi Soil Transmitted Helminths (STH) Pada Murid Kelas 1, 2 dan 3 SDN Pertiwi Lamgarot Kecamatan Ingin Jaya Kabupaten Aceh Besar. *J Biot* [Internet]. 2014;2(hubungan pengetahuan,sikap,dan tindakan dengan infeksi soil transmitted helminths (STH) PADA MURID KELAS 1.2 dan 3 SDN PERTIWI LAMGAROT KECAMATAN INGIN JAYA KABUPATEN ACEH BESAR):77–137. Available from: <https://jurnal.ar-raniry.ac.id/index.php/biotik/article/view/244>
10. Sandy S, Sumarni S, Soeyoko. Analisis Model Faktor Risiko Yang Mempengaruhi Infeksi Kecacingan Yang Ditularkan Melalui Tanah Pada Siswa Sekolah Dasar di Distrik Arso Kabupaten Keerom, Papua. *Media Litbangkes*. 2015;25(1):1–14.
11. Kamila AD, Margawati A, Nuryanto. Hubungan Kecacingan dengan Status Gizi dan Prestasi Belajar pada Anak Sekolah Dasar Kelas IV dan VI di Kelurahan Bandarharjo Semarang. *J Nutr Coll*. 2018;7(2):77–83.
12. Mahmudah U. Hubungan Sanitasi Lingkungan Rumah Terhadap Kejadian Infeksi Kecacingan Pada Anak Sekolah Dasar. *J Kesehat*. 2017;10(1):32–9.
13. Suprpti, Hapsari D. Status Gizi Balita Berdasarkan Kondisi Lingkungan dan Status Ekonomi (Data Riskesdas 2007). *J Ekol Kesehat*. 2007;10(2):103–13.
14. Azizaturridha A, Istiana, Hayatie L. Pengaruh infeksi kecacingan terhadap status gizi pada anak di SDN 2 Barabai Darat Kalimantan Selatan Tahun 2015. *Berk Kedokt*. 2016;12(2):165–73.
15. Indriyati L, Waris L, Luciasari E. Kerugian Finansial Akibat Kecacingan: Studi di Kabupaten Nunukan. *Penelit Gizi Makan*. 2014;37(2):155–60.

ORIGINALITY REPORT

16%

SIMILARITY INDEX

7%

INTERNET SOURCES

12%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Vierto Irennius Girsang, Friska Vaulina Manik, Seri Asnawati Munthe. "Determinants of roundworms among primary school students in North Sumatera Province, Indonesia", Malahayati International Journal of Nursing and Health Science, 2019
Publication 3%
- 2 "Proceedings of the Andalas International Public Health Conference 2017", BMC Public Health, 2017
Publication 2%
- 3 journal2.uad.ac.id
Internet Source 2%
- 4 Chairil Anwar, Saraswati Annisa, Dalilah Dalilah, Novrikasari Novrikasari. "The Relationship Between Soil Transmitted Helminthes (STH) Infection and Nutritional Status in Students of State Elementary School Number (SDN) 200 Palembang Indonesia", BioScientia Medicina, 2018
Publication 2%

5

Sri Lestari Ramadhani Nasution, Ali Napiah Nasution, Suhartomi, Sri Wahyuni Nasution. "The Effect of Habits on Wearing Footwear And hand washing after playing on the ground Against Worms in Primary School Al-Wasliyah In Medan Deli", Journal of Physics: Conference Series, 2019

Publication

1%

6

www.who.int

Internet Source

1%

7

Dewi Masyithah Darlan, Tania Silvia Alexandra, Zaimah Z. Tala. "Soil Transmitted Helminth Infections in Medan: a cross-sectional study of the correlation between the infection and nutritional status among elementary school children", Family Medicine & Primary Care Review, 2017

Publication

1%

8

Liena Sofiana, Erni Gustina, Luthva Luviandani Pratiwi. "HUBUNGAN ANTARA KECACINGAN DENGAN ANEMIA PADA ANAK SEKOLAH DASAR DI WILAYAH KERJA PUSKESMAS MOYUDAN, SLEMAN", Medika Respati : Jurnal Ilmiah Kesehatan, 2019

Publication

1%

9

es.slideshare.net

Internet Source

1%

10	Submitted to University of Houston System Student Paper	1%
11	Submitted to Universitas Sumatera Utara Student Paper	1%
12	positori.usu.ac.id Internet Source	1%
13	Submitted to iGroup Student Paper	<1%
14	jurnal.uns.ac.id Internet Source	<1%
15	eprints.undip.ac.id Internet Source	<1%
16	www.neliti.com Internet Source	<1%
17	Submitted to University of Leeds Student Paper	<1%
18	Septian Emma Dwi Jatmika, Muchsin Maulana, Kuntoro Kuntoro, Santi Martini, Beni Setya Anjani. "The Readiness of Smokers to Quit Smoking", Global Journal of Health Science, 2018 Publication	<1%

Exclude quotes On

Exclude matches Off

Exclude bibliography On

9. 933-2593-1-PB

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
