

CYP2E1 polymorphism, acetylator profiles and drug-induced liver injury incidence of Indonesian tuberculosis patients


By ENDANG DARMAWAN



4
Available online at www.sciencedirect.com

ScienceDirect

journal homepage: <http://www.journals.elsevier.com/indian-journal-of-tuberculosis/>



Original Article

8
CYP2E1 polymorphism, acetylator profiles and drug-induced liver injury incidence of Indonesian tuberculosis patients

Dyah A. Perwitasari^{a,*}, Lalu M. Irham^a, Endang Darmawan^a, Ully A. Mulyani^b, J. Atthobari^c

¹⁸
¹⁵
¹³
¹⁴
¹⁵
¹⁶
¹⁷
¹⁸
¹⁹
²⁰
²¹
²²
²³
²⁴
²⁵
²⁶
²⁷
²⁸
²⁹
³⁰
³¹
³²
³³
³⁴
³⁵
³⁶
³⁷
³⁸
³⁹
⁴⁰
⁴¹
⁴²
⁴³
⁴⁴
⁴⁵
⁴⁶
⁴⁷
⁴⁸
⁴⁹
⁵⁰
⁵¹
⁵²
⁵³
⁵⁴
⁵⁵
⁵⁶
⁵⁷
⁵⁸
⁵⁹
⁶⁰
⁶¹
⁶²
⁶³
⁶⁴
⁶⁵
⁶⁶
⁶⁷
⁶⁸
⁶⁹
⁷⁰
⁷¹
⁷²
⁷³
⁷⁴
⁷⁵
⁷⁶
⁷⁷
⁷⁸
⁷⁹
⁸⁰
⁸¹
⁸²
⁸³
⁸⁴
⁸⁵
⁸⁶
⁸⁷
⁸⁸
⁸⁹
⁹⁰
⁹¹
⁹²
⁹³
⁹⁴
⁹⁵
⁹⁶
⁹⁷
⁹⁸
⁹⁹
¹⁰⁰
¹⁰¹
¹⁰²
¹⁰³
¹⁰⁴
¹⁰⁵
¹⁰⁶
¹⁰⁷
¹⁰⁸
¹⁰⁹
¹¹⁰
¹¹¹
¹¹²
¹¹³
¹¹⁴
¹¹⁵
¹¹⁶
¹¹⁷
¹¹⁸
¹¹⁹
¹²⁰
¹²¹
¹²²
¹²³
¹²⁴
¹²⁵
¹²⁶
¹²⁷
¹²⁸
¹²⁹
¹³⁰
¹³¹
¹³²
¹³³
¹³⁴
¹³⁵
¹³⁶
¹³⁷
¹³⁸
¹³⁹
¹⁴⁰
¹⁴¹
¹⁴²
¹⁴³
¹⁴⁴
¹⁴⁵
¹⁴⁶
¹⁴⁷
¹⁴⁸
¹⁴⁹
¹⁵⁰
¹⁵¹
¹⁵²
¹⁵³
¹⁵⁴
¹⁵⁵
¹⁵⁶
¹⁵⁷
¹⁵⁸
¹⁵⁹
¹⁶⁰
¹⁶¹
¹⁶²
¹⁶³
¹⁶⁴
¹⁶⁵
¹⁶⁶
¹⁶⁷
¹⁶⁸
¹⁶⁹
¹⁷⁰
¹⁷¹
¹⁷²
¹⁷³
¹⁷⁴
¹⁷⁵
¹⁷⁶
¹⁷⁷
¹⁷⁸
¹⁷⁹
¹⁸⁰
¹⁸¹
¹⁸²
¹⁸³
¹⁸⁴
¹⁸⁵
¹⁸⁶
¹⁸⁷
¹⁸⁸
¹⁸⁹
¹⁹⁰
¹⁹¹
¹⁹²
¹⁹³
¹⁹⁴
¹⁹⁵
¹⁹⁶
¹⁹⁷
¹⁹⁸
¹⁹⁹
²⁰⁰
²⁰¹
²⁰²
²⁰³
²⁰⁴
²⁰⁵
²⁰⁶
²⁰⁷
²⁰⁸
²⁰⁹
²¹⁰
²¹¹
²¹²
²¹³
²¹⁴
²¹⁵
²¹⁶
²¹⁷
²¹⁸
²¹⁹
²²⁰
²²¹
²²²
²²³
²²⁴
²²⁵
²²⁶
²²⁷
²²⁸
²²⁹
²³⁰
²³¹
²³²
²³³
²³⁴
²³⁵
²³⁶
²³⁷
²³⁸
²³⁹
²⁴⁰
²⁴¹
²⁴²
²⁴³
²⁴⁴
²⁴⁵
²⁴⁶
²⁴⁷
²⁴⁸
²⁴⁹
²⁵⁰
²⁵¹
²⁵²
²⁵³
²⁵⁴
²⁵⁵
²⁵⁶
²⁵⁷
²⁵⁸
²⁵⁹
²⁶⁰
²⁶¹
²⁶²
²⁶³
²⁶⁴
²⁶⁵
²⁶⁶
²⁶⁷
²⁶⁸
²⁶⁹
²⁷⁰
²⁷¹
²⁷²
²⁷³
²⁷⁴
²⁷⁵
²⁷⁶
²⁷⁷
²⁷⁸
²⁷⁹
²⁸⁰
²⁸¹
²⁸²
²⁸³
²⁸⁴
²⁸⁵
²⁸⁶
²⁸⁷
²⁸⁸
²⁸⁹
²⁹⁰
²⁹¹
²⁹²
²⁹³
²⁹⁴
²⁹⁵
²⁹⁶
²⁹⁷
²⁹⁸
²⁹⁹
³⁰⁰
³⁰¹
³⁰²
³⁰³
³⁰⁴
³⁰⁵
³⁰⁶
³⁰⁷
³⁰⁸
³⁰⁹
³¹⁰
³¹¹
³¹²
³¹³
³¹⁴
³¹⁵
³¹⁶
³¹⁷
³¹⁸
³¹⁹
³²⁰
³²¹
³²²
³²³
³²⁴
³²⁵
³²⁶
³²⁷
³²⁸
³²⁹
³³⁰
³³¹
³³²
³³³
³³⁴
³³⁵
³³⁶
³³⁷
³³⁸
³³⁹
³⁴⁰
³⁴¹
³⁴²
³⁴³
³⁴⁴
³⁴⁵
³⁴⁶
³⁴⁷
³⁴⁸
³⁴⁹
³⁵⁰
³⁵¹
³⁵²
³⁵³
³⁵⁴
³⁵⁵
³⁵⁶
³⁵⁷
³⁵⁸
³⁵⁹
³⁶⁰
³⁶¹
³⁶²
³⁶³
³⁶⁴
³⁶⁵
³⁶⁶
³⁶⁷
³⁶⁸
³⁶⁹
³⁷⁰
³⁷¹
³⁷²
³⁷³
³⁷⁴
³⁷⁵
³⁷⁶
³⁷⁷
³⁷⁸
³⁷⁹
³⁸⁰
³⁸¹
³⁸²
³⁸³
³⁸⁴
³⁸⁵
³⁸⁶
³⁸⁷
³⁸⁸
³⁸⁹
³⁹⁰
³⁹¹
³⁹²
³⁹³
³⁹⁴
³⁹⁵
³⁹⁶
³⁹⁷
³⁹⁸
³⁹⁹
⁴⁰⁰
⁴⁰¹
⁴⁰²
⁴⁰³
⁴⁰⁴
⁴⁰⁵
⁴⁰⁶
⁴⁰⁷
⁴⁰⁸
⁴⁰⁹
⁴¹⁰
⁴¹¹
⁴¹²
⁴¹³
⁴¹⁴
⁴¹⁵
⁴¹⁶
⁴¹⁷
⁴¹⁸
⁴¹⁹
⁴²⁰
⁴²¹
⁴²²
⁴²³
⁴²⁴
⁴²⁵
⁴²⁶
⁴²⁷
⁴²⁸
⁴²⁹
⁴³⁰
⁴³¹
⁴³²
⁴³³
⁴³⁴
⁴³⁵
⁴³⁶
⁴³⁷
⁴³⁸
⁴³⁹
⁴⁴⁰
⁴⁴¹
⁴⁴²
⁴⁴³
⁴⁴⁴
⁴⁴⁵
⁴⁴⁶
⁴⁴⁷
⁴⁴⁸
⁴⁴⁹
⁴⁵⁰
⁴⁵¹
⁴⁵²
⁴⁵³
⁴⁵⁴
⁴⁵⁵
⁴⁵⁶
⁴⁵⁷
⁴⁵⁸
⁴⁵⁹
⁴⁶⁰
⁴⁶¹
⁴⁶²
⁴⁶³
⁴⁶⁴
⁴⁶⁵
⁴⁶⁶
⁴⁶⁷
⁴⁶⁸
⁴⁶⁹
⁴⁷⁰
⁴⁷¹
⁴⁷²
⁴⁷³
⁴⁷⁴
⁴⁷⁵
⁴⁷⁶
⁴⁷⁷
⁴⁷⁸
⁴⁷⁹
⁴⁸⁰
⁴⁸¹
⁴⁸²
⁴⁸³
⁴⁸⁴
⁴⁸⁵
⁴⁸⁶
⁴⁸⁷
⁴⁸⁸
⁴⁸⁹
⁴⁹⁰
⁴⁹¹
⁴⁹²
⁴⁹³
⁴⁹⁴
⁴⁹⁵
⁴⁹⁶
⁴⁹⁷
⁴⁹⁸
⁴⁹⁹
⁵⁰⁰
⁵⁰¹
⁵⁰²
⁵⁰³
⁵⁰⁴
⁵⁰⁵
⁵⁰⁶
⁵⁰⁷
⁵⁰⁸
⁵⁰⁹
⁵¹⁰
⁵¹¹
⁵¹²
⁵¹³
⁵¹⁴
⁵¹⁵
⁵¹⁶
⁵¹⁷
⁵¹⁸
⁵¹⁹
⁵²⁰
⁵²¹
⁵²²
⁵²³
⁵²⁴
⁵²⁵
⁵²⁶
⁵²⁷
⁵²⁸
⁵²⁹
⁵³⁰
⁵³¹
⁵³²
⁵³³
⁵³⁴
⁵³⁵
⁵³⁶
⁵³⁷
⁵³⁸
⁵³⁹
⁵⁴⁰
⁵⁴¹
⁵⁴²
⁵⁴³
⁵⁴⁴
⁵⁴⁵
⁵⁴⁶
⁵⁴⁷
⁵⁴⁸
⁵⁴⁹
⁵⁵⁰
⁵⁵¹
⁵⁵²
⁵⁵³
⁵⁵⁴
⁵⁵⁵
⁵⁵⁶
⁵⁵⁷
⁵⁵⁸
⁵⁵⁹
⁵⁶⁰
⁵⁶¹
⁵⁶²
⁵⁶³
⁵⁶⁴
⁵⁶⁵
⁵⁶⁶
⁵⁶⁷
⁵⁶⁸
⁵⁶⁹
⁵⁷⁰
⁵⁷¹
⁵⁷²
⁵⁷³
⁵⁷⁴
⁵⁷⁵
⁵⁷⁶
⁵⁷⁷
⁵⁷⁸
⁵⁷⁹
⁵⁸⁰
⁵⁸¹
⁵⁸²
⁵⁸³
⁵⁸⁴
⁵⁸⁵
⁵⁸⁶
⁵⁸⁷
⁵⁸⁸
⁵⁸⁹
⁵⁹⁰
⁵⁹¹
⁵⁹²
⁵⁹³
⁵⁹⁴
⁵⁹⁵
⁵⁹⁶
⁵⁹⁷
⁵⁹⁸
⁵⁹⁹
⁶⁰⁰
⁶⁰¹
⁶⁰²
⁶⁰³
⁶⁰⁴
⁶⁰⁵
⁶⁰⁶
⁶⁰⁷
⁶⁰⁸
⁶⁰⁹
⁶¹⁰
⁶¹¹
⁶¹²
⁶¹³
⁶¹⁴
⁶¹⁵
⁶¹⁶
⁶¹⁷
⁶¹⁸
⁶¹⁹
⁶²⁰
⁶²¹
⁶²²
⁶²³
⁶²⁴
⁶²⁵
⁶²⁶
⁶²⁷
⁶²⁸
⁶²⁹
⁶³⁰
⁶³¹
⁶³²
⁶³³
⁶³⁴
⁶³⁵
⁶³⁶
⁶³⁷
⁶³⁸
⁶³⁹
⁶⁴⁰
⁶⁴¹
⁶⁴²
⁶⁴³
⁶⁴⁴
⁶⁴⁵
⁶⁴⁶
⁶⁴⁷
⁶⁴⁸
⁶⁴⁹
⁶⁵⁰
⁶⁵¹
⁶⁵²
⁶⁵³
⁶⁵⁴
⁶⁵⁵
⁶⁵⁶
⁶⁵⁷
⁶⁵⁸
⁶⁵⁹
⁶⁶⁰
⁶⁶¹
⁶⁶²
⁶⁶³
⁶⁶⁴
⁶⁶⁵
⁶⁶⁶
⁶⁶⁷
⁶⁶⁸
⁶⁶⁹
⁶⁷⁰
⁶⁷¹
⁶⁷²
⁶⁷³
⁶⁷⁴
⁶⁷⁵
⁶⁷⁶
⁶⁷⁷
⁶⁷⁸
⁶⁷⁹
⁶⁸⁰
⁶⁸¹
⁶⁸²
⁶⁸³
⁶⁸⁴
⁶⁸⁵
⁶⁸⁶
⁶⁸⁷
⁶⁸⁸
⁶⁸⁹
⁶⁹⁰
⁶⁹¹
⁶⁹²
⁶⁹³
⁶⁹⁴
⁶⁹⁵
⁶⁹⁶
⁶⁹⁷
⁶⁹⁸
⁶⁹⁹
⁷⁰⁰
⁷⁰¹
⁷⁰²
⁷⁰³
⁷⁰⁴
⁷⁰⁵
⁷⁰⁶
⁷⁰⁷
⁷⁰⁸
⁷⁰⁹
⁷¹⁰
⁷¹¹
⁷¹²
⁷¹³
⁷¹⁴
⁷¹⁵
⁷¹⁶
⁷¹⁷
⁷¹⁸
⁷¹⁹
⁷²⁰
⁷²¹
⁷²²
⁷²³
⁷²⁴
⁷²⁵
⁷²⁶
⁷²⁷
⁷²⁸
⁷²⁹
⁷³⁰
⁷³¹
⁷³²
⁷³³
⁷³⁴
⁷³⁵
⁷³⁶
⁷³⁷
⁷³⁸
⁷³⁹
⁷⁴⁰
⁷⁴¹
⁷⁴²
⁷⁴³
⁷⁴⁴
⁷⁴⁵
⁷⁴⁶
⁷⁴⁷
⁷⁴⁸
⁷⁴⁹
⁷⁵⁰
⁷⁵¹
⁷⁵²
⁷⁵³
⁷⁵⁴
⁷⁵⁵
⁷⁵⁶
⁷⁵⁷
⁷⁵⁸
⁷⁵⁹
⁷⁶⁰
⁷⁶¹
⁷⁶²
⁷⁶³
⁷⁶⁴
⁷⁶⁵
⁷⁶⁶
⁷⁶⁷
⁷⁶⁸
⁷⁶⁹
⁷⁷⁰
⁷⁷¹
⁷⁷²
⁷⁷³
⁷⁷⁴
⁷⁷⁵
⁷⁷⁶
⁷⁷⁷
⁷⁷⁸
⁷⁷⁹
⁷⁸⁰
⁷⁸¹
⁷⁸²
⁷⁸³
⁷⁸⁴
⁷⁸⁵
⁷⁸⁶
⁷⁸⁷
⁷⁸⁸
⁷⁸⁹
⁷⁹⁰
⁷⁹¹
⁷⁹²
⁷⁹³
⁷⁹⁴
⁷⁹⁵
⁷⁹⁶
⁷⁹⁷
⁷⁹⁸
⁷⁹⁹
⁸⁰⁰
⁸⁰¹
⁸⁰²
⁸⁰³
⁸⁰⁴
⁸⁰⁵
⁸⁰⁶
⁸⁰⁷
⁸⁰⁸
⁸⁰⁹
⁸¹⁰
⁸¹¹
⁸¹²
⁸¹³
⁸¹⁴
⁸¹⁵
⁸¹⁶
⁸¹⁷
⁸¹⁸
⁸¹⁹
⁸²⁰
⁸²¹
⁸²²
⁸²³
⁸²⁴
⁸²⁵
⁸²⁶
⁸²⁷
⁸²⁸
⁸²⁹
⁸³⁰
⁸³¹
⁸³²
⁸³³
⁸³⁴
⁸³⁵
⁸³⁶
⁸³⁷
⁸³⁸
⁸³⁹
⁸⁴⁰
⁸⁴¹
⁸⁴²
⁸⁴³
⁸⁴⁴
⁸⁴⁵
⁸⁴⁶
⁸⁴⁷
⁸⁴⁸
⁸⁴⁹
⁸⁵⁰
⁸⁵¹
⁸⁵²
⁸⁵³
⁸⁵⁴
⁸⁵⁵
⁸⁵⁶
⁸⁵⁷
⁸⁵⁸
⁸⁵⁹
⁸⁶⁰
⁸⁶¹
⁸⁶²
⁸⁶³
⁸⁶⁴
⁸⁶⁵
⁸⁶⁶
⁸⁶⁷
⁸⁶⁸
⁸⁶⁹
⁸⁷⁰
⁸⁷¹
⁸⁷²
⁸⁷³
⁸⁷⁴
⁸⁷⁵
⁸⁷⁶
⁸⁷⁷
⁸⁷⁸
⁸⁷⁹
⁸⁸⁰
⁸⁸¹
⁸⁸²
⁸⁸³
⁸⁸⁴
⁸⁸⁵
⁸⁸⁶
⁸⁸⁷
⁸⁸⁸
⁸⁸⁹
⁸⁹⁰
⁸⁹¹
⁸⁹²
⁸⁹³
⁸⁹⁴
⁸⁹⁵
⁸⁹⁶
⁸⁹⁷
⁸⁹⁸
⁸⁹⁹
⁹⁰⁰
⁹⁰¹
⁹⁰²
⁹⁰³
⁹⁰⁴
⁹⁰⁵
⁹⁰⁶
⁹⁰⁷
⁹⁰⁸
⁹⁰⁹
⁹¹⁰
⁹¹¹
⁹¹²
⁹¹³
⁹¹⁴
⁹¹⁵
⁹¹⁶
⁹¹⁷
⁹¹⁸
⁹¹⁹
⁹²⁰
⁹²¹
⁹²²
⁹²³
⁹²⁴
⁹²⁵
⁹²⁶
⁹²⁷
⁹²⁸
⁹²⁹
⁹³⁰
⁹³¹
⁹³²
⁹³³
⁹³⁴
⁹³⁵
⁹³⁶
⁹³⁷
⁹³⁸
⁹³⁹
⁹⁴⁰
⁹⁴¹
⁹⁴²
⁹⁴³
⁹⁴⁴
⁹⁴⁵
⁹⁴⁶
⁹⁴⁷
⁹⁴⁸
⁹⁴⁹
⁹⁵⁰
⁹⁵¹
⁹⁵²
⁹⁵³
⁹⁵⁴
⁹⁵⁵
⁹⁵⁶
⁹⁵⁷
⁹⁵⁸
⁹⁵⁹
⁹⁶⁰
⁹⁶¹
⁹⁶²
⁹⁶³
⁹⁶⁴
⁹⁶⁵
⁹⁶⁶
⁹⁶⁷
⁹⁶⁸
⁹⁶⁹
⁹⁷⁰
⁹⁷¹
⁹⁷²
⁹⁷³
⁹⁷⁴
⁹⁷⁵
⁹⁷⁶
⁹⁷⁷
⁹⁷⁸
⁹⁷⁹
⁹⁸⁰
⁹⁸¹
⁹⁸²
⁹⁸³
⁹⁸⁴
⁹⁸⁵
⁹⁸⁶
⁹⁸⁷
⁹⁸⁸
⁹⁸⁹
⁹⁹⁰
⁹⁹¹
⁹⁹²
⁹⁹³
⁹⁹⁴
⁹⁹⁵
⁹⁹⁶
⁹⁹⁷
⁹⁹⁸
⁹⁹⁹
¹⁰⁰⁰

^a Postgraduate Program of Clinical Pharmacy, Faculty of Pharmacy, Ahmad Dahlan University, Jl prof Dr Soepomo, Janturan 55164, Jakarta, Indonesia

^b National Institute for Health Research and Development, Jl. Percetakan Negara 29, Jakarta Pusat 10560, Jakarta, Indonesia

^c Faculty of Medicine, Universitas Gadjah Mada, Jl. Farmako Sekip 55281, Yogyakarta, Indonesia

14
ARTICLE INFO

Article history:

Received 14 January 2016

Accepted 2 August 2016

Available online xxx

Keywords:

Hepatotoxicity
Acetylator status
INH
Indonesian
CYP2E1

ABSTRACT

Objective: A polymorphism of CYP2E1 may be directly associated with the development of INH hepatotoxicity. We conducted this study to evaluate the association between polymorphisms of CYP2E1, Isoniazid (INH) concentration and the acetylator status of INH in cases of Indonesian tuberculosis patients with drug-induced liver disease (DILI).

Methods: We conducted our study with a cohort design consisting of 55 Indonesian adult tuberculosis (TB) patients. Acetylating phenotypes were studied in using the metabolic ratio of plasma AchZ/HZ. DILI was defined using CTCAV version 4.0. The allelic and genotypic frequency distributions of CYP2E1 rs 3813867 were studied using the polymerase chain reaction – amplification refractory mutation system (ARMS) methodology.

Results: Patients with an INH concentration of more than 7 µg/mL showed a higher risk of developing DILI when compared with patients who showed a therapeutic range of 3–6 µg/mL INH (OR: 1.3, 95% CI: 0.2–8.2). Slow acetylators had a higher incidence of DILI when compared with rapid acetylators (OR: 4.6, 95% CI: 1.3–15.9). Meanwhile, subjects with GC had a higher risk of DILI incidence (OR: 4.3, 95% CI: 0.8–24.4).

Conclusion: Our study shows that polymorphisms of CYP2E1 and slow acetylator may have role in the DILI incidence.

© 2016 Tuberculosis Association of India. Published by Elsevier B.V. All rights reserved.

4
1. Introduction

In 2015, the tuberculosis (TB) prevalence in the world reached 42% lower than in 1990. In 2014, of the 9.6 million of TB incidence, 58% were from South-East Asia and Western-Pacific

24
region. Still, India, Indonesia and China had the largest number of cases from global total number (23%, 10% and 10%, respectively). The treatment success rate for newly diagnosed TB patients reached 86% in 2013 which was still sustained since 2015.¹ Treatment success is the key outcome of TB burden reduction. Some factors may influence the

* Corresponding author. Tel.: +62 274 563515; fax: +62 274 563515.

7 E-mail address: diahperwitasari2003@yahoo.com (D.A. Perwitasari).

<http://dx.doi.org/10.1016/j.ijtb.2016.08.001>

0019-5707/© 2016 Tuberculosis Association of India. Published by Elsevier B.V. All rights reserved.

treatment success of TB, like hospital facilities, staffing education, population covered and centralized or decentralized health care system.²

The TB treatment regimen is a four-drug combination consisting of isoniazid (INH), rifampicin (R), pyrazinamide (Z), and ethambutol (E). This anti-tuberculosis therapy can cause adverse drug events. The most commonly reported adverse drug event of anti-tuberculosis drugs is hepatotoxicity.³⁻⁵ Among these compounds, isoniazid (INH) strongly associated with anti-tuberculosis drug-induced hepatotoxicity (ATDH) or drug-induced liver injury (DILI). Previous reports showed INH induced liver injury or hepatotoxicity in approximately 5% of patients.⁴⁻⁶ Other studies showed the development of ATDH in approximately 1 to 36% of patients.^{7,8} INH is metabolized by N-acetyltransferase-2 (NAT-2) into acetylisoniazid. Next, the metabolite is hydrolysed to acetylhydrazine (AchZ) and hydrazine (HZ) by hepatic N-acetyltransferase-2 (NAT-2).⁹ These compounds are then oxidized by cytochrome P450 2E1 (CYP2E1) to form intermediate hepatotoxins.^{10,11}

There are large variations in the metabolism of the anti-tuberculosis drug isoniazid, including polymorphisms of the NAT-2 gene. This enzyme is markedly decreased in the livers of slow acetylators (SA). The elimination of INH follows a bimodal or trimodal distribution consisting of slow (SA), intermediate (IA) and rapid acetylators (RA). There is a strong correlation between these phenotypes and NAT2 genotypes in Caucasians.¹¹⁻¹³ Previous studies have reported inconsistent results on whether slow or rapid acetylators are a risk factor for INH-induced hepatotoxicity.^{9,13-15}

The objective of this study was to determine the association between CYP2E1 polymorphisms and the development of DILI in Indonesians and determine the genotypes and phenotypes related to a change in the risk of DILI.

2. Materials and methods

2.1. Patients

This study used a cohort design. A total of 55 Indonesian adult patients with newly diagnosed TB at 20 Public Health Centres in the Provinces of Yogyakarta (10 Public Health Centres) and Lampung (10 Public Health Centres) were enrolled. The patient's recruitment was conducted from January until December 2013. The inclusion criteria included all adult patients (age >18 years) newly diagnosed as a pulmonary TB patient who were receiving category 1 TB medications and signed the informed consent for study participation. Exclusion criteria consisted of TB patients with HIV/AIDS, an AST and ALT two-fold higher than normal baseline concentrations, hepatitis or a history of hepatitis, a haemoglobin concentration <8 mg/dL, cessation of medication for more than 2 weeks, a history of kidney diseases, or refusal of blood sampling procedures and patients who refused to participate in this study.

All patients received a standard TB treatment for the first 2 months, including oral INH (300 mg), rifampicin (600 mg), pyrazinamide (20 mg/kg body weight), and ethambutol (800 mg). After 2 months of treatment, the patients were given INH and rifampicin for an additional 4 months. The total

duration of anti-tuberculosis treatment was 6 months. Serum alanine transaminase (ALT), aspartate transaminase (AST) and alkaline phosphatase were measured before the anti-tuberculosis therapy and then monthly until the end of treatment.

Genomic DNA was extracted from blood samples using a DNA Gene JET Genomic DNA Purification Kit (Thermo Scientific®, Nutrilab Pratama, Jakarta, Indonesia) according to the manufacturer's instructions. Analysis and Identification of SNP (-1055) C/G (Rs 3813867) was completed using an amplification refractory mutation system (ARMS) method. Two primer sets of CYP2E1 (Rs 3813867) were used: forward I "5-GTACAAAATTGCAACCTATG-3" to detect the CYP2E1 polymorphic gene (F-primer - 1), forward II "5-GTAGAAAATTGCAACCTATG-3" to confirm the normal gene fragment and (F-primer - 2) reverse "5-ATCTTGCTTTGTTGATCCC-3" (Genebank accession number P05181). The cycling conditions involved preliminary denaturation at 95 °C for 2 min, followed by 35 cycles of denaturation at 95 °C for 30 s, annealing at 55 °C for 30 s, and elongation at 72 °C for 30 s, followed by a final elongation step at 72 °C for 2 min. The PCR products of single nucleotide polymorphism (SNP) (-1055) C > G (rs3813867) were obtained for 55 subjects. The results showed a band at 230 bp. After electrophoresis using a 2% agarose gel, samples were observed under UV light and imaged. Samples with homozygote CC (cytosine-cytosine) appeared using primer set Forward 1-Reverse (F1-R), samples with homozygote GG (guanine-guanine) appeared using primer set Forward 2-Reverse (F2-R), and samples with homozygote CG (cytosine-guanine) appeared using both primer sets F1-R and F2-R.

Venous blood samples were collected 2 h after drug administration. Determination of acetylator status was made using the AchZ/HZ ratio with a cut-off point of 15.00. A slow acetylator (SA) was defined as an AchZ/HZ ratio ≤ 15.00, and a rapid acetylator (RA) was defined as an AchZ/HZ ratio > 15.¹⁶ The AST and ALT levels were measured with an automatic chemical analyser. Isoniazid and its metabolites, HZ and AchZ, were measured using HPLC. Statistical significance was analyzed using correlation tests. DILI incidence was defined as an ALT and/or AST level above the upper limit normal value (ULN) listed in Common Toxicity Criteria for Adverse Events version 4.0 (CTCAE v. 4). The normal values of ALT and AST are 0-50 mg/dL and 0-33 mg/dL, respectively, for males. The normal values for ALT and AST are 0-34 mg/dL and 0-27 mg/dL, respectively, for females. Increased ALT and/or AST or an increased ALP were categorized as grade 1+, (>1.0-2.5 × ULN), grade 2+ (>2.5-5.0 × ULN), grade 3+ (>5.0-20 × ULN), or grade 4+ (>20.0 × ULN).¹⁷

This study was approved by the National Ethics Committee of the National Institute of Health Research and Development, Ministry of Health, Republic of Indonesia, and written informed consent was obtained from all participants.

2.2. Statistical methods

Statistical analysis was conducted using the SPSS® statistical package, version 16.0. The odds ratio (OR) and confidence interval (CI) were calculated using binary logistic regression analysis. This analysis evaluated the risk and association of DILI, INH and acetylator status. Allele frequencies were

calculated, and the agreement between allele frequencies and Hardy-Weinberg equilibrium was tested using a chi-square test (d.f. = 1) for each locus.

3. Results

This study recruited 55 tuberculosis patients. Table 1 shows the demographics of the patients. There were 9 patients (16.36%) genotyped as CC (c1/c1), 29 patients (52.73%) genotyped as CG (c1/c2) and 17 patients (30.90%) genotyped as GG (c2/c2). The genotype frequencies were not significantly different from those predicted by the Hardy-Weinberg equation ($P > 0.05$). According to sex and age differences, the CG genotype was predominant over CC and GG. There were also no significant differences between ALT and AST means at baseline between males and females with different genotypes ($P > 0.05$).

Table 2 shows the association between INH concentration, acetylator status and DILI incidence. Among the 55 tuberculosis patients, there were 25 patients (45%) with DILI. There was no significant difference between INH serum concentrations and DILI incidence. However, patients with INH serum concentrations more than 7 $\mu\text{g/mL}$ had a greater tendency to develop DILI when compared with subjects who had normal (3–6 $\mu\text{g/mL}$) and low (3 $\mu\text{g/mL}$) INH concentrations (OR: 1.3, 95% CI: 0.2–8.2). There was a significant association between acetylator status and DILI incidence. A total of 17 patients (30.9%) were identified as RAs and the other 38 patients (69.09%) were SAs. SAs showed a higher risk of developing DILI when compared with RAs (OR: 4.6, 95% CI: 1.3–15.9).

Table 3 shows the distribution of INH serum concentration and acetylator status based on genotype. There were 10 patients (16.36%) genotyped as CC (c1/c1), 29 patients (52.73%) genotyped as CG (c1/c2) and 16 patients (30.90%) genotyped as GG (c2/c2). There were no significant differences in the frequencies of c1/c1, c1/c2, and c2/c2 genotypes between the patients with and without DILI. The heterozygous c1/c2 variant showed a higher DILI incidence when compared with the homozygous c2/c2 variant (OR: 4.3, 95% CI: 0.8–24.4 and OR: 2.5, 95% CI: 0.4–15.5, respectively). We did the stratified analysis according to the acetylator data to the INH serum

Table 1 – Demographics of Indonesian tuberculosis patients according to genotype.

	Genotype			P value
	CC N (%)	CG N (%)	GG N (%)	
Sex				
Male	4 (44.5)	15 (53.6)	10 (55.5)	
Female	5 (55.5)	13 (46.4)	8 (44.5)	
Total	9 (100.0)	28 (100.0)	18 (100.0)	
Age (years)				
18–35	2 (22.2)	20 (71.4)	10 (55.6)	
36–55	5 (55.5)	3 (10.7)	6 (33.3)	
>55	3 (33.3)	5 (17.9)	2 (11.1)	
Total	9 (100.0)	28 (100.0)	18 (100.0)	
	Mean \pm SD	Mean \pm SD	Mean \pm SD	
ALT (U/L) Baseline				
Male	21.8 \pm 15.98	23.2 \pm 16.81	20.8 \pm 11.72	0.07
Female	16.4 \pm 6.69	15.3 \pm 12.15	14.6 \pm 3.99	
AST (U/L) Baseline				
Male	19.5 \pm 9.03	24.4 \pm 13.54	23.4 \pm 7.19	0.35
Female	24.8 \pm 15.89	20.6 \pm 9.78	16.2 \pm 5.47	

concentration as the determinant; however, we did not find significant association ($P > 0.05$, data was not shown). Furthermore, the binary logistic regression analysis was performed; however we also did not find significant association ($P > 0.05$; data was not shown). After performing bivariate analysis, there was interesting pattern of results in this study, which are: patients who did not experience DILI had INH serum concentration between 0 and 7 ng/mL (OR; 1.8; 95% CI: 0.4–8.2). After stratified by genotyping, we found that the patients with GG genotype have more possibilities to experience DILI (OR: 0.7; 95% CI: 0.4–1.1). We conducted the stratified analysis according to acetylator status with DILI status as the determinant of INH serum concentration. We found that patients without DILI have lower serum concentration INH in slow acetylator (OR: 1.6; 95% CI: 0.3–7.6).

An INH serum concentration greater than 7 $\mu\text{g/mL}$ was found to be predominant in all genotypes. This pattern was also found for acetylator status. A SA status is dominant in all

Table 2 – Association between INH concentration, acetylator status and DILI incidence.

Variable	With DILI N = 25	Without DILI N = 30	Total N	OR (CI 95%)	P-value	
INH concentration ($\mu\text{g/mL}$)	0–3	4	4	8	Reference	
	3–6	3	4	7	0.6 (0.03–8.9)	0.63
	>7	18	22	40	1.2 (0.2–8.2)	0.83
Acetylators status	Rapid acetylators	12	5	17	Reference	
	Slow acetylators	13	25	38	4.6 (1.3–15.9)	0.02*
Genotype CYP2E1	CC(c1/c1)	2	7	9	Reference	0.09
	CG(c1/c2)	16	13	29	4.3 (0.8–24.4)	
Genotype CYP2E1	CC(c1/c1)	2	7	9	Reference	0.09
	GG(c2/c2)	7	10	17	2.4 (0.5–15.5)	

OR: odds ratio; CI: confidence interval.

* Significant difference ($P < 0.05$).

Please cite this article in press as: Perwitasari DA, et al. CYP2E1 polymorphism, acetylator profiles and drug-induced liver injury incidence of Indonesian tuberculosis patients, *Indian J Tuberc.* (2016), <http://dx.doi.org/10.1016/j.ijtb.2016.08.001>

Table 3 – Distribution of INH serum concentration and acetylator status based on genotype.

Variable		CYP2E1 genotype			P-value
		c1/c1 N = 10 (%)	c1/c2 N = 29 (%)	c2/c2 N = 16 (%)	
INH serum concentration ($\mu\text{g/mL}$)	0-3	1 (10.0)	4 (13.8)	3 (18.8)	0.70
	>3-6	3 (30.0)	2 (6.9)	2 (12.5)	0.53
	>7	6 (60.0)	23 (79.9)	11 (68.8)	0.83
Acetylator Status	SA	8 (80.0)	19 (65.5)	11 (68.8)	0.65
	RA	2 (20.0)	10 (34.5)	5 (31.3)	0.43

genotypes. The INH serum concentration had no significant influence on DILI risk.

4. Discussion

Our study shows that the CG genotype of CYP2E1 (rs 3813867) is the most frequent in Indonesian tuberculosis patients. Moreover, according to the acetylator status, SAs had a higher incidence of hepatotoxicity when compared with RAs and SAs with the heterozygous c1/c2 variant (CG). Previous studies explained a significant association between the c1/c1 genotype and the risk of developing DILI by a higher CYP2E1 activity in patients with the c1/c1 genotype and the inhibitory effect of INH.^{18,19} Another study showed that the CYP2E1*1A/*1A polymorphism may confer a greater risk of INH-induced hepatotoxicity in a heterogeneous ethnic population; however, the patients in this study were receiving monotherapy and potential interactions were not addressed.^{10,19} A recent study from Chamoro et al. reported that the CYP2E1 c1/c2 polymorphism did not show a significance.²⁰ Our study's findings are in line with previous studies which reported that acetylator status was a risk factor for DILI in tuberculosis patients.^{9,14,15,21}

There have been several reports on the relationship between INH concentration and ATDH. A previous study showed an association between INH serum concentration and NAT2 genotype in Chinese tuberculosis patients.²² INH is hydrolysed by NAT-2 and oxidized by CYP2E1 to form a hepatotoxic substance.²³ However, our study confirmed that there is no association between these parameters. Our study reports that around 40% TB patients had higher INH serum concentrations, and 45% among them are experienced DILI. Our study is in line with previous data that showed SA status is significantly associated with DILI in the Indonesian population.²⁴ Our patients underwent combination therapy for TB treatment, including INH, and drug-drug pharmacokinetic and pharmacodynamic interactions occurred. INH is known to have a biphasic effect on CYP2E1 activity that consists of inhibition followed by induction. Many studies have reported on drug combinations that increase the incidence of antituberculosis drug-induced hepatotoxicity up to 35%.²⁵⁻²⁸

Huang et al. showed that CYP2E1 by RsaI restriction polymorphism was associated with susceptibility to INH-induced hepatotoxicity.²⁰ INH is principally metabolized by N-acetyltransferase 2 (NAT-2) and cytochrome P450 2E1 (CYP2E1). NAT-2 plays a key role in the detoxification and elimination of drugs, and it is also involved in carcinogen

metabolism.²⁹ Modifications in NAT-2 activity can result in the accumulation of precursors, such as HZ and AcHZ, leading to the development of hepatotoxicity.³⁰⁻³² CYP2E1 is predominantly expressed in the liver where it activates (via oxidation) low molecular-weight lipophilic compounds, drugs, and procarcinogens,³³ and it is involved in INH metabolism and generates hepatotoxic intermediates.^{34,35} The wild-type and mutated CYP2E1 alleles have been named c1 and c2, respectively. There are several in vivo studies that have associated the c1/c1 genotype with increased transcriptional activity, protein levels, and enzymatic activity.^{20,21} Therefore, patients with the CYP2E1 (c1/c1) genotype generate more hepatotoxins. Other studies have found that the c1/c1 genotype may be a risk factor for INH-induced hepatotoxicity.^{34,35} These data are in contrast with our present study that found c1/c2 may be a risk factor for INH-induced hepatotoxicity in SAs.

This study has some limitations as follows: we did not consider the complexity mechanism of antituberculosis-induced hepatotoxicity which was also interfered by roles of other genes such as NAT2 and GST.³⁶ The other mechanism of antituberculosis-induced hepatotoxicity which we did not involve in this study was due to the rifampicin-INH interaction. Rifampicin may induce INH metabolism through CYP2E1 which also induces the hepatotoxicity.^{37,38} The small sample size in our study is caused by the one criteria of the study is newly diagnosed TB patients, which only small number during the study period.

5. Conclusions

SAs were associated with an increased DILI incidence. A large-scale population study is needed to confirm the association between INH serum concentration, CYP2E1 polymorphisms, acetylator status and DILI.

Conflict of interest

The authors have none to declare.

Acknowledgement

The authors thank the Indonesian National Health Institute for approval and funding to complete this study.

REFERENCES

- World Health Organization. *Global tuberculosis control: surveillance, planning, financing: WHO report 2015*. WHO/HTM/TB/2006.7. Geneva, Switzerland: World Health Organization; 2015.
- Davtyan K, Zachariah R, Davtyan H, et al. Performance of decentralised facilities in tuberculosis case notification and treatment success in Armenia. *Public Health Action*. (October):2014;(October):4.
- Higuchi N, Tahara N, Yanagihara K, et al. NAT2 6A, a haplotype of the N-acetyltransferase 2 gene, is an important biomarker for risk of anti-tuberculosis drug-induced hepatotoxicity in Japanese patients with tuberculosis. *World J Gastroenterol*. 2007;13:6003–6008.
- Tostmann A, Boeree MJ, Aarnoutse RE, de Lange WC, van der Ven AJ, Dekhuijzen R. Antituberculosis drug-induced hepatotoxicity: concise up-to-date review. *J Gastroenterol Hepatol*. 2008;23:192–202.
- Durant F, Jebrak G, Pessayre D, Fournier M, Bernau J. Hepatotoxicity of antitubercular treatments. Rationale for monitoring liver status. *Drug Saf*. 1996;15:394–405.
- Schaberg T, Rebhan K, Lode H. Risk factors for adverse drug event of isoniazid, rifampin and pyrazinamide in patients hospitalized for pulmonary tuberculosis. *Eur Respir J*. 1996;9:2026–2030.
- Lee WM. Drug-induced hepatotoxicity. *NEJM*. 1995;333:1118–1127.
- Yamamoto T, Suou T, Hirayama C. Elevated serum aminotransferase induced by isoniazid in relation to isoniazid acetyltransferase phenotype. *Hepatology*. 1996;6:295–298.
- Mitchell JR, Zimmerman HJ, Ishak KG, et al. Isoniazid liver injury: clinical spectrum, pathology, and probable pathogenesis. *Ann Intern Med*. 1976;84:181–192.
- Farrell GC. Drug-induced acute hepatitis. In: Farrell GC, ed. *Drug-induced Liver Disease*. Edinburgh: Churchill Livingstone; 1994:247–299.
- Ryan DE, Ramanathan L, Iida S, et al. Characterization of a major form of rat hepatic microsomal cytochrome P-450 induced by isoniazid. *J Biol Chem*. 1985;260:6385–6393.
- Spielberg SP. N-acetyltransferases: pharmacogenetics and clinical consequences of polymorphism drug metabolism. *J Pharmacokinet Biopharm*. 1996;24:509–519.
- Cascorbi I, Drakoulis N, Brockmüller J, Maurer A, Sperling K, Roots L. Arylamine N-acetyl transferase (NAT2) mutations and their allelic linkage in unrelated Caucasian individual: correlation with phenotypic activity. *Am J Hum Genet*. 1995;57:581–592.
- Pande JN, Singh SP, Khilnani GC, Khilnani S, Tandon RK. Risk factors for hepatotoxicity from antituberculosis drugs: a case-control study. *Thorax*. 1996;51:132–136.
- Dickinson DS, Bailey WC, Hirschowitz BI, Soong SC, Eidus L, Hodgkin MM. Risk factors for isoniazid (NIH)-induced liver dysfunction. *J Clin Gastroenterol*. 1981;3:271–279.
- Fukino K, Sasaki Y, Hirai S, et al. Effects of N-acetyltransferase 2 (NAT2), CYP2E1 and Glutathione-S-transferase (GST) genotypes on the serum concentrations of isoniazid and metabolites in tuberculosis patients. *J Toxicol Sci*. 2008;33:187–195.
- Department of Health and Human Services. *Common Terminology Criteria for Adverse Events*. Version 4.0. Bethesda, MD: National Institutes of Health; 2010.
- Huang YS, Chern HD, Su WJ, et al. Cytochrome P450 2E1 genotype and the susceptibility to antituberculosis drug induced hepatitis. *Hepatology*. 2003;37:924–930.
- Vuilleumier N, Rossier MF, Chiappe A, et al. CYP2E1 genotype and isoniazid-induced hepatotoxicity in patients treated for latent tuberculosis. *Eur J Clin Pharmacol*. 2006;62:423–429.
- Chamorro JG, Castagnino JP, Musella RM, et al. Sex, ethnicity, and slow acetylator profile are the major causes of hepatotoxicity induced by antituberculosis drugs. *J Gastroenterol Hepatol*. 2013;28:323–328.
- Mitchell JR, Thorgeirsson UP, Black M, et al. Increased incidence of isoniazid hepatitis in rapid acetylators: possible relation to hydranize metabolites. *Clin Pharmacol Ther*. 1975;18:70–79.
- Chen B, Li JH, Xu YM, Wang J, Cao XM. The influence of NAT2 genotypes on the plasma concentration of isoniazid and acetylisoniazid in Chinese pulmonary tuberculosis patients. *Clin Chim Acta*. 2006;365:104–108.
- Wang T, Yu HT, Wang W, Pan YY, HE LX, Wang ZY. Genetic polymorphisms of cytochrome P450 and glutathione S-transferase associated with antituberculosis drug induced hepatotoxicity in Chinese tuberculosis patients. *J Int Med Res*. 2010;38:977–986.
- Cho JH, Koh WJ, Ryu RJ, et al. Genetic polymorphisms of NAT2 and CYP2E1 associated with antituberculosis drug-induced hepatotoxicity in Korean patients with pulmonary tuberculosis. *Tuberculosis*. 2007;87:551–556.
- Hwang SJ, Wu JC, Lee CN, et al. A prospective clinical study of isoniazid-rifampicin-pyrazinamide-induced liver injury in an area endemic for hepatitis B. *J Gastroenterol Hepatol*. 1997;12:87–91.
- Wong WM, Wu PC, Yuen MF. Antituberculosis drug-related liver dysfunction in chronic hepatitis B infection. *Hepatology*. 2000;31:201–206.
- Branch RA, Adedoyin A, Frye RF, Wilson JW, Romkes M. In vivo modulation of CYP enzymes by quinidine and rifampicin. *Clin Pharmacol Ther*. 2000;68:401–411.
- Zand R, Nelson SD, Slattey JT, et al. Inhibition and induction of cytochrome P4502E1-catalysed oxidation by isoniazid in humans. *Clin Pharmacol Ther*. 1993;1003:142–149.
- Sabbagh A, Langaney A, Darlu P, Gerard N, Krishnamoorthy R, Poloni ES. Worldwide distribution of NAT2 diversity: implications for NAT2 evolutionary history. *BMC Genet*. 2008;9:21–35.
- Timbrell JA. Studies on the role of acetylhydrazine in isoniazid hepatotoxicity. *Arch Toxicol*. 1979;2:1–8.
- Lauterburg BH, Smith CV, Todd EL, Mitchell JR. Oxidation of hydrazine metabolites formed from isoniazid. *Clin Pharmacol Ther*. 1985;38:566–571.
- Ohno M, Yamaguchi I, Yamamoto I, et al. Slow N-acetyltransferase 2 genotype affects the incidence of isoniazid and rifampicin-induced hepatotoxicity. *Int J Tuberc Lung Dis*. 2000;4:256–261.
- Carriere V, Berthou F, Baird S, Belloc C, Beaune P, de Waziers I. Human cytochrome P450 2E1 (CYP2E1): from genotype to phenotype. *Pharmacogenetics*. 1996;6:203–211.
- Lee SW, Chung LS, Huang HH, Chuang TY, Liou YH, Wu LS. NAT2 and CYP2E1 polymorphisms and susceptibility to first-line anti-tuberculosis drug-induced hepatitis. *Int J Tuberc Lung Dis*. 2010;14:622–626.
- Ryan DE, Iida S, Wood AW, Thomas PE, Lieber CS, Levin W. Characterization of three highly purified cytochromes P-450 from hepatic microsomes of adult male rats. *J Biol Chem*. 1984;259:1239–1250.
- Perwitasari DA, Aththobari J, Wilffert B. Pharmacogenetics of isoniazid-induced hepatotoxicity. *Drug Metab Rev*. 2015;47:222–228.
- Jeong I, Park JS, Cho YJ, et al. Drug-induced hepatotoxicity of anti-tuberculosis drugs and their serum levels. *J Korean Med Sci*. 2015;30:167–172.
- Huang YS. Recent progress in genetic variation and risk of antituberculosis drug-induced liver injury. *J Chin Med Assoc*. 2014;77:169–173.

CYP2E1 polymorphism, acetylator profiles and drug-induced liver injury incidence of Indonesian tuberculosis patients

ORIGINALITY REPORT

17%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.mdpi.com Internet	54 words — 2%
2	kundoc.com Internet	40 words — 1%
3	search.crossref.org Internet	36 words — 1%
4	repository.unair.ac.id Internet	35 words — 1%
5	Yang Xiang, Long Ma, Weidong Wu, Wei Liu et al. "The Incidence of Liver Injury in Uyghur Patients Treated for TB in Xinjiang Uyghur Autonomous Region, China, and Its Association with Hepatic Enzyme Polymorphisms NAT2, CYP2E1, GSTM1 and GSTT1", PLoS ONE, 2014 Crossref	34 words — 1%
6	Wang, T., H. Yu, W. Wang, Y. Pan, L. He, and Z. Wang. "Genetic Polymorphisms of Cytochrome P450 and Glutathione S-transferase Associated with Antituberculosis Drug-induced Hepatotoxicity in Chinese Tuberculosis Patients", Journal of International Medical Research, 2010. Crossref	23 words — 1%
7	repository.wima.ac.id Internet	22 words — 1%
8	eprints.uad.ac.id Internet	19 words — 1%

- 9 Nouha Souayed, Malek Chennoufi, Fida Boughattas, Mohsen Hassine et al. "Circadian-time dependent tolerance and haematological toxicity to isoniazid in murine", *Biomedicine & Pharmacotherapy*, 2015
Crossref 18 words — 1%
-
- 10 Pengcheng Wang, Komal Pradhan, Xiao-bo Zhong, Xiaochao Ma. "Isoniazid metabolism and hepatotoxicity", *Acta Pharmaceutica Sinica B*, 2016
Crossref 17 words — < 1%
-
- 11 R. M. Wu, C. W. Cheng, K. H. Chen, D. E. Shan, J. W. Kuo, Y. F. Ho, H. D. Chern. *Journal of Neural Transmission*, 2002
Crossref 17 words — < 1%
-
- 12 journals.plos.org
Internet 15 words — < 1%
-
- 13 onlinelibrary.wiley.com
Internet 14 words — < 1%
-
- 14 ijocta.balikesir.edu.tr
Internet 14 words — < 1%
-
- 15 bmcnutr.biomedcentral.com
Internet 14 words — < 1%
-
- 16 bm Cresnotes.biomedcentral.com
Internet 13 words — < 1%
-
- 17 Rashmi R Shah. "Can pharmacogenetics help rescue drugs withdrawn from the market?", *Pharmacogenomics*, 09/2006
Crossref 12 words — < 1%
-
- 18 Abdul Rahman W, Nurkhasanah Nurkhasanah, Nanik Sulistyani. "The Effect of Ethanolic Extract of Roselle (*Hibiscuss sabdariffa* L) on Increasing SOD Activity", *International Journal of Public Health Science (IJPHS)*, 2016
Crossref 12 words — < 1%

19	www.wjgnet.com Internet	11 words — < 1%
20	www.nature.com Internet	11 words — < 1%
21	www.aids2008.org Internet	11 words — < 1%
22	S. V. Rana, R. P. Ola, Sanjeev K. Sharma, S. K. Arora, S. K. Sinha, P. Pandhi, K. Singh. "Comparison between acetylator phenotype and genotype polymorphism of n-acetyltransferase-2 in tuberculosis patients", <i>Hepatology International</i> , 2011 Crossref	10 words — < 1%
23	Stephens, Camilla, Maria Lucena, and Raúl Andrade. "Pharmacogenomics and Clinical Assessment of Drug-Induced Hepatotoxicity toward Improving Clinical Outcomes", <i>Handbook of Personalized Medicine</i> , 2014. Crossref	10 words — < 1%
24	Kerry Atkinson, David Mabey. "The Burden of Communicable Diseases in Low- and Middle-Income Countries", Wiley, 2019 Crossref	10 words — < 1%
25	web.stanford.edu Internet	9 words — < 1%
26	www.pubmedcentral.nih.gov Internet	9 words — < 1%
27	Sumita Verma, Neil Kaplowitz. "Hepatotoxicity of Antitubercular Drugs", Elsevier BV, 2013 Crossref	9 words — < 1%
28	baadalsg.inflibnet.ac.in Internet	9 words — < 1%

-
- 29 preview-systematicreviewsjournal.biomedcentral.com 9 words — < 1%
Internet
-
- 30 www.medcraveonline.com 9 words — < 1%
Internet
-
- 31 Haiyue Jiang, Xiaoyue Zhang, Jiayang Shen, Yu Zhang, Yiyang Gu, Tian Tian, Minjie Chu, Xun Zhuang, Yulong Lian. "Association between CYP2E1 and GOT2 gene polymorphisms and susceptibility and low-dose N,N-dimethylformamide occupational exposure-induced liver injury", International Archives of Occupational and Environmental Health, 2019
Crossref
-
- 32 Liu, S.. "Elucidation of CYP2E1 5' regulatory RsaI/PstI allelic variants and their role in risk for oral cancer", Oral Oncology, 200107
Crossref
-
- 33 www.sigaa.ufs.br 8 words — < 1%
Internet
-
- 34 R. Deng, T. Yang, Y. Wang, N. Tang. "<I>CYP2E1 RsaI/PstI</I> polymorphism and risk of anti-tuberculosis drug-induced liver injury: a meta-analysis [Review article]", The International Journal of Tuberculosis and Lung Disease, 2012
Crossref
-
- 35 isbweb.org 8 words — < 1%
Internet
-
- 36 Liu, Fang, An-xia Jiao, Xi-rong Wu, Wei Zhao, Qing-qin Yin, Hui Qi, Wei-wei Jiao, Jing Xiao, Lin Sun, Chen Shen, Jian-ling Tian, Dan Shen, Evelyne Jacqz-Aigrain, and A-dong Shen. "Impact of Glutathione S-Transferase M1 and T1 on Anti-Tuberculosis Drug-Induced Hepatotoxicity in Chinese Pediatric Patients", PLoS ONE, 2014.
Crossref

37

Rob Aarnoutse. "Chapter 18: Pharmacogenetics of Antituberculosis Drugs", S. Karger AG, 2011

Crossref

7 words — < 1%

EXCLUDE QUOTES ON

EXCLUDE
BIBLIOGRAPHY ON

EXCLUDE MATCHES OFF