

# HASIL\_CEK\_jurnl internasional

*by 2020 Jurnl Internasional*

---

**Submission date:** 15-Apr-2021 01:46PM (UTC+0700)

**Submission ID:** 1559769898

**File name:** 2020-Jurnal\_Internasional\_AMSJ-2020-N9-76\_2.pdf (1.37M)

**Word count:** 4028

**Character count:** 19257

**BACKWARD BIFURCATION IN A WITHIN-HOST TUBERCULOSIS MODEL**YUDI ARI ADI<sup>1</sup> AND ARIS THOBRIN

ABSTRACT. A within-host tuberculosis model describing the interaction between macrophages and Mycobacterium tuberculosis was developed and analyzed in this paper. We derive  $R_0$  as the threshold value of the model and analyze the existence and stability of equilibrium points. Furthermore, bifurcation analysis performed based on the use of the application of the center manifold theory. Explicit condition for the existence of backward bifurcation is given. Finally, numerical simulations are presented to support the theoretical findings.

**I. INTRODUCTION**

Mycobacterium tuberculosis (Mtb) is an agent that causes Tuberculosis (TBC), an infectious disease that has a high mortality rate in various countries. Until now, tuberculosis is still one of the main causes of human death in the world. Based on the 2019 Global Tuberculosis Report released by WHO, eight countries that accounted for two-thirds of the total global TB cases were India (27 %), China (9 %), Indonesia (8 %), Philippines (6 %), Pakistan (6 %), Nigeria (4 %), Bangladesh (4 %) and South Africa (3 %) [1].

The Mtb bacteria commonly infect the lungs with alveolar macrophages as its primary target [2]. Responding to the presence of Mtb bacteria, the immune system forms granulomas consisting of immune cells known as macrophages which are responsible for controlling and separating the pathogens that infect

<sup>1</sup>corresponding author

2010 Mathematics Subject Classification. 34C23, 92B05.

Key words and phrases. Tuberculosis model, stability, backward bifurcation.

the lungs. Macrophages that function as the main host cell niche for the growth and survival of Mtb, are found in several organs of the human body [3, 4]. Macrophages ingest Mtb bacteria and isolate them in cellular compartments where bacteria cannot grow back. After the droplet core containing Mtb bacteria reaches the pulmonary alveoli, these bacteria are digested by alveolar macrophages and most can be destroyed or inhibited. A small amount can multiply intracellularly and is released when macrophages die. The dies macrophages do not necessarily cause bacteria to die but instead becomes a fuel for bacteria to grow. Living bacteria can spread through the lymphatic channels or through the bloodstream to distant tissues and organs, including areas of the body where TB disease is likely to develop, such as lymph nodes, lung spleen, kidneys, brain, and bones [5–7]. MTb does not have classical virulence factors and can remain on the host during long-term latency and will not cause transmission or significant damage if the host's immunity is not impaired [8].

In recent years mathematical models have been developed to describe the dynamics of tuberculosis in the human body, see [9–13]. In [9], Ibarguen-Mondragon et al. developed and analyzed a mathematical model that described dynamic interactions between macrophage cells, Mtb bacteria, and T-cell. Furthermore, Shi et al. in [10] added the chemotherapy drug variable and analyzed it with control theory. Recent research in [12] based on the model in [11] shows how T cell responses, which are immune cells to the presence of Mtb bacterial intervention. Bifurcation analysis is carried out to show conditions so that TB can disappear from the human body. Furthermore, Zhang et al. [13] added an analysis of global stability and conducted a bifurcation analysis for the disease-free equilibrium point. However, these models have not been able to clearly determine the important factors that play a role in TB infection. Motivated by this, in this article we simplified Zhang's model in [13] by combining T cells into a single immune response, which reacts against MTb bacteria that infect macrophages. Furthermore, in this study, we will investigate a qualitative analysis and show that the model undergoes a backward bifurcation. The bifurcation analysis in this study will be carried out based on the use of the center manifold theory as introduced by Castillo-Chavez and Song in [15].

This paper is organized as follows. We begin by formulating our model and establishing the properties of solutions in Section 2. In Section 3, we discuss the existence of equilibria and stability of the disease-free equilibrium point. In

section 4, we investigate the condition for the existence of backward bifurcation. To support the theoretical results, some numerical simulations are given in Section 5. Finally, discussion and conclusion are given in Section 6.

2. MODEL FORMULATION

Based on a within-host tuberculosis model in [13], we study a 3-dimensional ODE model that describes the interaction between uninfected macrophages ( $M_u$ ), infected macrophages ( $M_i$ ), and MTb bacteria ( $B$ ). Our model is given as:

$$(2.1) \quad \begin{aligned} \frac{dM_u}{dt} &= \Lambda - \mu M_u - \frac{\beta M_u B}{1 + \alpha B}, \\ \frac{dM_i}{dt} &= \frac{\beta M_u B}{1 + \alpha B} - c M_i - \frac{k M_i}{1 + \varepsilon M_i}, \\ \frac{dB}{dt} &= r c M_i - \gamma M_u B - d B. \end{aligned}$$

In model (2.1), we assume that uninfected macrophages reproduce at a constant rate of  $\Lambda$  and decrease by a natural death at a rate of  $\mu$ . Uninfected macrophages became infected at a saturated incidence rate of  $\frac{\beta M_u B}{1 + \alpha B}$ , with  $\beta$  is the maximal transmission of infection rate, while  $\frac{1}{1 + \alpha B}$  is an inhibition effect. Inside the infected macrophages, MTb bacteria multiply up to a limit in which the infected macrophages bursts, followed by the release of new bacteria. In this model,  $c$  is the rate of macrophages burst and  $r$  is the average number of the MTb bacteria released by infected macrophages. The releasing bacteria then infect macrophages or ingested and killed by uninfected macrophages. The parameters  $\gamma$  and  $d$  are the MTb bacteria death rate by uninfected macrophages and MTb bacteria natural death rate. The infected macrophages die caused by the adaptive immune response which modeled in a density-dependent term,  $\frac{k M_i}{1 + \varepsilon M_i}$ . In this model,  $k$  and  $\varepsilon$  are maximum killing rate and half-saturation constant, respectively.

The following theorem verifies that the system (2.1) has biological meaningful, that is, all solutions with nonnegative initial conditions remain nonnegative for all  $t \geq 0$ .

**Theorem 2.1.** Let  $\bar{\mu} = \min\{\mu, c\}$ . The set

$$\Omega = \{(M_u, M_i, B) \in \mathbf{R}_+^3 : M_u + M_i \leq \frac{\Lambda}{\bar{\mu}}, B \leq \frac{d\bar{\mu}}{rc\Lambda}\}$$

is a positive invariant and attracting set for equation (2.1).

*Proof.* From the first equation of (2.1), we have

$$\frac{dM_u}{dt} \leq \Lambda - \mu M_u.$$

Thus, from standard comparison theorem we have  $\lim_{t \rightarrow \infty} \sup M_u(t) \leq \frac{\Lambda}{\mu}$ .

Let  $M(t) = M_u(t) + M_i(t)$ . Then we get

$$\begin{aligned} \frac{dM}{dt} &= \Lambda - \mu M_u - cM_i - \frac{kM_i}{1 + \varepsilon M_i} \\ &\leq \Lambda - \mu M_u - cM_i \leq \Lambda - \bar{\mu}, \end{aligned}$$

with  $\bar{\mu} = \min\{\mu, c\}$ . Thus, we have

$$0 \leq M(t) \leq \frac{\Lambda}{\bar{\mu}}.$$

Clearly,  $M_i \leq \frac{\Lambda}{\bar{\mu}}$ .

From the third equation of (2.1), we have

$$\frac{dB}{dt} = rcM_i - \gamma M_u B - dB \leq rc\frac{\Lambda}{\bar{\mu}} - dB.$$

Hence,  $\lim_{t \rightarrow \infty} \sup B(t) \leq \frac{rc\Lambda}{d\bar{\mu}}$ . Therefore, all solution of (2.1) that starting in  $\Omega$  remain in  $\Omega$  for all  $t \geq 0$ .  $\square$

### 3. EXISTENCE AND STABILITY OF EQUILIBRIA

In this sections, we will consider the existence and stability of the equilibrium points of system (2.1). The crucial quantity is the basic reproduction number that measures the expected number of secondary infections that result from one newly infected cell. Applying the procedure of next-generation matrix described in [14], we obtain

$$(3.1) \quad R_0 = \sqrt{\frac{rc\beta\Lambda}{(c+k)(\gamma\Lambda + d\mu)}}.$$

Then we get the following theorem.

41

**Theorem 3.1.** Consider the system (2.1).

- (i) If  $R_0 < 1$  and  $c + k - \varepsilon\Lambda \geq 0$ , then the system (2.1) only has a disease free equilibrium  $E_0 = \left(\frac{\Lambda}{\mu}, 0, 0\right)$ .
- (ii) If  $R_0 < 1$  and  $c + k - \varepsilon\Lambda < 0$ , in addition to  $E_0$ , then the system (2.1) may has two or no positive disease equilibrium points.
- (iii) If  $R_0 > 1$ , in addition to  $E_0$ , the system (2.1) at least has one positive disease equilibrium point.

*Proof.* Solving the system (2.1) by setting the right-hand side equal to zero, we get a disease-free equilibrium point, denoted by  $E_0$ , and it is given by

$$E_0 = \left(\frac{\Lambda}{\mu}, 0, 0\right)$$

and disease equilibrium points, denoted by  $E_{1j^*}$ , and given by

$$E_{1j^*} = (M_u^*, M_i^*, B^*),$$

with

$$(3.2) \quad M_u^* = \frac{\Lambda(1 + \alpha B^*)}{\mu + (\alpha\mu - \beta)B^*}, M_i^* = \frac{(\gamma M_u^* + d)B^*}{rc},$$

and  $B^*$  is any positive root of the cubic equations

$$(3.3) \quad a_3 B^3 + a_2 B^2 + a_1 B + a_0 = 0,$$

where

$$\begin{aligned} a_3 &= (\varepsilon\alpha^2(d^2\mu^2 + 2d\gamma\mu\Lambda + \gamma^2\Lambda^2) + \varepsilon\alpha(2d^2\mu\beta + 2d\gamma\Lambda\beta) + \varepsilon d^2\beta^2), \\ a_2 &= \alpha^2\mu r(cd\mu + \gamma c\Lambda + \mu kd + k\gamma\Lambda) + r\alpha\beta(c + k - \Lambda\varepsilon)(d\mu + \gamma\varepsilon) \\ &\quad + 2d\Lambda\gamma\beta\varepsilon + (c + k)d\mu\alpha\beta + \alpha\varepsilon(2d^2\mu^2 + 4d\Lambda\gamma\mu + 2\gamma^2\Lambda^2) \\ &\quad + rd\beta^2(c + k - \varepsilon\Lambda) + 2d^2\mu\beta\varepsilon, \\ a_1 &= r(\beta + \mu\alpha)(c + k)(d\mu + \Lambda\gamma)(1 - R_0^2) + r\beta(c + k - \varepsilon\Lambda)(d\mu + \Lambda\gamma) \\ &\quad + r\mu\alpha(c + k)(d\mu + \Lambda\gamma) + \Lambda\gamma\varepsilon(\gamma\Lambda + 2d\mu) \\ a_0 &= r\mu(c + k)(d\mu + \Lambda\gamma)(1 - R_0^2). \end{aligned}$$

In order to get a positive disease equilibrium, the condition  $M_u^* > 0$  must be hold. From (3.2), it is clear that  $M_u^* > 0$  if only if  $\mu + (\alpha\mu - \beta)B^* > 0$  or  $B^* < \frac{\mu}{\beta - \alpha\mu}$ , with  $B^*$  is positive roots of equation (3.3). The coefficient  $a_3$  in equation (3.3)

is always positive, while  $a_2$ ,  $a_1$ , and  $a_0$  can be positive or negative. Furthermore, we have

- (i) If  $R_0 < 1$  and  $c + k - \varepsilon\Lambda \geq 0$ , then both of the coefficients  $a_2$  and  $a_1$  are positive, so there are no changes of coefficient sign of equation (3.3). According the Descartes rule of sign, there are no positive real root of equation (3.3). Hence, the system (2.1) only has a disease-free equilibrium  $E_0 = \left(\frac{\Lambda}{\mu}, 0, 0\right)$ .
- (ii) If  $R_0 < 1$  and  $c + k - \varepsilon\Lambda < 0$ , then the coefficient  $a_0$  is always positive, while both  $a_1$  and  $a_2$  can be positive or negative. According the Descartes rule of sign, equation (3.3) may have two or no positive real roots. Therefore, in addition to  $E_0$ , then the system (2.1) may has two or no positive disease equilibrium points.
- (iii) If  $R_0 > 1$ , the coefficient  $a_0$  is always negative. Since  $a_3$  is always positive, at least there is one change of coefficient sign of equation (3.3). According the Descartes rule of sign, there is at least one positive root of equation (3.3). The possible changes of sign for the coefficient of equation (3.3) can be determined from Table 1. Therefore, in addition to  $E_0$ , the system (2.1) at least has one positive disease equilibrium point.

This completes the proof. □

TABLE 1. Sign of the coefficients of equation (3.3) for  $R_0 > 1$ .

$a_3$	$a_2$	$a_1$	$a_0$
+	+	+	-
+	+	-	-
+	-	+	-
+	-	-	-

The following theorem shows that  $R_0$  is a threshold quantity for the stability of disease-free equilibrium point of system (2.1). The stability determined by the sign of real part of eigenvalues of the Jacobian matrix at the given equilibrium point.

**Theorem 3.2.** *The disease-free equilibrium  $E_0 = \left(\frac{\Lambda}{\mu}, 0, 0\right)$  is locally asymptotically stable if  $R_0 < 1$  and unstable if  $R_0 > 1$ .*

<sup>11</sup> *Proof.* The Jacobian matrix of system (2.1) at  $E_0 = \left(\frac{\Lambda}{\mu}, 0, 0\right)$  is

$$J(E_0) = \begin{bmatrix} -\mu & 0 & -\frac{\beta\Lambda}{\mu} \\ 0 & -(c+k) & \frac{\beta\Lambda}{\mu} \\ 0 & rc & -\frac{\gamma\Lambda+d\mu}{\mu} \end{bmatrix}.$$

that gives the following characteristic equation

$$(3.4) \quad (\lambda + \mu)(\lambda^2 + P_1\lambda + P_0) = 0.$$

where

$$P_1 = k + c + \frac{\gamma\Lambda + d\mu}{\mu} \geq 0,$$

$$P_0 = \frac{(k + c)(\gamma\Lambda + d\mu) - rc\beta\Lambda}{\mu}.$$

<sup>50</sup> It is clear that all eigenvalues have negative real part if  $P_0 > 0$ , that is  $(k+c)(\gamma\Lambda + d\mu) - rc\beta\Lambda > 0$  or  $\frac{rc\beta\Lambda}{(k+c)(\gamma\Lambda+d\mu)} < 1$ . It implies that the disease-free equilibrium  $E_0$  is locally asymptotically stable if  $R_0 < 1$  and unstable if  $R_0 > 1$ .  $\square$

Now, consider the critical case  $R_0 = 1$ . It implies that  $rc\beta\Lambda = (k+c)(\gamma\Lambda + d\mu)$ . In this case the characteristic equation (3.4) has two negative eigenvalues and one zero eigenvalue. In the next section, we will use the application of center manifold theorem introduced in [15] to study the stability of the equilibrium point  $E_0$ .

#### 4. THE EXISTENCE OF BACKWARD BIFURCATION

Based on the formula of basic reproduction number,  $R_0$  in (3.1), we find that the stability of equilibrium point significantly affected by the transmission of infection rate  $\beta$ , the average number of the MTb bacteria  $r$ , and the rate of macrophages burst  $c$ . <sup>35</sup> In this section, we investigate the effect of parameters to the number of equilibrium points and their stability change by varying the parameters value. We thus choose the transmission of infection rate  $\beta$  as a bifurcation parameter and derive the sufficient conditions for the occurrence of backward and forward bifurcation. <sup>23</sup> We have the following theorem.

**Theorem 4.1.** *The system (2.1) in  $R_0 = 1$  exhibits backward bifurcation if  $k\Lambda\epsilon\mu^2 - 1 + \Lambda(c+k)(\gamma - \alpha\beta) > 0$ .*



*Proof.* From (3.1) we find that  $R_0 = 1$  iff  $\beta = \beta^* = \frac{(c+k)(\gamma\Lambda+d\mu)}{rc\Lambda}$  and  $E_0$  is locally asymptotically stable if  $\beta < \beta^*$ , whereas it loses its stability if  $\beta > \beta^*$ . Evaluating the eigenvalues of the Jacobian matrix of system (2.1) at  $(E_0, \beta^*)$  we get  $\lambda_1 = -\mu$ ,  $\lambda_2 = 0$ , and  $\lambda_3 = -\frac{\mu(c+k)(\gamma\Lambda+d\mu)}{\mu}$ . It is clear that  $\lambda_2 = 0$  is a single zero eigenvalue and the others eigenvalues are real and negative. Hence,  $E_0$  is a non hyperbolic equilibrium point and the system (2.1) can undergo backward bifurcation at  $\beta = \beta^*$ . Furthermore, we can determine the eigenvector associated with the zero eigenvalue.

Let  $\mathbf{w} = (w_1, w_2, w_3)^T$  be the right eigenvector of the Jacobian matrix  $J(E_0, \beta^*)$ . We have

$$\begin{aligned} -\mu w_1 - \frac{(c+k)(\gamma\Lambda+d\mu)}{\mu rc} w_3 &= 0, \\ -(c+k)w_2 + \frac{(c+k)(\gamma\Lambda+d\mu)}{\mu rc} w_3 &= 0, \\ rcw_2 - \frac{(\gamma\Lambda+d\mu)}{\mu} w_3 &= 0. \end{aligned}$$

Then we get

$$(4.1) \quad \mathbf{w} = \left( -\frac{(c+k)(\gamma\Lambda+d\mu)}{\mu^2 rc}, \frac{\gamma\Lambda+d\mu}{\mu rc}, 1 \right).$$

The left eigenvector of the Jacobian matrix  $J(E_0, \beta^*)$  satisfying  $\mathbf{v} \cdot \mathbf{w} = 1$  is given by

$$(4.2) \quad \mathbf{v} = \left( 0, -\frac{\mu^2 rc}{(c+k)(\gamma\Lambda+d\mu)}, \frac{\mu}{\gamma\Lambda+d\mu+\mu} \right).$$

Now, we define the coefficient  $a$  and  $b$ :

$$a = \sum_{i,j,k=1}^3 v_k w_i w_j \frac{\partial^2 f_k}{\partial x_i \partial x_j}(E_0, \beta^*), \quad b = \sum_{i,k=1}^3 v_k w_i \frac{\partial^2 f_k}{\partial x_i \partial \beta}(E_0, \beta^*).$$

In view of (4.1), (4.2), and

$$\begin{aligned} f_1 &= \Lambda - \mu M_u - \frac{\beta M_u B}{1 + \alpha B}, \\ f_2 &= \frac{\beta M_u B}{1 + \alpha B} - c M_i - \frac{k M_i}{1 + \varepsilon M_i}, \\ f_3 &= rc M_i - \gamma M_u B - dB, \end{aligned}$$

we then get

$$a = \frac{2(\gamma\Lambda + d\mu)\kappa}{\Lambda\mu rc(c+k)(\gamma\Lambda + d\mu + \mu)}$$

with  $\kappa = \Lambda\epsilon k\mu^2 - 1 + \Lambda(c+k)(\gamma - \alpha c)$  and

$$b = \frac{\Lambda rc}{(c+k)(\gamma\Lambda + d\mu + \mu)}.$$

We find that  $b$  is always positive. Next, by applying the Theorem 4 in [15], we conclude that the system (2.1) exhibits a backward bifurcation if  $\kappa > 0$ , i.e  $\Lambda\epsilon k\mu^2 - 1 + \Lambda(c+k)(\gamma - \alpha c) > 0$ . Otherwise, forward bifurcation is occur. This completes the proof.  $\square$

The existence of backward bifurcation plays an important role in a management of disease. The backward bifurcation shows that a stable disease-free equilibrium coexist with both a stable and unstable disease equilibrium. Biologically, the backward bifurcation at  $R_0 = 1$  makes treatment more difficult, because the condition  $R_0 < 1$  is not completely sufficient to eradicate infectious cases.

18  
5. NUMERICAL SIMULATIONS

In this section, we give some numerical simulations to demonstrate the theoretical results in the previous section. In the simulation, we use the parameter values obtained from literature [13]. We consider a set of parameter values:

$$(5.1) \quad \Lambda = 3500, \mu = 0.01, \beta = 0.8 \times 10^{-8}, \alpha = 0.01, c = 0.3, \\ k = 1, \epsilon = 30, r = 30, \gamma = 0.125 \times 10^{-8}, d = 0.05.$$

With these set of parameter values, the condition of Theorem 3.1(ii) is satisfied. We have  $R_0 = 0.61994 < 1$ ,  $c + k - \epsilon\Lambda = -1.0499 \times 10^5 < 0$  and  $E_0 = (350000, 0, 0)$ . From Theorem 3.2 we know that  $E_0$  is locally asymptotically stable, see Figure 1. In this case, the system (2.1) has no positive disease equilibrium point. Figure 1 confirm that all trajectories approach to the equilibrium point  $E_0$ .

Now, we will analyze the effect of variation in parameter  $\beta$  and keeping all other parameters fixed as given in (5.1). We found that, if  $0 < \beta < 2.0815476 \times 10^{-8}$ , then  $c+k-\epsilon\Lambda < 0$  and  $R_0 < 1$ . We noticed that, if  $0 < \beta < 0.9619088 \times 10^{-8}$ , then the equation (3.3) has no positive real root, so the system (2.1) has no positive disease equilibrium point. If  $0.9619088 \times 10^{-8} < \beta < 2.0815476 \times 10^{-8}$ ,




FIGURE 1. The system (2.1) has only one disease free equilibrium  $E_0 = (3.5 \times 10^5, 0, 0)$  and it is locally asymptotically stable.

then the equation (3.3) has two positive real roots, so the system (2.1) has two positive disease equilibrium points. As illustration in Figure 4, with  $\beta = 1.08 \times 10^{-8}$ , in addition to  $E_0 = (3.5 \times 10^5, 0, 0)$  we found two disease equilibrium points, i.e unstable equilibrium point  $E_{11}^* = (3.49995 \times 10^5, 0.082393, 14.702156)$  and stable equilibrium point  $E_{12}^* = (3.49998 \times 10^5, 0.47267, 84.34271)$ .

If we set  $\beta = 2.28 \times 10^{-8}$ , we have  $R_0 = 1.04658 > 1$  and the condition of Theorem 3.1(iii) is satisfied. From Theorem 3.2, we know that the disease-free equilibrium point  $E_0$  is unstable. The system (2.1) has only one stable disease equilibrium  $E_{11}^* = (3.49937 \times 10^5, 1.9586, 349.4906)$ . Figure 3 confirm that all trajectories approach the equilibrium point  $E_{11}^*$ .

Finally, we find that with the set of parameter values as in (5.1), the condition in Theorem 4.1 is satisfied, that is  $k\Lambda\varepsilon\mu^2 - 1 + \Lambda(c+k)(\gamma - \alpha\beta) = 9.5 > 0$ . If the equilibrium point  $E_0$  continued with respect to parameter  $\beta$ , then the system(2.1) exhibits a backward bifurcation at  $R_0 = 1$  associated with  $\beta = \beta^* = 2.0815476 \times 10^{-8}$ . Subsequently, there is a saddle-node bifurcation at  $\beta = \beta^* = 0,961908 \times 10^{-8}$  since the number of disease equilibrium point changes from one to three as  $\beta$  passes through this point. Figure 4 shows that bistable behaviour exist whenever  $0,961908 \times 10^{-8} < \beta < 2.0815476 \times 10^{-8}$ . In this interval, both




FIGURE 2. Phase portraits Infected macrofag vs Mtb bacteria with  $\beta = 1.08 \times 10^{-8}$ . The system (2.1) has one disease-free equilibrium  $E_0 = (3.5 \times 10^5, 0, 0)$  and it is locally asymptotically stable.




FIGURE 3. Phase portraits confirm that the system (2.1) has one stable disease equilibrium point  $E_{11}$  whenever  $\beta = 2.28 \times 10^{-8}$ .




FIGURE 4. Backward bifurcation at  $\beta = \beta^* = 2,0815476 \times 10^{-8}$ .

the disease-free equilibrium point  $E_0$  and the disease equilibrium point  $E_{12}^*$  are stable, while the disease equilibrium point  $E_{11}^*$  is unstable. Such conditions indicate that to eliminate the disease, it is not sufficient to reduce  $R_0$  below 1. From Figure 4, we know that,  $\beta$  must be below  $0,961908 \times 10^{-8}$ , associated with  $R_0$  below  $0.679788$  to guarantee that the system (2.1) has only disease-free equilibrium  $E_0$ .

## 25 6. DISCUSSION AND CONCLUSION

In this paper, we have studied a within-host tuberculosis model describing the interaction between Microbacterium tuberculosis and macrophages. The dynamics of the model depends on the stability of the equilibrium points. In order to determine the TBC treatment, we have to find the analytical condition so that all of the trajectories converge to the disease-free equilibrium. This condition is given in Theorem ref th-01. We found the threshold value  $R_0$ , so that the disease free equilibrium  $E_0$  is stable when  $R_0 < 1$  and unstable when  $R_0 > 1$ . Medically, if all of the trajectories converge to disease-free equilibrium, then Mtb bacteria and infected macrophage will eliminate so that the patient is ultimately cured of TBC. Meanwhile, if trajectories converge to the disease

equilibrium  $E_{1j}^*$ , then Mtb bacteria and infected macrophage will persist in the body.

Further, with the help of bifurcation analysis, we have shown that the continuation of the diseases free equilibrium point  $E_0$  with respect to the parameter  $\beta$ , causes the system (2.1) undergoes a backward bifurcation at  $R_0 = 1$ . Through studying the backward bifurcation in our model we know that the basic reproduction number below unity is not fully adequate to eradicate the disease. From this result, we suggest that in order to successfully eradicate the disease, it is necessary to optimize and improve the efficiency of treatment that can reduce Mtb bacterial transmission. In other words, medical methods and technology need to be improved. A more realistic in-host TB model can also be found by selecting the most relevant treatment method with respect to actual data, so it can help clinicians to suggest the right treatment for TB patients. We will learn more in future research.

#### ACKNOWLEDGMENT

The authors would like to thank the editor and the anonymous referees for their careful reading the original paper and their valuable comments to the improvement of this work. This work is supported by the University of Ahmad Dahlan.

#### REFERENCES

- [1] WHO : *Global tuberculosis report 2019*, WHO, 2019.
- [2] A. AZAD, A. PAPP, X. ZHOU, ET AL. : *Alveolar macrophages responses to Mycobacterium tuberculosis reveal human to human variation in important immunobiology pathways gleaned from functional genomics study*, *The J. Immunology*, **202**(1) (2019), ID62:4.
- [3] J. I. MOLIVA, J. TURNER, J. B. TORRELLES : *Immune Responses to Bacillus Calmette-Guerin Vaccination: Why Do They Fail to Protect against Mycobacterium tuberculosis?*, *Front. Immunol.*, **8**(407) (2017), doi: 10.3389/fimmu.2017.00407.
- [4] E. GUIRADO, L. S. SCHLESINGER, G. KAPLAN : *Macrophages in Tuberculosis: Friend or Foe*, *Semin Immunopathol* **35**(5) (2013), 563–583.
- [5] E. GUIRADO, L. S. SCHLESINGER: *Modeling the Mycobacterium tuberculosis granuloma—the critical battlefield in host immunity and disease*, *Front. Immunol.*, **4**(98) (2013), doi:10.3389/fimmu.2013.00098.

- [6] D. MAHAMED, M. BOILLE, Y. GANGA, ET AL.: *Intracellular growth of Mycobacterium tuberculosis after macrophage cell death leads to serial killing of host cells*, eLife (2017);6:e22028 doi: 10.7554/eLife.22028.
- [7] S. B. COHEN, B. H. GERN, J. L. DELAHAYE, K. N. ADAMS, C. R. PLUMLEE, J. WINKLER, D. R. SHERMAN, M. Y. GERNER, K. B. URDAH : *Alveolar macrophages provide an early Mycobacterium tuberculosis niche and initiate dissemination*, Cell Host Microbe, **24** (3) (2018), 439446.e4. doi:10.1016/j.chom.2018.08.001.
- [8] Q. CHAI, Y. ZHANG, C. H. LIU : *Mycobacterium tuberculosis: An Adaptable Pathogen Associated With Multiple Human Diseases*, Front. Cell. Infect. Microbiol., **8**(158) (2018), doi: 10.3389/fcimb.2018.00158.
- [9] E. IBARGUEN-MONDRAGON, L. ESTEVA, L. CHAVEZ-GALAN: *A mathematical model for cellular immunology of tuberculosis*, Math Biosci Eng., **8**(4) 2011), 973 – 986. doi: 10.3934/mbe.2011.8.973.
- [10] R. SHI, Y. LI, S. TANG : *A Mathematical Model with Optimal Controls for Cellular Immunology of Tuberculosis*, Taiwanese Journal of Mathematics, **18**(2) (2014), 575 – 597.
- [11] Y. DU, J. WU, J. M. HEFFERNAN : *A simple in-host model for Mycobacterium tuberculosis that captures all infection outcomes*, Math. Popul. Stud., **24**(1) (2017), 37 – 63.
- [12] W. ZHANG : *Analysis of an in-host tuberculosis model for disease control*, Applied Mathematics Letters, **99** (2020), <https://doi.org/10.1016/j.aml.2019.07.014>.
- [13] W. ZHANG, F. FRASCOLI, J. M. HEFFERNAN : *Analysis of solutions and disease progressions for a within-host tuberculosis model*, Mathematics in Applied Sciences and Engineering, **1**(1) (2020), 39 – 49.
- [14] P. VAN DEN DRIESSCHE, J. WATMOUGH : *Reproduction Numbers and sub-threshold endemic equilibria for compartmental model of disease transmission*, Mathematical Biosciences, **180** (2002), 29 – 48.
- [15] C. CASTILLO-CHAVEZ, B. SONG: *Dynamical Model of Tuberculosis and Their Applications*, Math. Biosci. Eng., **1**(2) (2004), 361 – 404.

DEPARTMENT OF MATHEMATICS  
FACULTY OF APPLIED SCIENCE AND TECHNOLOGY  
UNIVERSITY OF AHMAD DAHLAN  
YOGYAKARTA 55191, INDONESIA  
Email address: yudi.adi@math.uad.ac.id

DEPARTMENT OF MATHEMATICS  
FACULTY OF APPLIED SCIENCE AND TECHNOLOGY  
UNIVERSITY OF AHMAD DAHLAN  
YOGYAKARTA 55191, INDONESIA  
Email address: aris.thobi@math.uad.ac.id

# HASIL\_CEK\_jurnl internasional

---

## ORIGINALITY REPORT

---

19%

SIMILARITY INDEX

9%

INTERNET SOURCES

17%

PUBLICATIONS

6%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

- |   |   |    |
|---|---|----|
| 1 | Kassa, S.M.. "Epidemiological models with prevalence dependent endogenous self-protection measure", <i>Mathematical Biosciences</i> , 201101<br>Publication   | 1% |
| 2 | <a href="https://www.research-publication.com">research-publication.com</a><br>Internet Source  | 1% |
| 3 | Submitted to Grand Canyon University<br>Student Paper   | 1% |
| 4 | Santosh Mittal, Devindra Kumar Vohra, Parul Mittal. "A Study to Evaluate Magnitude of Rifampicin Resistance in Cases of Sputum Positive Pulmonary Tuberculosis in a Tertiary Care Centre in Western U.P.", <i>Journal of Evidence Based Medicine and Healthcare</i> , 2020<br>Publication | 1% |
| 5 | <a href="https://www.mafiadoc.com">mafiadoc.com</a><br>Internet Source  | 1% |
-



6

Zain Ul Abadin Zafar, Kashif Rehan, M. Mushtaq. "HIV/AIDS epidemic fractional-order model", Journal of Difference Equations and Applications, 2017

Publication

1 %

7

Nakul Chitnis, J. M. Cushing, J. M. Hyman. "Bifurcation Analysis of a Mathematical Model for Malaria Transmission", SIAM Journal on Applied Mathematics, 2006

Publication

1 %

8

Wang, Xueying, and Jin Wang. "Analysis of cholera epidemics with bacterial growth and spatial movement", Journal of Biological Dynamics, 2015.

Publication

1 %

9

Eduardo Ibarquen-Mondragon, Lourdes Esteva. "On CTL response against mycobacterium tuberculosis", Applied Mathematical Sciences, 2014

Publication

1 %

10

Submitted to Higher Education Commission Pakistan

Student Paper

1 %

11

Miaoran Yao, Yongxin Zhang, Wendi Wang. "Bifurcation analysis for an in-host Mycobacterium tuberculosis model", Discrete & Continuous Dynamical Systems - B, 2017

Publication

1 %

12 Wenjun Jing, Zhen Jin, Juping Zhang. "An SIR pairwise epidemic model with infection age and demography", Journal of Biological Dynamics, 2018  
Publication 1 %

---

13 Mehdi Lotfi, Azizeh Jabbari, Hossein Kheiri. "A mathematical analysis of a tuberculosis epidemic model with two treatments and exogenous re-infection", International Journal of Biomathematics, 2020  
Publication <1 %

---

14 Submitted to Indian Institute of Technology  
Student Paper <1 %

---

15 Submitted to University of Nottingham  
Student Paper <1 %

---

16 Evelyn Guirado, Larry S. Schlesinger, Gilla Kaplan. "Macrophages in tuberculosis: friend or foe", Seminars in Immunopathology, 2013  
Publication <1 %

---

17 S. MUSHAYABASA, C. P. BHUNU, G. MAGOMBEDZE, A. G. R. STEWART. "ON THE ROLE OF SCREENING AND EDUCATIONAL CAMPAIGNS ON CONTROLLING HCV IN CORRECTIONAL INSTITUTIONS", Journal of Biological Systems, 2013  
Publication <1 %

---

18

Xia Wang, Sanyi Tang, Xinyu Song, Libin Rong. "Mathematical analysis of an HIV latent infection model including both virus-to-cell infection and cell-to-cell transmission", *Journal of Biological Dynamics*, 2016

Publication

&lt;1 %

19

[polen.itu.edu.tr](http://polen.itu.edu.tr)

Internet Source

&lt;1 %

20

Edoardo Beretta, Yang Kuang. "Modeling and analysis of a marine bacteriophage infection with latency period", *Nonlinear Analysis: Real World Applications*, 2001

Publication

&lt;1 %

21

Larissa Sens, Ana Caroline Arruda de Souza, Lucas Antonio Pacheco, Angela Camila Orbem Menegatti et al. "Synthetic thiosemicarbazones as a new class of *Mycobacterium tuberculosis* protein tyrosine phosphatase A inhibitors", *Bioorganic & Medicinal Chemistry*, 2018

Publication

&lt;1 %

22

Qiyao Chai, Yong Zhang, Cui Hua Liu. "Mycobacterium tuberculosis: An Adaptable Pathogen Associated With Multiple Human Diseases", *Frontiers in Cellular and Infection Microbiology*, 2018

Publication

&lt;1 %

23 Pei Yu, Wenjing Zhang, Lindi M. Wahl. <1 %  
"Dynamical analysis and simulation of a 2-  
dimensional disease model with convex  
incidence", Communications in Nonlinear  
Science and Numerical Simulation, 2016  
Publication

---

24 [home.cc.umanitoba.ca](http://home.cc.umanitoba.ca) <1 %  
Internet Source

---

25 [sciencepubco.com](http://sciencepubco.com) <1 %  
Internet Source

---

26 [www.intechopen.com](http://www.intechopen.com) <1 %  
Internet Source

---

27 A. O. Egonmwan, D. Okuonghae. <1 %  
"Mathematical analysis of a tuberculosis  
model with imperfect vaccine", International  
Journal of Biomathematics, 2019  
Publication

---

28 DANIEL OKUONGHAE, VINCENT AIHIE. "CASE  
DETECTION AND DIRECT OBSERVATION  
THERAPY STRATEGY (DOTS) IN NIGERIA: ITS  
EFFECT ON TB DYNAMICS", Journal of  
Biological Systems, 2011  
Publication

---

29 Danilo Alonso Ortega Bejarano, Eduardo  
Ibarguen-Mondragon, Enith Amanda Gomez-  
Hernandez. "A stability test for non linear  
systems of ordinary differential equations

based on the Gershgorin circles",  
Contemporary Engineering Sciences, 2018

Publication

30

Sk Shahid Nadim, Joydev Chattopadhyay.  
"Occurrence of backward bifurcation and  
prediction of disease transmission with  
imperfect lockdown: A case study on COVID-  
19", Chaos, Solitons & Fractals, 2020

Publication

<1 %

31

Wenjing Zhang. "Analysis of an in-host  
tuberculosis model for disease control",  
Applied Mathematics Letters, 2020

Publication

<1 %

32

[prod--journal.elifesciences.org](http://prod--journal.elifesciences.org)

Internet Source

<1 %

33

CÉDRIC WOLF. "A Multi-Patch Epidemic Model  
with Periodic Demography, Direct and Indirect  
Transmission and Variable Maturation Rate",  
Mathematical Population Studies, 9/1/2006

Publication

<1 %

34

Miguel Fudolig, Reka Howard. "The local  
stability of a modified multi-strain SIR model  
for emerging viral strains", Cold Spring Harbor  
Laboratory, 2020

Publication

<1 %

35

Nasri, M.. "Study of a system of non-linear  
difference equations arising in a deterministic

<1 %

model for HIV infection", Applied Mathematics and Computation, 20051215

Publication

36

Sharomi, O.. "Curtailling smoking dynamics: A mathematical modeling approach", Applied Mathematics and Computation, 20080201

Publication

<1 %

37

[hal.univ-lorraine.fr](http://hal.univ-lorraine.fr)

Internet Source

<1 %

38

BING LI, SHENGQIANG LIU. "A DELAYED HIV-1 MODEL WITH MULTIPLE TARGET CELLS AND GENERAL NONLINEAR INCIDENCE RATE", Journal of Biological Systems, 2014

Publication

<1 %

39

Das, P.. "Qualitative study of a model of Chagas' disease", Mathematical and Computer Modelling, 200602

Publication

<1 %

40

Ma, Z.P.. "Stability and Hopf bifurcation analysis on a predator-prey model with discrete and distributed delays", Nonlinear Analysis: Real World Applications, 200904

Publication

<1 %

41

Peng Shi, Carlos E. De Souza, Lihua Xie. "Bounded real lemma for linear systems with finite discrete jumps", International Journal of Control, 1997

Publication

<1 %

42	S. MUSHAYABASA, C. P. BHUNU. "MODELING THE IMPACT OF VOLUNTARY TESTING AND TREATMENT ON TUBERCULOSIS TRANSMISSION DYNAMICS", International Journal of Biomathematics, 2012 Publication	<1 %
43	Stanca M. Ciupe, Jane M. Heffernan. "In-host modeling", Infectious Disease Modelling, 2017 Publication	<1 %
44	Submitted to University of KwaZulu-Natal Student Paper	<1 %
45	ZHANG, Xianghua, Fu CHEN, Ke WANG, and Hong DU. "Stochastic sirs model driven by lévy noise", Acta Mathematica Scientia, 2016. Publication	<1 %
46	article.sciencepublishinggroup.com Internet Source	<1 %
47	www.iosrjournals.org Internet Source	<1 %
48	Imane Agmour, meriem Bentounsi, Naceur Achtaich, Youssef El Foutayeni. "Carrying capacity influence on the incomes of seiners exploiting marine species in the Atlantic coast of Morocco", Mathematical Biosciences, 2018 Publication	<1 %

49

Jifa Jiang, Zhipeng Qiu. "The Complete Classification for Dynamics in a Nine-Dimensional West Nile Virus Model", SIAM Journal on Applied Mathematics, 2009

Publication

<1 %

50

Ma, Z.. "Stability analysis for differential infectivity epidemic models", Nonlinear Analysis: Real World Applications, 200312

Publication

<1 %

51

Ting Guo, Zhipeng Qiu, Libin Rong. "Modeling the role of macrophages in HIV persistence during antiretroviral therapy", Journal of Mathematical Biology, 2020

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On