

Iran J Public Health(18583)

External

Inbox



Iranian Journal of Public Health <ijph1971@gmail.com>

Sun, Dec 15, 2019,
12:37 PM

to me

Dear Corresponding Author

Thank you for your contribution with this journal. Please note that your paper Submitted

to this journal now has been evaluated by our panel of referees. Please read the attached comments of referee(s) carefully and reply to each in a **separate letter** to be sent for related referee. Please **highlight each revision by color or anything else in the text** so that the referee(s) can see the required changes.

Please note that for any reason if we do not receive any reply **in 60 days, we consider your manuscript "unprintable" and archive it.** If for accepted reasons, you are not capable to send us the reply in this course, let us know immediately.

Waiting your prompt reply.

Please send us your response just via replying this mail not via website.

Best Regards

Dr MB Rokni

Editor

E-mail: ijph@tums.ac.ir

Tell/Fax: +98-21-88950184



Iranian Journal of Public Health

Official Publication of the Iranian Public Health Association

Date: October 23, 2019

Art. No.: 18583

Ref No.: A

Art. Title: A high-altitude as the risk factor of Malaria in Indonesia

Reviewer Checklist for the Editor/Author

No	Requested Questions	Yes	No	Comments
1	Is this article ...			
	An original work?	<input checked="" type="checkbox"/>		
	A new subject?		<input checked="" type="checkbox"/>	
2	Is the title ...		<input checked="" type="checkbox"/>	See additional comments
3	Is the abstract ...		<input checked="" type="checkbox"/>	See additional comments
4	Is the introduction ...			
	Sufficient?	<input checked="" type="checkbox"/>		
	Informative?	<input checked="" type="checkbox"/>		
5	Are the materials and methods			
	Clear?	<input checked="" type="checkbox"/>		
	Adequate?	<input checked="" type="checkbox"/>		
	Ethical?	<input checked="" type="checkbox"/>		
6	Are the results...			
	Efficient?	<input checked="" type="checkbox"/>		
	Satisfactory in statistical analysis?	<input checked="" type="checkbox"/>		
	Well presented?	<input checked="" type="checkbox"/>		
7	Are the tables ...			
	Satisfactory?	<input checked="" type="checkbox"/>		
	Clear?	<input checked="" type="checkbox"/>		
	Necessary?	<input checked="" type="checkbox"/>		
	Adequate in number?	<input checked="" type="checkbox"/>		
8	Are the figures ...			
	Satisfactory?	<input checked="" type="checkbox"/>		
	Clear/in good quality of art?	<input checked="" type="checkbox"/>		

P.O. Box: 14155-6446, Tehran, Iran, Tel/Fax: + 98-21-88950184

Email: ijph@tums.ac.ir

URL: <http://ijph.ir>

System Preferences

		Necessary?	<input type="checkbox"/>	
		Adequate in number?	<input type="checkbox"/>	
9	Does the discussion ...	Include other relevant studies?	<input type="checkbox"/>	
		Suitable?	<input type="checkbox"/>	
		Sufficient?	<input type="checkbox"/>	
10	Are the references ...	Up to date?	<input checked="" type="checkbox"/>	
		Adequate in number?	<input type="checkbox"/>	
		Reduction in any part of the manuscript?	<input type="checkbox"/>	Reduce the text in results section
11	Would you suggest ...	Addition in any part of the manuscript?	<input type="checkbox"/>	Abstract needs rewriting
12	Is the quality of scientific language...	Satisfactory?	<input checked="" type="checkbox"/>	There are a lot of mistakes in the text it needs revised completely
13	Is the acknowledgement...	Include?	<input checked="" type="checkbox"/>	Acknowledgement should be added
14	Is the ethical consideration..	Include?	<input type="checkbox"/>	

Art. No: 18583

Ref No.: A

Reviewer comments for the author

Additional comments

It is better to consider this comments to improve this manuscript:

1- The title should be change and including whole of variable item in it not just altitude.

2- The abstract section needs completely change.

3- The number of malaria cases should be based on the last report of WHO.

4- Table 1:

Education section needs revised.

5- There are a lot of grammar mistake in the text. It needs to improve.

6- To prove the influences of altitude on number of malaria cases you need to have a P value on this variables:

(In this study, most of the cases were found with an average height of 311 meters with Standard Deviation (SD) 127.78 and controls 349 meters with SD = 198.97.)

System Preferences



Dear Corresponding authors

Please note the attached file as your edited article submitted to Iran J Public Health. Please amend the article accordingly and send the revised article in **maximum 2 days**.

Plz don't change the name of the file. Just add a word of **corrected** to the end of the file name.

Bes

IJPH Office

SOLIKHAH SOLIKHAH <solikhah@ikm.uad.ac.id>

Jan 17, 2020, 3:14 PM

to Iranian

Dear Editor

Considering comments from reviewers, we send our revised manuscript. Here noted revised of our manuscript below:

1. **Reviewers comment:** The title should be change and including whole of variable item in it not just altitude.

Answer: Thank you very much for your suggestion. Revised done

2. **Reviewers comment:** The abstract section needs completely change.

Answer: Abstract already revised in our manuscript

3. **Reviewers comment:** The number of malaria cases should be based on the last report of WHO

Answer: Thank you for your kindness and our revised manuscript done.

4. **Reviewers comment:** Table 1:Education section needs revised.

Answer: our revised manuscript highlighted with different colour.

5. **Reviewers comment:** There are a lot of grammar mistake in the text. It needs to improve.

Answer: Thank you for your suggestion. Our revised manuscript done

6. **Reviewers comment:** To prove the influences of altitude on number of malaria cases you need to have a p value on this variables: (In this study, most of the cases were found with an average height of 311 meters with Standard)

Answer: in this case we analyse and describe our data with univariate only, while our multivariate result of this study has been written in our manuscript with highlighted.

7. **Reviewer comment:** Acknowledgement should be added

Answer: our manuscript already done revised, Thank you very much.

Please kindly attached file below for our revised manuscript. Thank you very much,

Warm regards,

Solikhah Solikhah

<Referee Form18583.pdf>

Attachments area



Iranian Journal of Public Health <ijph1971@gmail.com>

Jan 18, 2020,
12:45 PM

to me

Thanks a lot.

On Fri, Jan 17, 2020 at 11:44 AM SOLIKHAH SOLIKHAH <solikhah@ikm.uad.ac.id> wrote:

Dear Editor

Considering comments from reviewers, we send our revised manuscript. Here noted revised of our manuscript below:

1. **Reviewers comment:** The title should be change and including whole of variable item in it not just altitude.

Answer: Thank you very much for your suggestion. Revised done

2. **Reviewers comment:** The abstract section needs completely change.

Answer: Abstract already revised in our manuscript

3. **Reviewers comment:** The number of malaria cases should be based on the last report of WHO

Answer: Thank you for your kindness and our revised manuscript done.

4. **Reviewers comment:** Table 1:Education section needs revised.

Answer: our revised manuscript highlighted with different colour.

5. **Reviewers comment:** There are a lot of grammar mistake in the text. It needs to improve.

Answer: Thank you for your suggestion. Our revised manuscript done

6. **Reviewers comment:** To prove the influences of altitude on number of malaria cases you need to have a p value on this variables: (In this study, most of the cases were found with an average height of 311 meters with Standard)

Answer: in this case we analyse and describe our data with univariate only, while our multivariate result of this study has been written in our manuscript with highlighted.

7. **Reviewer comment:** Acknowledgement should be added

Answer: our manuscript already done revised, Thank you very much.

Please kindly attached file below for our revised manuscript. Thank you very much,

Warm regards,
Solikhah Solikhah

[UNIVERSITAS AHMAD DAHLAN](#)

Kampus 1: Jln. Kapas No. 9 Yogyakarta

Kampus 2: Jl. Pramuka 42, Sidikan, Umbulharjo, Yogyakarta 55161

Kampus 3: Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta 55164

Kampus 4: Jl. Ringroad Selatan, Yogyakarta

Kampus 5: Jl. Ki Ageng Pemanahan 19, Yogyakarta

Kontak

Email: info@uad.ac.id

Telp. : (0274) 563515, 511830, 379418, 371120

Fax. : (0274) 564604

--

Dr. MB Rokni

Editor

Iran J Public Health

Professor

Ph.D. in Medical Parasitology

Head-Indexing & Visibility Committee- Eastern Mediterranean Association of Medical Editors (EMAME)

Member of WHO's Foodborne Disease Burden Epidemiology Reference Group (FERG)

P.O.BOX: 6446 Tehran 14155, IRAN

Telfax: +98 21 88950184



SOLIKHAH SOLIKHAH <solikhah@ikm.uad.ac.id>

Feb 29, 2020,
5:16 PM

to Iranian

Dear Editor in Chief

We have send our responses to reviewer on January 17, 2020.
Please let me know the progress.
Thank you very much

Best Regards,
Solikhah Solikhah



Iranian Journal of Public Health <ijph1971@gmail.com>

Mar 2, 2020,
4:58 PM

to me

Dear Dr Solikhah

Thank you for your cooperation with this journal. Regarding your manuscript submitted to IJPH, it is under peer review process and thus far we need the final approval by the referees' panel. Your revised manuscript must be approved by the referees in the end and this assists the Editorial Board to reach a clear-cut conclusion. Receiving this outcome we will let you know the result. Thank you for your positive reception.

Best

Best Regards

Dr MB Rokni
Editor
E-mail: ijph@tums.ac.ir
Tell/Fax: +98-21-88950184

Thank you for your response.
Noted with thanks.

SOLIKHAH SOLIKHAH <solikhah@ikm.uad.ac.id>

Thu, Aug 12, 2021,
9:48 PM

to Iranian

Dear Editor,
Please kindly find a revised manuscript. Thank you
Warm regards,
Dr Solikhah S

Dear Corresponding author

Would you please have a look at the comments in the text and correct the paper accordingly.

- ✓ **Because some modifications have been made on your paper, please revise the present issue only not your original one.**
- ✓ **Please do not delete any comments of the text.**

Please reply in maximum **3 days**.

Best regard
Dr MB Rokni
Editor
Iranian J Publ Health

Dear Editor

Please kindly check our revised manuscript with highlight. Thank you

Warm Regard
Dr. Solikhah S

Original Article

Risk Factors Analysis of Malaria Transmission at Cross-Boundaries Area in Menoreh Hills, Java, Indonesia

Dwi Sarwani Sri Rejeki ¹, *Solikhah Solikhah ², Siwi Pramatama M Wijayanti ¹

1. *Department of Public Health, Faculty of Health Sciences, Universitas Jenderal Soedirman, Purwokerto, Indonesia*

2. *Faculty of Public Health, Universitas Ahmad Dahlan, Yogyakarta, Indonesia*

*Corresponding Author: E-mail: solikhah@ikm.uad.ac.id

Abstract

Background: Risk factors of Malaria transmission at cross-boundaries area is important to be identified. This study aimed to identify the risk factors of Malaria transmission at cross-boundaries area in Menoreh Hills, Java, Indonesia.

Methods: The design of the study was an observational study with a case-control design. Data on malaria cases and controls were obtained from the Primary Health Care in Menoreh. All malaria positive patients with clinical and laboratory examinations recorded in health services during 1 Jan 2015-31 Dec 2015. Overall, 138 cases and 138 controls were included. Several variables were collected such as altitude, night out behavior, the use of mosquito nets, nighttime bed, travel history, mosquito bite prevention activities, cattle ownership, distance to mosquito breeding site, etc. Data were obtained by structured questionnaires and observation. Data were analyzed by univariate, bivariate and multivariate

Results: The altitude of house >500 m above sea level proved to be influential as a risk factor for Malaria (OR 3.62, 95% CI 1.61-8.16, $P=0.002$). Several variables were identified as a risk factor of Malaria such as the wall of the house from bamboo/wood, no insecticide and distance of house <100 m from mosquito breeding site.

Conclusion: An awareness for the local health sector particularly to provide a recommendation for house construction to protect a community from Malaria transmission.

Keywords: Altitude; Malaria; Menoreh; Risk factors; Indonesia

Introduction

Malaria is a vector-borne disease that has a major impact on global public health with around 3.4 billion people at risk. This disease is caused by five species of Plasmodium parasites, with *P. falciparum* and *P. vivax* are considered as the greatest threat. Plasmodium is transmitted by the bites of female Anopheles mosquitoes (1). In 2018, an estimated 228 million cases of malaria occurred worldwide compared to 2010 with 251 million cases and 231 million cases in 2017. Most of the malaria cases in 2018 were in the African Region (213 million or 93%), followed by Asia Southeast with 3.4% and Eastern Mediterranean Region with 2.1%. The incidence of malaria has decreased globally between 2010 and 2018, from 71 to 57 cases per 1000 population at risk (2). In Indonesia, Malaria has become the target of elimination gradually by 2030. Based on Indonesia **Regular Health Research (3)**, the incidence of Malaria in Indonesia in 2013 was 1.9%, with 10 provinces have Annual Parasitic Incidence (API) values between 1-5 per 1000 inhabitants, 6 provinces have API values > 5 per 1000 inhabitants, and there is area who have an API value of > 50 per 1000 inhabitants (Badan Penelitian dan Pengembangan Kesehatan Kemenkes RI, 2013). About 70.3% of the area in Indonesia categorized as unstable transmission and most of the population lives in this unstable transmission area (4).

Malaria transmission is determined by various factors such as the environment and behavior of people. Environmental factors which related to Malaria include temperature, rainfall, humidity, altitude, a composition of the vegetation index, conditions of the house (conditions of walls, roofs, floors, ventilation, the existence of livestock, density of the rooms, near breeding sites/rivers etc.) (5-7). The behavioral determinants that correlated with malaria such as the use of personal protection, the use of mosquito nets and insecticide spraying (8,9). Each country and areas have their own determinant risk depend on environmental, ecology and local socioeconomic factors (8). Integrated understanding about the patterns of malaria transmission, risk factors, the environment, socio-economic conditions of each region needs to be understood in the context of developing a malaria control program (10).

In addition to environmental and behavior factor, altitude is considered one of the factors that determine the geographical coverage of malaria transmission. This is related to the decrease in average temperature since Malaria decreases at increasing altitude. Malaria transmission rarely occurs at altitudes above 2000m above sea level (masl). However, currently, there are a higher density of Anopheles mosquito and Malaria incidence in altitude more than 1500 masl (10). The incidence of malaria is also higher at altitudes >100 m asl in hilly areas in

Bangladesh and statistically significant (11). In Irian Jaya Mountains, Indonesia, which was previously rare in malaria, currently Malaria is more commonly found (12). Indeed, it is interesting to examine whether high-altitude is a factor that plays a role in malaria incidence in Indonesia. The results of the study in Kenya showed that differences in altitude of place reflected the differences of Malaria prevalence. In general, at higher altitudes associated with malaria, this is related to lower temperatures which slows the development of Anopheles vectors and malaria transmission. In Kenya, the elevation is not significant for malaria prevalence on the district scale (13).

This study would explore the correlation of several environmental factors including altitude with Malaria incidence in the Menoreh region, Central Java Indonesia. In this region, the determinant of malaria is strongly influenced by ecosystems, vegetation and policies to eradicate malaria vector. Menoreh is an area that is administratively located in the border of Magelang Regency and Purworejo Regency, Central Java Province. Menoreh area is an area that forms a unique ecosystem that is the source of life for living things. This area is a karst and breccia area, as a buffer for cultural heritage objects, one of which is Borobudur Temple. Menoreh hills are a malaria-endemic area. In addition, the Menoreh region borders two surrounding endemic areas, namely in Purworejo and Magelang Districts. Vector eradication in this area facing difficulties since geographically this area covered by different local government. There were more than 3500 cases of malaria occurring in Menoreh (cross-border area of Central Java Province and Yogyakarta Special Region) for 11 yr from 2005-2015. Menoreh Hills were the highest contributor of malaria cases in the two regencies (39%-47% during 2011–2015) (14). Rainfall, humidity and migration affected the incidence of malaria (15). However, the information about the risk factors of Malaria transmission in cross-boundaries area is still limited.

We aimed to find out the risk factors which correlated with the Malaria transmission at cross-boundaries area in Menoreh Hills, Java, Indonesia. Results of this study would provide valuable information and input for Malaria prevention and control.

Materials and Methods

Study design and the area of study

The design of the study was an observational study with a case-control design. Data on malaria cases and controls were obtained from the Primary Health Care in Menoreh, while interviews and observations were carried out at each case and control's houses. The area of this study is the Menoreh ecosystem which located in an altitude of around 100-900 m above sea level, and there are Moderate Case Incidence/High Case Incidence villages for 5 consecutive years (2010-2014) and bordering each other. This region is suspected as a source of transmission every year on Java. There are two provinces bordering each other which are this region, namely the Province of Central Java, and the Special Region of Yogyakarta. Purworejo Regency and Kulonprogo Regency are regencies in the 2 Provinces bordered, which consist of 3 malaria-endemic sub-districts which include Kaligesing District, Bagelen District, and Kokap District. Population of cases is all malaria positive patients with clinical and laboratory examinations recorded in health services during 1 Jan 2015-31 Dec 2015. The control population, i.e. non-malaria sufferers, was recorded in the Primary Health Care and health services during the period of 1 Jan 2015-31 Dec 2015. Samples were 138 cases and 138 controls.

Most cases were distributed in Kaligesing Community Health Center (81 cases), Dadirejo Community Health Center (30 cases), Bagelen Community Health Center (15 cases), Kokap I Community Health Center (9 cases) and Kokap II Community Health Center (3 cases). The 5 villages with the most cases were Somongari, Kaligono, Semono, Sokoagung, and Kalirejo. Location of study could be seen in Fig. 1.




Fig. 1: Location of cross-boundary area in Menoreh hills Indonesia

Collecting data & analysis

Several variables were collected in this study such as altitude, night out behavior, the use of mosquito nets, nighttime bed, travel history, mosquito bite prevention activities, cattle ownership, distance to mosquito breeding site etc. Data were obtained by structured questionnaires and observation. Data were analyzed by univariate, bivariate and multivariate. Bivariate analysis to identify strong independent variables related to malaria. Multivariable analysis with logistic regression to determine the variables that most influence the incidence of malaria by first doing analysis in stages to ensure the existence of confounding or not. All data were analyzed with SPSS 19.0 for Windows.

Ethical approval

The protocol was approved by the ethical committee of the Faculty of Medicine Universitas Gadjah Mada number: KE/ FK/222/EC/2016.

Results

The results of the univariate analysis showed that among the respondents, the gender variables between cases and controls were almost the same. The average age of control was 35.2 yr, slightly more than the case of 32.1 years. The univariate analysis of behavior and socioeconomic characteristic could be seen in Table 1.

A night out behavior in the case and control groups were almost the same, more respondents who did not go out the night before the illness. In the ownership variables of the mosquito net the control group has more mosquito nets than the case group, and more cases groups two do not use bed nets increase the risk of exposure to mosquitoes. While for insecticide spraying, it is more in the control group than case group.

For the effort to prevent mosquito bites, the control group has made more prevention efforts through the use of repellents, the use of mosquito repellent, wearing long clothes at night and planting mosquito repellent plants. Mosquito bite prevention efforts are most often carried out in both the case and control groups, such as the use of long clothes at night. The percentage of case and control groups is almost the same in terms of the history of traveling to other regions, only about 40% have a history of traveling to malaria-endemic areas within 1 month before visiting the primary health care.

The education level of the control group was better than the case group. About 20% of the control group had a high school education and above and only 11% in the case group. The employment variables between the case group and control characteristics are almost the same, most as farmers and housewives/school children. In the physical condition of the house, the condition of the control group is better than that in the case group, there are about 62% of the controls that have permanent walls of brick/brick/brick, and in the case group around 40%. In the condition of the house ceiling and ventilation, both the case house and the control of most ceilings are open and there is no gauze ventilation, which allows the mosquito to enter the house. Most cases and control groups have livestock and are at risk because livestock is placed close to settlements which are <10 meters away. Goats and chickens are livestock mostly maintained by respondents. The physical condition of the house and environmental of the case and control groups are shown in Table 2.

Table 1: Behavior and socioeconomic characteristics

Variables	Group: n(%)	
	Cases	Controls
Night out behavior		
1. Yes, Night out	64 (46.4)	62 (44.9)
2. No	74 (53.6)	76 (55.1)
Ownership of mosquito nets		
1. 0 (do not have)	49 (35.5)	42 (30.4)
2. 1 - 2 pieces	76 (55.1)	84 (60.9)
3. 3 - 4 pieces	13 (26.1)	11 (8.0)
4. More than 4 pieces	0 (0.0)	1 (0.7)
Use of mosquito nets		
1. Do not use	83 (60.1)	86 (62.3)
2. Use	55 (39.9)	52 (38.7)
Insecticide Spraying in last 1 year		
1. No	118 (85.5)	100 (72.5)
2. Yes	20 (14.5)	38 (27.5)
Prevent to mosquito bites		
1. Did not do prevention	53 (38.4)	36 (26.1)
2. Do prevention	85 (61.6)	102 (73.9)
Mosquito bites prevention activities		
1. Using repellent	30 (21.7)	31 (22.5)
2. Using anti-mosquito drug	39 (28.3)	53 (38.4)
3. Wear long clothes	43 (31.2)	65 (47.1)
4. Plant mosquito repellent plants	0 (0.0)	1 (0.7)
Nighttime bed		
1. Outside the room	15 (10.9)	17 (12.3)
2. In the room	123 (89.1)	121 (87.7)
Travel history		
1. Yes, traveling	61 (44.2)	55 (39.9)
2. No	77 (55.8)	83(60.1)
Education		
1. Uneducated	19 (13.8)	13 (9.4)
2. Has not finished elementary school	35 (25.4)	28 (20.3)
3. Graduated from elementary school	38 (27.5)	37 (26.8)
4. Graduated from junior high school	30 (21.7)	34 (24.6)
	15 (10.9)	20 (14.5)
	1 (0.7)	0 (0.0)
	0 (0.0)	5 (3.6)

5.	Graduated from senior high school	0 (0.0)	1 (0.7)
Occupation			
1.	Farmer	67 (48.6)	53 (38.49)
2.	Plantation	0 (0.0)	1 (0.7)
3.	Labor	10 (7.2)	14 (10.1)
4.	Private work	4 (2.9)	9 (6.5)
5.	Traders	3 (2.2)	7 (5.1)
6.	Retired	0 (0.0)	1 (0.7)
7.	Civil servant	0 (0.0)	4 (2.9)
8.	unemployed	54 (39.1)	49 (3.5)

Table 2: Behavior, socioeconomic and environmental characteristics

Variables	Group: n(%)		
	Cases	Controls	
House wall			
1.	Bamboo	27 (19.6)	10 (7.2)
2.	Wood	58 (42.0)	42 (30.4)
3.	Brick	13 (9.4)	27 (19.6)
4.	Brick plaster	37 (26.8)	59 (42.8)
5.	Others, such as brick making	3 (2.2)	0 (0.0)
Roof/ Ceiling			
1.	Open, there is no roof of the ceiling	127 (92.0)	119 (86.2)
2.	Bamboo covered	1 (0.7)	2 (1.4)
3.	Wood covered	1 (0.7)	0 (0.0)
4.	Plasterboard covered	9 (6.5)	17 (12.3)
Gauze ventilation			
1.	Not available	135 (97.8)	134 (97.1)
2.	Available	3 (2.2)	4 (2.9)
Cattle ownership			
1.	Yes, at-risk (large cattle, such as cows, goats and enclosure <10 meters)	40(29.0)	43(31.2)
2.	No cattle	98(71.0)	95(68.8)
Type of cattle			
1.	Cow	3 (2.2)	2 (1.4)
2.	Goats	59 (42.8)	70 (50.7)
3.	Chicken/duck	74 (53.6)	65 (47.1)
Type of mosquito breeding place near the house			
1.	River	61 (29.3)	72 (36.6)
2.	Wellspring	57 (27.4)	63 (32.0)
3.	Small river	52 (25.0)	28(14.2)
4.	Basin/ Inundation	29 (13.9)	16(8.1)
5.	Pond	9 (4.3)	10(5.1)
6.	Rice fields	-	6(3.0)
7.	Lake	-	2(1.0)
Average distance		83.9 m	167.0 m
The distance of the house to mosquito breeding place			
1.	≤ 100 meters	114 (62.8)	96 (69.6)
2.	> 100 meters	24 (17.4)	42 (30.4)
The average distance to mosquito breeding place		83.9 m	167.0 m
Distance of the house to the Primary Health Care			
1.	< 4000 meters	65 (47.1)	83 (60.1)

2.	≥ 4000 meters	4646.4 m	4743.5 m
Average distance to Primary Health Care			
Altitude (above sea level)			
1.	≤ 500 meters asl	9(6.5)	37(26.8)
2.	> 500 meters asl	129(93.5)	101(73.2)
Average altitude		311 m	349 m

Mosquito breeding sites that are found in both cases and controls are almost the same, such as rivers, springs, water, inundation, ponds, basins/stalls. The average breeding distance to the house in the case group was closer (83.9 m) than the control group (167 m). As for the distance of the house to the Primary Health Care, in the case group (4646 m) was almost the same compared to the control group (4743.5 m). In the variable altitude of residence, the average house height of the case group (311 m asl) was lower in the control group (349 m asl). The physical

Bivariate analysis showed that there were 7 variables with a value of $P < 0.25$ which were spraying, preventing mosquito bites, education, walls of the house, roof/ceiling, a distance of the house to mosquito breeding and the height of the house. The next step is to enter the 7 variables into multivariate analysis with logistic regression analysis. The following are the best model results for factors that influence the incidence of malaria.

The best model results showed that the altitude of the house > 500 meters asl is a greater risk of occupants suffering from malaria compared to houses which located in altitude ≤ 500 meters asl with OR=3.625 ($P=0.002$). Details of the analysis presented in Table 3.

Table 3: The best analysis results of factors that influence the incidence of malaria

No.	Variables	B	P	OR	95% CI
1.	Altitude (> 500 meters)	1.288	0.002	3.625	1,610-8,160
2.	House's wall (bamboo/wood)	0.790	0.003	2.203	1,307-3,713
3.	Spraying (not sprayed in the last 1 year)	0.730	0.026	2.075	1,090-3,951
4.	Distance of house with mosquito breeding place (≤ 100 meters)	0.662	0.040	1.939	1,032-3,643

Discussion

This study highlighted that altitude of house location influences the Malaria incidence in the area of study. Menoreh ecosystem is a hilly area with the highest altitude of 1017 m above sea level. In this study, most of the cases were found with an average height of 311 meters with Standard Deviation (SD) 127.78 and controls 349 m with SD=198.97. The results of this study are in accordance with the results of a study in Bangladesh, which stated that forest and highland areas were at risk of malaria. This is related to the presence of Anopheles vectors found in forests and forest margins in Bangladesh (11). Altitude >100 meters is a risk factor for malaria in Bangladesh with OR=3.44. While in Uganda, higher altitudes of 1,500 m have been shown to be associated with lower malaria risk (16). This fact in accordance in Indonesia, which stated that the incidence of Malaria at an altitude of >1200 m is less than the incidence in 200 m (17). In the tropical area, altitude has a role in the spread of malaria in relation to the Anopheles mosquito. The higher a place will lower the average temperature. This temperature will affect the sporogony cycle and age of mosquitoes (12). The conducive area for Anopheles mosquito breeding site is at an altitude of 500-1000 m, because in that region has a warm temperature, which will shorten the duration of the extrinsic cycle (sporogony cycle). The study in Ethiopia which divides altitude into 6 categories, namely <1000 m, 1000-1500 m, 1500-2000 m, 2000-2500 m, 2500-3000 m and > 3000 m, showed that the increasing prevalence of malaria prevalence diminishes. Malaria incidence is more common in areas with a height of <1000 m (5). *An. aconitus* malaria vector was found in rice field ecosystems at an altitude of 400-1000 meters (16). Our study, we found that the Menoreh ecosystem which showed that the incidence of malaria is more prevalent in areas with an altitude between 500-800 m. The results also similar to a study in Tanzania (18) and Malawi (19) which stated that the prevalence of malaria parasites decreased with increasing the altitude of residence. The most prevalence of malaria is found in residents with a height of 500 m in Tanzania and 650 m in Malawi, and decreasing to a height of 2000 m.

The rivers in the area of study flow during the rainy season and the dry season, and in those places where mosquitoes breed. There are many rivers and rivers (small rivers) at the research location and the distance of the river/river to the houses of people close enough (<200 meters) is still within the range of flying Anopheles mosquitoes are at risk of increasing malaria cases. *Anopheles. Balabacencis* larvae breeding place usually is a pool of water covered by shrubs in the form of a water basin/basin in the river, in a pond, open puddles like a puddle. While *An. maculatus* breeds in mountainous areas, such as small rivers, springs, ponds, puddles on the

banks of rivers and seepage of fields whose water flows slowly (12). A natural resting place for Malaria vectors in Kokap Subdistrict, Kulonprogo Regency, which are shaded holes and used as household waste disposal sites (7).

The advantage of this research is that this study uses a comprehensive analysis of malaria risk factors which include environmental, behavioral and socioeconomic factors. In the multivariate analysis, the confound control is carried out, so that internal validity is better. Meanwhile, the limitation of this study is that this study uses a case-control study design, namely with reverse inference methodology starting from new diseases traced to exposure, so the possibility of the emergence of recall bias is greater, such as mosquito bite prevention behavior, night out behavior and use of mosquito nets. The malaria case used was only malaria cases recorded in health services based on micro-examination, where there may still be asymptomatic cases not reported in health services. Finally, we concluded the main determinant of Malaria in Menoreh's hilly ecosystem is altitude. In this study, several variables were identified as a risk factor of Malaria such as the wall of the house from bamboo/wood, no insecticide and distance of house <100 m from mosquito breeding site. This result should be an awareness for local health sector particularly to provide a recommendation for house construction for a new house to protect a community from Malaria. It is also important to increase the prevention effort of Malaria to minimize them from Anopheles mosquito bites.

Ethical considerations

Ethical issues (Including plagiarism, informed consent, misconduct, data fabrication and/or falsification, double publication and/or submission, redundancy, etc.) have been completely observed by the authors.

Funding

This study was supported by the Ministry of Research, Technology and Higher Education of the Republic of Indonesia.

Acknowledgements

The authors acknowledge the contribution of all participants in the Kokap District and all staffs of Primary Health Care in Menoreh.

Conflict of interest

All authors declare that there is no conflict of interest.

References

1. Kantele A, Jokiranta TS (2011). Review of cases with the emerging fifth human malaria parasite, *Plasmodium knowlesi*. *Clin Infect Dis*, 52(11): 1356–62.
2. Global Malaria Programme, WHO (2019). WHO Malaria Policy Advisory Committee (MPAC) meeting: Meeting Report. 2019;(October):1–23. Available from: <http://www.who.int/malaria/mpac/mpacmembers/en/>
3. Badan penelitian dan pengembangan kesehatan Kemenkes RI (2013). Riset Kesehatan Dasar. Kementerian Kesehatan Republik Indonesia, Jakarta, pp.: 2.
4. Elyazar IRF, Gething PW, Patil AP, et al (2011). *Plasmodium falciparum* Malaria Endemicity in Indonesia in 2010. *PLOS ONE*, 6(6): e21315.
5. Graves PM, Richards FO, Ngondi J, et al (2009). Individual, household and environmental risk factors for malaria infection in Amhara, Oromia and SNNP regions of Ethiopia. *Trans R Soc Trop Med Hyg*, 103(12): 1211–20.
6. Guthmann JP, Hall AJ, Jaffar S, et al (2001). Environmental risk factors for clinical malaria: a case-control study in the Grau region of Peru. *Trans R Soc Trop Med Hyg*, 95(6): 577–83.
7. Handayani FD, Darwin A (2006). Habitat vektor malaria di daerah endemis Kecamatan Kokap Kabupaten kulonprogo Propinsi Daerah Istimewa Yogyakarta. *Indones J Health Ecol*, 5(2): 438–446.
8. Yadav K, Dhiman S, Rabha B, et al (2014). Socio-economic determinants for malaria transmission risk in an endemic primary health centre in Assam, India. *Infect Dis Poverty*, 3(19): 19.
9. Protopopoff N, Bortel WV, Speybroeck N, et al (2009). Ranking Malaria Risk Factors to Guide Malaria Control Efforts in African Highlands. *PLOS ONE*, 4(11): e8022.
10. De Silva PM, Marshall JM (2012). Factors Contributing to Urban Malaria Transmission in Sub-Saharan Africa: A Systematic Review. *J Trop Med*, 2012(Article ID 819563): 1–10.
11. Haque U, Sunahara T, Hashizume M, et al (2011). Malaria Prevalence, Risk Factors and Spatial Distribution in a Hilly Forest Area of Bangladesh. *PLOS ONE*, 6(4): e18908.
12. Susanna D (2011). *Dinamika Penularan Malaria*. UI-Press, Jakarta, pp.: 48.
13. Ernst KC, Adoka SO, Kowuor DO, et al (2006). Malaria hotspot areas in a highland Kenya site are consistent in epidemic and non-epidemic years and are associated with ecological factors. *Malar J*, 5(1): 78.

- 14 Rejeki DSS, Fuad A, Widartono BS, et al (2019). Spatiotemporal patterns of malaria at cross - boundaries area in Menoreh Hills , Java , Indonesia. *Malaria Journal*, 18(80): 1–12.
15. Rejeki DSS, Nurhayati N, Aji B, et al (2018). A Time Series Analysis: Weather Factors, Human Migration and Malaria Cases in Endemic Area of Purworejo, Indonesia, 2005–2014. *Iran J Public Health*, 47(4): 499–509.
16. De Beaudrap P, Nabasumba C, Grandesso F, et al (2011). Heterogeneous decrease in malaria prevalence in children over a six-year period in south-western Uganda. *Malar J*, 10(132): 2–9.
17. Hanandita W, Tampubolon G (2016). Geography and social distribution of malaria in Indonesian Papua: a cross-sectional study. *Int J Health Geogr*, 15(13): 2–15.
18. Drakeley CJ, Carneiro I, Reyburn H, et al (2005). Altitude-dependent and -independent variations in *Plasmodium falciparum* prevalence in northeastern Tanzania. *J Infect Dis*, 191(10): 1589–98.
19. Kazembe LN, Kleinschmidt I, Holtz TH, et al (2006). Spatial analysis and mapping of malaria risk in Malawi using point-referenced prevalence of infection data. *Int J Health Geogr*, 5(41): 1–9.

Galley proof of your paper in IJPH

External

Inbox



Iranian Journal of Public Health <ijph1971@gmail.com>

Mon, Aug 30, 2021,
12:11 AM

to me

Dear Corresponding Author

Please note the attached file as the galley proof of your manuscript submitted to **Iranian J Publ Health** and let me know if it is OK or need amendment. Please **highlight** the background for any changes by **Yellow** color (Don't color the text).

In this step, **absolutely**, you **cannot add** any authors or **large sentences**, besides you are not allowed to change the corresponding author. Please check the MS only in terms of any typos, grammar mistake, reference number, Table and Figure numbers etc.

Considering that the journal is published monthly, if for any reason, after **48 hours from now**, we do not receive any reply of you, we consider your manuscript **approved** by you, and no changes will be accepted after that. Surely, after that, you will be responsible for any mistakes in the paper.

Note: After approving the galley proof no changes would be accepted and the responsibility would be on your side.

توجه مهم: با توجه به تمهیدات پایگاه های اطلاعاتی بعد از تایید گالی پروف به هیچ عنوان امکان تغییر مقاله نیست. لطفا دقت بفرمایید چون مسنولیت با خود شماست.

IJPH Office

--

Dr. MB Rokni

Editor

Iran J Public Health

Professor

Ph.D. in Medical Parasitology

Head-Indexing & Visibility Committee- Eastern Mediterranean Association of Medical Editors (EMAME)

Member of WHO's Foodborne Disease Burden Epidemiology Reference Group (FERG)

P.O.BOX: 6446 Tehran 14155, IRAN

Telfax: +98 21 88950184



SOLIKHAH SOLIKHAH <solikhah@ikm.uad.ac.id>

Mon, Aug 30, 2021,
9:06 AM

to Iranian

Dear Editor,
Thank you for nice job, however please kindly arrange neatly for yellow highlight.

Thank you
Warm regards,
Dr. Solikhah S



Iranian Journal of Public Health <ijph1971@gmail.com>

Mon, Aug 30, 2021,
10:06 AM

to me

Dear Dr
Thanks a lot