

Depression among diabetes mellitus patients

By Putri Azizah

Depression among diabetes mellitus patients: A study of the protective factors

Putri Nur Azizah¹, Herlina Siwi Widiana², Siti Urbayatun³

^{1,2,3}Faculty of Psychology, Universitas Ahmad Dahlan, Indonesia

Article Info

Article history:

Received

Revised

Accepted

Keywords:

Depression

Diabetes mellitus

Protective factors

Resilience

Self-esteem

ABSTRACT

Diabetes mellitus is a chronic disease with patients that continue to increase per year. This study aimed to understand the role of self-esteem and religious orientation on depression with resilience as a mediator among patients with diabetes mellitus (DM). A total of 100 patients were recruited from three public health centers. The Center for Epidemiologic Studies Depression Scale (CES-D) was used to measure depression. In contrast, self-esteem was measured using two aspects of Rosenberg's self-esteem scale, namely self-liking and self-competence. Religious orientation was measured using Allport and Ross's religious orientation scale, while resilience was measured using Connor and Davidson's resilience scale. The results were analyzed using path analyses. The study found that resilience mediated the relationship between self-esteem and depression among DM patients. Self-esteem was also shown to correlate with resilience, and resilience also showed a significant relationship with depression. In contrast to initial predictions, however, there was no significant effect of religious orientation on resilience. The research implications suggest that resilience serves as an important protective factor toward depression among patients suffering from DM.

This is an open access article under the CC BY-SA license.



Corresponding Author:

Herlina Siwi Widiana,

Faculty of Psychology,

Universitas Ahmad Dahlan,

Jl. Kapas No. 9 Semaki Yogyakarta 55166, Indonesia.

Email: herlina.widiana@psy.uad.ac.id

1. INTRODUCTION

In 2020, Wuhan was struck by a peculiar case of pneumonia caused by the Sars-Cov-19 virus [1]. Patients with a history of hypertension and diabetes mellitus are at particular risk from COVID-19 infection [2]. Diabetes is a chronic disease, and according to the International Diabetes Foundation, it constitutes a major severe disease. Diabetes occurs due to genetic abnormalities in metabolism, which result in increases in blood sugar levels; however, it may also be marked by deformities in insulin production [3]. As a result, the total number of diabetes patients is increasing, particularly in developing countries. Indonesia being among these developing countries, has shown that 6% of people with diabetes are above 15 years old; however, only 9% receive medical care [4].

Diabetes patients must undergo long-term medication to prevent complications in the body due to insulin imbalance [5]. However, prolonged medication may lead to deteriorating mental health marked by feelings of hopelessness and a fractured self-concept, leading to depression [6]. It is common among diabetes patients to experience emotional problems, such as stress due to lifestyle changes. Stress can also emerge among DM patients due to its symptoms, which may lead to depressive symptoms or worry since the patient is unable to carry on with normal life activities. A study on 122 DM patients in Padang found that 43.4% of the patients experienced mild depression [7]. Depression among these patients was marked by unhappiness,

difficulties sleeping, and intentions to commit suicide [7]. The stress which emerges due to the illness also impacts the personal management of diabetes [8]. This research showed that people with depressive symptoms were more likely to have poor compliance and poor abilities to control their glycemic concentration in their blood, while patients without depressive symptoms were more able to regulate their lives and glycemic concentration in their blood. This result indicates the importance of monitoring depressive symptoms among DM patients [9].

DM patients who have sufficient knowledge about their illness can also minimize the risks of developing depression [10]. Depression is a mood disorder that involves emotional problems (prolonged sadness), lack of interest in daily activities, cognitions (helplessness), and affects bodily functions (easily fatigued and loss of appetite) [11]. When diabetes patients experience depression, this leads to a poor diet and other complications and disorders, for example, psychological disorders and emotional problems. In addition, emotional problems can affect the quality of life, poor self-grooming, poor management of glycemic levels, and increases the risk of death [12].

Depression is a general clinical disorder; however, it is not easily detected like a physical illness, and therefore patients with depression are not seriously treated. Some factors affect the extent of depression in DM patients, namely economic condition, family status, obesity, smoking habits, and poor lifestyle [13]. The prevalence of depression among diabetes patients has increased twofold, where research has shown an increase of 24-30% [3]. In addition, the prevalence of comorbidities between depression and other chronic diseases is higher than the prevalence of depression [14]. This prevalence shows that having a chronic disease for a long period without the healing potential may lead to depression.

Each person has once experienced an adverse life event that leads to stress, and living with diabetes mellitus may constitute such a negative event. The stress experienced by each diabetes patient can lead to some negative impacts, for example, declines in well-being, post-traumatic stress disorder, anxiety disorders, and depression [15]. However, each person has different abilities in managing stress, otherwise known as resilience. Resilience has many meanings, however, it concerns positive adaptation in managing risk or stress [16]. Resilience refers to a person's ability to maintain a stable and healthy psychological and physical state when dealing with a stressful situation [17]. Resilience affects DM patients' ability to control glucose levels and protects them from the adverse risks of psychological distress on blood sugar levels [18].

Resilience comes from three factors, namely "I am," "I have," and "I can" [19]. The factor "I am" comes from within the individual and can form self-esteem or self-worth. Self-esteem refers to beliefs that one can think and survive in managing life's challenges and beliefs that one has the right to succeed, be happy and content, and enjoy the outcome of their achievements. According to some research, self-esteem plays a role in life quality and emotional health [20]. Among individuals with comorbid depression, there is a cycle between inferiority and tendencies to experience mental disorders [21].

A study in Iran found that religious orientation positively correlated with self-esteem [22]. The study evaluated the aspects that affected religious orientation, for example, anxiety and self-esteem. The psychological distress experienced by diabetes patients was associated with religious observance, whereby past research has shown that a lack of spiritual quality may lead to depression among patients [23]. Attitudes were associated with two religious aspects, namely intrinsic and extrinsic orientation [24]. Individuals tend to view religiosity intrinsically when their religious practices and motivation originate from personal beliefs. People who adhere to this way of life follow one that strictly aligns with their religion. The research has shown that religious beliefs play a role in dealing with unpleasant situations and therefore make it relevant to mental health [25]. In addition, religion and spirituality, which comprises beliefs of a divine relationship with God, increase a person's resilience [26].

The intrinsic aspect of religious orientation plays a key role in mental health because it can ameliorate stress [27]. Koenig [28] found a relationship between religious orientation and positive psychological adaptation consisting of life satisfaction, happiness, optimism, life goals, and stress reduction. Religiosity, particularly the self-oriented type, was associated with patients who have life goals, are motivated to engage with daily activities, and reduce the probability of experiencing depression [29].

From the elaborations above, it becomes important to understand the factors that affect depression among diabetes mellitus patients. Furthermore, this study tests the role of religious orientation, self-esteem toward depression among diabetes mellitus patients with resilience as the mediator that previous studies have not explored. Thus, the study may contribute to knowledge development, mainly related to protective factors of depression among patients with diabetes mellitus.

2. RESEARCH METHOD

The respondents in this study were diabetes mellitus patients aged from 21-65 years. The respondents were outpatients from three public health centres (*Puskesmas*) in Yogyakarta namely *Puskesmas* Kotagede I, *Puskesmas* Gondomanan, *Puskesmas* Mergangsan, and *Puskesmas* Jetis. Table 1 shows the demographic data

of the respondents. The scales were administered to the respondents directly in the public health centers and using Google Forms distributed through WhatsApp groups. All respondents agreed to complete the scales and completed an informed consent form that the researcher provided. The Center for Epidemiologic Studies Depression Scale (CES-D) was used to measure depression. The scale was translated into Indonesian for this research. In [11] study, the item discrimination power ranged from .44-.54, with a reliability of .85. Furthermore, Rosenberg's self-esteem scale was used to measure self-esteem. The discrimination power of the items ranged from .415-.703 with reliability of .858. Religious orientation was measured using Allport & Ross's scale [24], which consists of 14 items, and the orientation scale had an item discriminatory power ranging from .385-.897 and a reliability of .86. Finally, resilience was measured using the CD-RISC, composed of 25 items and translated into Indonesia [30]. In this study, the item's discriminatory power ranged from .560-.905 and had a reliability coefficient of .975. The current research used path analysis to analyze the data with IBM SPSS AMOS 22. Path analysis is a technique that analyzes causal relations when an independent variable affects a dependent variable not only directly but also indirectly. Path analysis is a development of the regression model used to test the fit of two models being compared.

Table 1. Demographic characteristics

Category	Frequency	Percentage
Gender		
Male	38	38%
Female	62	62%
Marital status		
Married	86	86%
Not married	14	14%
Most Recent Education		
Elementary school	24	24%
Junior High School	15	15%
Senior High School	36	36%
Diploma	3	3%
Undergraduate	17	17%
Others	5	5%
Duration as DM patient		
< 1 years	20	20%
1-5 years	43	43%
>5 years	37	37%
Treatment		
Medicine	83	83%
Insulin	16	16%
Others	1	1%
Diabetes Type		
Type I	8	8%
Type II	22	22%
Others (Do not know)	70	70%

3. RESULTS AND DISCUSSIONS

Table 2 shows that a majority of the respondents experienced depression in the mild category (47%). In contrast, a majority of the subjects had high self-esteem (97%), high religious orientation (100%), and high resilience (97%). The subjects showed high self-esteem, which supports past research, which suggests that DM patients have high external motivation and can maintain their self-esteem. This condition is marked by a sense of appreciation that can minimize the risks of developing emotional problems, such as helplessness, anxiety, and depression among patients with chronic illness [31]. In addition, the patients also showed high resilience (97%), which indicates that patients have the insight to adapt their lifestyle to manage the adverse effects of diabetes. This resilience would motivate the patient to take care of themselves and further lead to resilience [32]. The religious orientation shown by the participants was also in the high category, where patients use religious orientation to affect all aspects of life. They are therefore able to control their anger and be accountable for their actions [33].

Table 2. Categorization of subjects based on hypothetical data

Scale	Category	Score Intervals	Frequency	Percentage
Depression	Severe	>24	20	20%
	Moderate	16 < X ≤ 23	33	33%
	Mild	< 16	47	47%
Self-esteem	High	>21	97	97%
	Moderate	14 < X ≤ 21	3	3%
	Low	< 14	0	0%
Religious orientation	High	>28	100	100%
	Moderate	17 < X ≤ 28	0	0%
	Low	< 17	0	0%
Resilience	High	>91	99	99%
	Moderate	58 < X ≤ 91	0	0%
	Low	<58	1	1%

The path analyses showed that the structural model in Figure 1 meets the criteria for model fit, shown by the Chi-square value of 2.695 with a $p = .260$.




Figure 1. Results of analysis data using a path model

Furthermore, the values of GFI = .987, AGFI = .934, NFI = .942, TLI = .948, a ¹⁵RMSEA = .059 indicates the indexes meets with the cut off. The overall goodness of fit index is presented in Table 3.

Table 3. Index of good fit for the structural equation model

Goodness of fit index	Value	Cut-off value	Conclusion
CMIN/DF	1.347	≤5.00	Fit
Probabilitas	.260	≥.05	Fit
GFI	.987	≥.90	Fit
AGFI	.934	≥.90	Fit
TLI	.948	≥.90	Fit
NFI	.942	≥.90	Fit
PGFI	.197	0-1.0	Fit
RMSEA	.059	≤.08	Fit

Table 4 shows resilience has a negatively sign¹⁷ificant relationship with depression. The higher resilience among DM patients, the lower depression will be. This result is in line with the finding from previous research that resilience has a significant influence on reducing depression levels [34]. A person with resilience can have different²⁰ views when facing problems, fostering a strength to encounter the problems. A previous study on patients with breast cancer also found resilience to become a protective factor of depression [35].

This current study shows that most respondents have mild depression (47%), and only 20% have severe depression. Comorbid between depression and diabetes mellitus is common among adult patients, as they face multiple stressors in their life. For example, one condition is a denial of the illness among DM patients, resulting in their not adhering to diet control; therefore, their lifestyle is less effective [36].

Table 4. ¹⁶ression weights

Variable		Estimate	S.E.	C.R.	P
Resilience <--	Self-esteem	2.116	.417	5.069	***
Resilience <--	Religious orientation	-.003	.404	-.009	.993
Depression <--	Resilience	-.206	.067	-3.056	.002

¹ Based on the results presented in Table 4, self-esteem has a significant relationship with resilien¹. Self-esteem refers to a person's evaluation of his or her characteristics and abilities, and it supports the

performance of adaptive behaviors, which may strengthen resilience among DM patients [37]. Resilience is characterized by the ability to rebound and function normally following an unpleasant situation. Self-esteem is one of the characteristics of resilient people because it allows for better adaptation to the environment and has a large capacity to confront stress [38]. In addition, a previous study among DM patients in Central Java found self-confidence and self-esteem will encourage them to think positively [39].

The results showed that there was no relationship between religious orientation and resilience. Both extrinsic and intrinsic religious orientation did not affect resilience. Research in Africa showed that caregivers might negatively affect a person's resilience when they excessively communicate about religious ideas [40]. The current research is limited in that from the total 18 public health centers in Yogyakarta, and only 3 approved the research. Furthermore, more than half of the participants wanted the researcher to read the materials due to impaired vision.

4. CONCLUSION

There was a relationship between resilience and depression among DM patients. DM patients who can deal with and rebound from the DM diagnosis can positively adapt their physical conditions and prevent developing depression. In addition, self-esteem is associated with resilience. DM patients have good self-esteem, and they would try to make a positive image of themselves and adapt to the changes that occur around them, which provides the basis for a resilient individual. Self-esteem is one of the important factors in forming resilience. When individuals are confident of their abilities, this will increase their resilience, which eventually becomes a protective factor toward depression among DM patients.

ACKNOWLEDGEMENTS

The authors would like to thank the Directorate of Research and Community Service, Deputy of Strengthening Research and Development, Ministry of Research and Technology/National Research and Innovation Agency for providing this research grant (Letter of Grant Agreement on Implementation of Postgraduate Thesis Grant Research No. PTM-029/SKPP.TT/LPPM UAD/VI/2020).

REFERENCES

- [1] H. A. Rothan and S. N. Byrareddy, "The epidemiology and pathogenesis of coronavirus disease (COVID-19) outbreak," *J. Autoimmun.*, vol. 109, p. 102433, 2020.
- [2] L. Fang, G. Karakiulakis, and M. Roth, "Are patients with hypertension and diabetes mellitus at increased risk for COVID-19 infection? The Lancet," *Respir. Med.*, vol. 8, no. 4, p. e21, 2020.
- [3] W. Katon, M. Maj, and N. Sartorius, *Depression and diabetes*. UK: Wiley Blackwell, 2010.
- [4] Ministry-of-Health, "Basic health research," 2018. [Online]. Available: <https://www.kemkes.go.id/resources/download/info-terkini/hasil-risikesdas-2018.pdf>.
- [5] W. Tarwoto and I. M. Taufiq, *Keperawatan medikal bedah gangguan sistem endokrin (Surgical medical nursing of endocrine system disorders)*. Jakarta: TIM, 2012.
- [6] Nurbani, "Terapi spiritual emotional freedom technique mengurangi depresi pada pasien diabetes mellitus (Emotional freedom technique spiritual therapy reduces depression in diabetes mellitus patients)," *J. Keperawatan Jiwa*, vol. 8, no. 2, pp. 129–138, 2020.
- [7] R. Sarfika, "Hubungan keputusasaan dengan depresi pada pasien diabetes melitus di Padang (Relationship between hopelessness and depression in diabetes mellitus patients in Padang)," *NERS J. Keperawatan*, vol. 15, no. 1, pp. 14–24, 2019.
- [8] R. D. Goldney, P. J. Phillips, L. J. Fisher, and D. H. Wilson, "Diabetes, depression, and quality of life: a population study," *Diabetes Care*, vol. 27, no. 5, pp. 1066–1070, 2004.
- [9] K. J. W. Baucom, S. L. Turner, E. L. Tracy, C. A. Berg, and D. J. Wiebe, "Depressive symptoms and diabetes management from late adolescence to emerging adulthood," *Heal. Psychol.*, vol. 37, no. 8, pp. 716–724, 2018.
- [10] M. K. Putri, T. A. Larasati, and D. I. Angraini, "Tingkat depresi dan kontrol gula darah puasa pada penderita diabetes melitus tipe II di Bandar Lampung (Level of depression and control of fasting blood sugar among patients with diabetes mellitus type II in Bandar Lampung)," *J. Agromedicine*, vol. 5, no. 1, pp. 395–398, 2018.
- [11] C. Wade, C. Tavis, and M. Garry, *Psychology*, Eleventh e. Pearson, 2014.
- [12] M. J. Knol, J. W. Twisk, A. T. Beekman, R. J. Heine, F. J. Snoek, and F. Pouwer, "Depression as a risk factor for the onset of type 2 diabetes mellitus. A meta-analysis," *Diabetologia*, vol. 49, no. 5, pp. 837–845, 2006.
- [13] J. S. Gonzales, L. Fisher, and W. H. Polonsky, "Depression in diabetes: have we been missing something important?," *Diabetes Care*, vol. 34, no. 1, pp. 236–239, 2011.
- [14] S. Moussavi, S. Chatterji, E. Verdes, A. Tandon, V. Patel, and B. Ustun, "Depression, chronic diseases, and decrements in health: results from the World Health Surveys," *Lancet (London, England)*, vol. 370, no. 9590, pp. 851–858, 2007.
- [15] A. Monat, R. S. Lazarus, and G. Reeve, *The Praeger handbook on stress and coping*. Praeger, 2007.

- [16] J. W. Reich, A. J. Zautra, and J. S. Hall, *Handbook of adult resilience*. The Guildford Press, 2010.
- [17] G. A. Bonanno, "Loss, Trauma, and Human Resilience: Have We Underestimated the Human Capacity to Thrive After Extremely Aversive Events?," *Am. Psychol.*, vol. 59, no. 1, pp. 20–28, 2004.
- [18] C. N. Massey, E. H. Feig, L. Duque-Serrano, and J. C. Huffman, "Psychological well-being and type 2 diabetes," *Curr. Res. Diabetes Obes. J.*, vol. 4, no. 4, p. 555641, 2017.
- [19] E. H. Grotberg, "The international resilience project: Research, application, and policy," *Pap. Present. Symp. Int. Stress e Violencia*, pp. 27–30, 1995.
- [20] M. Soheylizad, A. E. Yahaghi, D. Amini, and B. Gholamaliev, "Relationship between self-esteem, resilience, and quality of life in patients with type 2 diabetes in Hamadan," *Pojouhan Sci. J.*, vol. 15, no. 1, pp. 1–8, 2016.
- [21] M. H. Guindon, *Self-esteem across the lifespan*. Routledge, 2010.
- [22] S. Mousavimoghadam, M. Nourmohammadil, N. Ranjbarian, and M. Rashidahal, "The relationship between religious orientation and students self esteem," *Heal. Spiritual, Med Ethics*, vol. 1, no. 4, pp. 16–21, 2014.
- [23] N. Jafari, Z. Farajzadegan, A. Loghmani, M. Majlesi, and N. Jafari, "Spiritual well-being and quality of life of Iranian adults with type 2 diabetes," *Evidence-based Complement. Altern. Med. eCAM*, vol. 2014, p. 619028, 2014.
- [24] G. W. Allport and J. M. Ross, "Personal religious orientation and prejudice," *J. Pers. Soc. Psychol.*, vol. 5, no. 4, pp. 432–443, 1967.
- [25] M. Mahmoodabad *et al.*, "Extrinsic or intrinsic religious orientation may have an impact on mental health," *Res. J. Med. Sci.*, vol. 10, no. 4, pp. 232–236, 2016.
- [26] F. Dehgani and V. Andishmand, "The relationship of religious orientation and spiritual health to resilience among high school Sophomores in Kerman," *J. Res. Relig. Heal.*, vol. 3, no. 4, pp. 66–67, 2017.
- [27] G. H. Javanmard, "Religious beliefs and resilience in academic students," *Procedia - Soc. Behav. Sci.*, vol. 84, no. 2013, pp. 744–748, 2013.
- [28] H. G. Koenig, "Research on religion, spirituality, and mental health: a review," *Can. J. Psychiatry*, vol. 54, no. 5, pp. 283–291, 2009.
- [29] M. Heidarzadeh and M. Aghamohammadi, "Spiritual growth in patients with type II diabetes mellitus: A qualitative study," *J. Res. Dev. Nurs. Midwifery*, vol. 14, no. 2, pp. 34–44, 2017.
- [30] Y. Kurniawan and N. Noviza, "Peningkatan resiliensi pada penyintas kekerasan terhadap perempuan berbasis terapi kelompok pendukung (Support group therapy to increase resilience among survivor of violence against women)," *Psikohumaniora J. Penelit. Psikol.*, vol. 2, no. 2, pp. 125–142, 2017.
- [31] S. E. Ernawati and T. Kurniawan, "Pengaruh Motivasi Internal dan Eksternal terhadap Diabetes Self Management di Wilayah Kecamatan Gombong (The effect of internal and external motivation on diabetes self management in Gombong district)," *JSK*, vol. 1, no. 2, pp. 55–62, 2015.
- [32] U. Fadila and H. Laksmiwati, "Perbedaan resiliensi pada penderita diabetes mellitus tipe II berdasarkan jenis kelamin (The difference of resilience among patients with type II diabetes mellitus based on gender)," *Character*, vol. 3, no. 2, pp. 1–6, 2014.
- [33] N. Mazlom, M. Ardekani, and A. Dadgari, "Relationship between the religion orientation and coping with diabetes in patients with type 2 diabetes," *Iran. J. Diabetes Obes.*, vol. 5, no. 1, pp. 33–39, 2013.
- [34] E. Mujahidah and R. A. Listiyandini, "Pengaruh resiliensi dan empati terhadap gejala depresi pada remaja (The effect of resilience and empathy on depression among adolescents)," *J. Psikol.*, vol. 14, no. 1, pp. 60–75, 2018.
- [35] G. Ristevska-Dimitrovska, P. Stefanovski, S. Smichkoska, M. Raleva, and B. Dejanova, "Depression and Resilience in Breast Cancer Patients," *Open access Maced. J. Med. Sci.*, vol. 3, no. 4, pp. 661–665, 2015.
- [36] G. Semiardji, *Stres emosional pada penyandang diabetes (Emotional stress among patients with diabetes)*. Jakarta: Balai Penerbit FKUI, 2009.
- [37] C. J. Mruk, *Self-esteem and positive psychology*. New York: Springer Publishing Company, 2015.
- [38] H. Setiorini, T. Pahria, and T. Sutini, "Gambaran harga diri pasien diabetes mellitus yang mengalami ulkus diabetik di Rumah Perawatan Luka Bandung (Description of self-esteem among patients with diabetes mellitus who have diabetic ulcers in Rumah Perawatan Luka Bandung)," *J. Keperawatan Komprehensif*, vol. 5, no. 2, pp. 118–126, 2019.
- [39] P. H. Livana, I. P. Sari, and Hermanto, "Gambaran tingkat depresi pasien diabetes mellitus di Kabupaten Kendal (Description of depression level among patients with diabetes mellitus in Kendal district)," *J. Kesehat. Poltekkes Ternate*, vol. 11, no. 2, pp. 48–57, 2018.
- [40] M. Mhaka-Mutepfa and T. Maundeni, "The role of faith (spirituality/religion) in resilience in Sub-Saharan African children," *Int. Community Soc. Dev.*, vol. 1, no. 3, pp. 211–233, 2019.

Depression among diabetes mellitus patients

ORIGINALITY REPORT

13%

SIMILARITY INDEX

PRIMARY SOURCES

1	www.neliti.com Internet	34 words — 1%
2	Triantoro Safaria, Hadi Suyono. "The role of parent-child relationship, school climate, happiness, and empathy to predict cyberbullying behavior", International Journal of Evaluation and Research in Education (IJERE), 2020 Crossref	33 words — 1%
3	repository.unair.ac.id Internet	29 words — 1%
4	worldwidescience.org Internet	27 words — 1%
5	www.hindawi.com Internet	25 words — 1%
6	pt.scribd.com Internet	17 words — 1%
7	www.annfammed.org Internet	16 words — 1%
8	www.hsrd.research.va.gov Internet	15 words — 1%

-
- 9 Nigus Alemnew Engidaw, Abate Dargie Wubetu, Elyas Admasu Basha. "Prevalence of depression and its associated factors among patients with diabetes mellitus at Tirunesh-Beijing general hospital, Addis Ababa, Ethiopia", BMC Public Health, 2020
Crossref 14 words — 1%
-
- 10 eprints.qut.edu.au
Internet 14 words — 1%
-
- 11 www.oalib.com
Internet 13 words — < 1%
-
- 12 Melchor A. Sarmiento, Mariano T Maglutac Jr, Ma. Susan Yanga-Mabunga. "Antibiotic prescribing practices of Filipino Dentists", International Journal of Public Health Science (IJPHS), 2019
Crossref 11 words — < 1%
-
- 13 mafiadoc.com
Internet 11 words — < 1%
-
- 14 aicosh.uin-suka.ac.id
Internet 10 words — < 1%
-
- 15 ijibm.elitehall.com
Internet 10 words — < 1%
-
- 16 e-journal.uajy.ac.id
Internet 9 words — < 1%
-
- 17 Akrom Akrom, Wima Anggitasari. "Adherence and quality of life among diabetic patients with hypertension", International Journal of Public Health Science (IJPHS), 2018
Crossref 8 words — < 1%

-
- 18 doaj.org
Internet 8 words — < 1%
-
- 19 psj.umsha.ac.ir
Internet 8 words — < 1%
-
- 20 www.elsevier.es
Internet 8 words — < 1%
-
- 21 www.tandfonline.com
Internet 8 words — < 1%
-
- 22 Fahmi Anwar, Abdul Fadlil, Imam Riadi. "Validation Analysis of Scalable Vector Graphics (SVG) File Upload using Magic Number and Document Object Model (DOM)", International Journal of Advanced Computer Science and Applications, 2020
Crossref 7 words — < 1%
-
- 23 Dohoon Kim, Hyejung Chang. "Key functional characteristics in designing and operating health information websites for user satisfaction: An application of the extended technology acceptance model", International Journal of Medical Informatics, 2007
Crossref 6 words — < 1%
-

EXCLUDE QUOTES OFF

EXCLUDE MATCHES OFF

EXCLUDE BIBLIOGRAPHY ON