

artikel

by Laela Hayu Nurani

Submission date: 19-Jun-2022 02:29PM (UTC+0700)

Submission ID: 1859305374

File name: 11_fr-2021-331_nurani.pdf (415.08K)

Word count: 5783

Character count: 31336

Isolation of active compound from *Nephelium lappaceum* L. rind as an antioxidant¹Nurani, L.H., ¹Edityaningrum, C.A., ^{2,3}Suhaera, ⁴Windarsih, A., ²Riyanto, S. and ^{2,5,*}Rohman, A.¹Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta 55164, Indonesia²Department of Pharmaceutical Chemistry, Faculty of Pharmacy, Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia³Study Program of Pharmacy, Health Institute of Mitra Bunda, Batam, 29444, Riau Island, Indonesia⁴Research Center for Food Technology and Processing (PRTTP), National Research and Innovation Agency (BRIN), Yogyakarta, 55861, Indonesia⁵Institute of Halal Industry and Systems (IHIS), Universitas Gadjah Mada, Yogyakarta, 55281, Indonesia**Article history:**

Received: 16 May 2021

Received in revised form: 25 June 2021

Accepted: 27 September 2021

Available Online: 11 May 2022

Keywords:Rambutan rind,
In vitro antioxidant,
Phenolic content,
Total Flavonoid,
1,2-benzenedicarboxylic acid**DOI:**[https://doi.org/10.26656/fr.2017.6\(3\).331](https://doi.org/10.26656/fr.2017.6(3).331)**Abstract**

The consumption of rambutan (*Nephelium lappaceum* L.) results in a vast amount of rambutan rind. It is very interesting to explore rambutan rind as natural antioxidants. This study aimed to explore the antioxidant activities of rambutan rind *in vitro* and to correlate those activities with a group of compounds (phenolics and flavonoids). Rambutan rind was cleaned, dried using a conventional oven, powdered, and the powder obtained was subjected to maceration. The macerate was fractionated using petroleum ether and chloroform to get chloroform fraction. The sub fractionation was done toward chloroform fraction. The chloroform fraction and its subfractions were then subjected to antioxidant activity measurement, determination of phenolics and flavonoid contents, and the active compound present in the most active subfractions was isolated using column chromatography. The isolate was identified using several methods including thin-layer chromatography (TLC), Fourier transform infrared (FTIR) spectroscopy as well as gas chromatography-mass spectrometry (GC-MS). The results showed that the chloroform fraction and its subfractions were correlated with the phenolic and flavonoid contents found in rambutan rinds. The phenolic and flavonoid content of chloroform fraction was 32.32 ± 0.79 and 55.05 ± 3.900 , respectively, whereas the IC_{50} of antioxidant activity was 49.77 ± 0.22 for DPPH and 45.36 ± 0.27 for ABTS. The reducing power was 34.478 ± 0.245 (mg vitamin C equivalent/g sample) and the metal chelating activity was 332.753 ± 0.695 (mg Na EDTA equivalent/g sample). The phenolic contents contributed towards the metal chelating activities compared to other antioxidant activities with an R^2 value of 0.8726, while flavonoid contents contributed more to ABTS radical scavenging activity with an R^2 value of 0.8916. One of the active compounds present in rambutan rind having antioxidant activity was identified as 1,2-benzenedicarboxylic acid. Thus, the under-utilized part of rambutan rind could be explored as a natural antioxidant to be used as a food supplement.

1. Introduction

The imbalance between reactive oxygen species (ROS) and reactive nitrogen species (RNS) with endogenous antioxidants has led to the formation of oxidative stress. ROS and RNS have been associated with some degenerative diseases including diabetes, cancer, ageing, neurodegenerative and cardiovascular diseases (White *et al.*, 2014). Some efforts have been driven by a group of researchers to explore exogenous antioxidants either natural or synthetic antioxidants (Palanisamy *et al.*, 2008). The antioxidant can be defined

as any compounds or materials capable of delaying or inhibiting the oxidation reactions of lipids, proteins and other molecules by terminating free radical reaction (Pavithra and Banu, 2017). Natural antioxidants derived from fruits and vegetables have been explored and commercialized such as grape seed extracts. One of the potential natural antiradical derivatives is from Rambutan (*Nephelium lappaceum* L.).

Rambutan (king of fruits) with the scientific name of *Nephelium lappaceum* L. (Family of Sapindaceae) is a tropical fruit commonly found in South East Asian

*Corresponding author.

Email: abdulkimfar@gmail.com

regions like Indonesia, Thailand Malaysia, and Vietnam (Fidrianny *et al.*, 2015). Rambutan is most commonly consumed in cultivars Aceh, Binjai and Rapih (Rohman *et al.*, 2016). The consumption of Rambutan fruit has resulted in a vast amount of waste, including rind and peel. Therefore, it is very interesting to explore of underutilized part of rambutan fruit as a natural antioxidant and identify the active components contributing to antioxidant activities.

Several biological activities of *Nephelium lappaceum* L. fruit including its rind and seed have been reported, namely antiradical due to its phenolic compounds (Thitilertdech and Rakariyatham, 2011), *in vitro* and *in vivo* antioxidant activities using different standardized methods (Pavithra and Banu, 2017; Iman Kamaludin *et al.*, 2016; Mistriyani *et al.*, 2018), lipid peroxidation inhibition (Setyawati *et al.*, 2015) anti-inflammatory effects (Chingsuwanrote *et al.*, 2016) anti-hypercholesterolemia activities (Muhtadi *et al.*, 2016) antibacterial activities against some pathogenic bacterial strains (Thitilertdech *et al.*, 2008), inhibitors of alpha-amylase and alpha-glucosidase activities *in vitro* (Thinkratok *et al.*, 2014) hypoglycemic effects (Soeng *et al.*, 2015), anti-hyperglycemic activity (Palanisamy *et al.*, 2011), α - and β -glucosidases inhibition (Widowati *et al.*, 2015), and anti-diabetic activity on rats induced by alloxan (Muhtadi *et al.*, 2016)

Some active compounds have been identified in rambutan including geraniin, corilagin and ellagic acid (Hernández *et al.*, 2017). A total of 39 compounds were also identified in rambutan rind, including 1 simple phenolic acid, 1 flavone, 5 hydrolyzable tannins, 5 hydroxybenzoic acids, 10 flavonols, 11 flavonols, 6 ellagic acid and conjugates (Zhuang *et al.*, 2017). However, the study on the antioxidant activities of *Nephelium lappaceum* L. rind fraction is still limited. To the best of our knowledge, the isolation of active antioxidant components from *Nephelium lappaceum* L. rind chloroform fraction has not been reported. Mistriyani *et al.* (2021) have carried out the isolation of antioxidant components in the ethyl acetate fraction of Rambutan peel. In this study, the chloroform fraction of Rambutan rind was chosen for further fractionation to investigate the active antioxidant compound because the chloroform fraction and subfraction showed good antioxidant activities. Furthermore, the active component contributing to antioxidant activity was identified.

23

2. Materials and methods

2.1 Materials

Rambutan fruit was obtained from Bantul, Yogyakarta, Indonesia. The authentication of rambutan

samples was performed in the Department of Pharmaceutical Biology, Faculty of Pharmacy with assistance from Dr Djoko Santosa. The chemical reagents and solvents used were of pro-analytical grade.

2.2 Preparation of chloroform extract of rambutan

The chloroform fraction was prepared according to Rohman *et al.* (2016). Rambutan rind was cleaned, cut into small pieces using a commercial cutter and dried in a conventional oven at 65°C for 2 days then grounded into a powder. The powder was subjected to extraction using the maceration technique using methanol as extracting solvent and occasionally shaken. The solvent was evaporated using a vacuum rotary evaporator to obtain the methanolic extract. The extract was then added with aquadest and subjected to partition using petroleum ether, chloroform, and ethyl acetate to get corresponding fractions, namely petroleum ether (PE), chloroform (CH) and ethyl acetate (EA) fractions. CH fraction was then further sub-fractionated to get sub-fractions. The most active sub-fraction as an antioxidant was subjected to isolation to identify the compounds responsible for antioxidant activity. The methanol extracts and their fractions (water, PE, CL, and EA) were evaluated by determining antiradical activities using 2,2'-diphenyl-1-picrylhydrazyl (DPPH) and 2,2'-azinobis (3-ethylbenzo thiazoline-6-sulphonic acid) diammonium salt (ABTS), reducing power, and metal chelating activity.

2.3 Evaluation of radical scavenging activity using DPPH radical

Scavenging activity of DPPH radicals was performed according to (Khalil *et al.*, 2018) with slight modification. A-50 μ L of extract and fraction samples with different concentrations was added to 1.0 mL DPPH 0.4 mM and 3.950 mL methanol. The mixture was allowed to react at room temperature for 20 mins. Then, the absorbance was measured at 515 nm with a calibrated UV-visible spectrophotometer. All measurements were done in triplicate.

$$\% \text{ Antiradical activity} = \frac{A_0 - A_1}{A_0} \times 100$$

Where A_0 is the absorbance of the control (without samples) and A_1 is the absorbance of samples.

2.4 Radical scavenging activity using ABTS technique

The ABTS radical assay of extract and fraction samples was performed according to Fidrianny *et al.* (2015). The mixture of potassium persulphate 2.45 mM and ABTS 7 mM (1:1) was allowed to stand in the dark for 12–16 hrs at room temperature to produce ABTS radical cation ($ABTS^+$). This $ABTS^+$ solution was

diluted with methanol to obtain absorbance values of 0.600-0.800 at 734 nm. The ABTS⁺ working solution (3 mL) and 30 μ L of blank (methanol), standard or sample were mixed and the absorbance was measured at 734 nm after 6 mins using a spectrophotometer.

2.5 Reducing power evaluation

The evaluation of ferric reducing activity power (FRAP) was carried out according to Mistriyani *et al.* (2018). FRAP reagent was prepared by mixing 300 mmol/L acetate buffer (pH 3.6), 10 mmol/L 2,3,5-triphenyl-1,3,4-triazol-2-azoniacyclopenta-1,4-diene chloride (TPTZ) (in 40 mmol/L HCl), and 20 mmol/L ferric chloride (10:1:1, v:v:v). To the 4.5 mL of reagent, 150 μ L ethanol plant extract was added. The absorbance readings were analysed after 5 mins and performed at a wavelength of 593 nm. The blank consisted of FRAP reagent. The final absorbance of each sample was compared with those obtained from the standard curve made from ferric sulphate (FeSO₄.7H₂O) (200–1000 μ mol/L). Results were expressed in nmol Fe²⁺/mg dried extract.

2.6 Metal chelating activity

Metal chelating activity of extract and fractions was measured as described by (Wong *et al.*, 2014). A certain amount (mL) of samples was added with 0.1 mM FeSO₄ (0.2 mL) and 0.25 mM ferrozine (0.4 mL). After incubating at room temperature for 10 min, the absorbance of the mixture was recorded at 562 nm. The chelating activity was calculated using the following formula:

$$\text{Chelating Activity (\%)} = \frac{A_0 - A_1}{A_0} \times 100$$

Where A₀ is the absorbance of the control reaction (without any samples) and A₁ is the absorbance in the presence of extracts or fractions.

2.7 Determination of total phenolics and flavonoid contents

Total phenolics contents of extracts and fractions were determined according to (Kovarovič *et al.*, 2019), while total flavonoids were determined using the spectroscopic method according to Rohman *et al.* (2010). For phenolics content analysis, Folin Ciocalteu reagent (FCR) was used. Briefly, extracts and fractions in methanol were mixed with 0.4 mL of FCR. The solution was allowed to stand at room temperature for 8-10 mins, added with 4 mL Na₂CO₃ 7% and made to 10.0 mL with bidistilled water. The mixture was allowed to stand for 2 hrs and subsequently measured at 725 nm. The phenolic contents were expressed as mg gallic acid equivalents (GAE) per gram of sample (mg/g). For total flavonoid

analysis, samples in methanol were added with 4 mL of distilled water and 0.3 mL of NaNO₂ 5%. After 5 mins, 0.3 mL AlCl₃ 10% was added to the mixture. At 6 mins, 2 mL NaOH 1 M was added to the mixture. Immediately, the mixture was diluted to a volume of 2.4 mL distilled water and thoroughly mixed and its absorbance was measured at 510 nm versus a blank containing all reagents except samples of extracts or fractions. The total flavonoid content of the extracts and fractions were expressed as mg rutin equivalents (RE) per gram of sample (mg/g).

2.8 Fractionation of chloroform extract

Chloroform fraction has good antioxidant activities, therefore, it was further subjected to fractionation using vacuum liquid using silica gel G 60 GF₂₅₄ as in (Rohman *et al.*, 2010). The chloroform fraction was dissolved in chloroform, then anhydrous sodium sulphate was added that was previously heated for 2 hrs at 110°C, and allowed to stand for one night and filtered using a filter paper. The solvent was concentrated using a vacuum rotary evaporator. Samples were added to a column that has been let stand overnight. The chloroform fraction was eluted using the solvents of chloroform, ethyl acetate, and methanol in various compositions. A volume of 50 mL of eluates each was collected for each and evaporated using a vacuum rotary evaporator.

2.9 Fractionation of active fraction

The sub-fractions were loaded into TLC using the stationary phase of GF254 with suitable eluting solvents. The plate was sprayed with DPPH 0.02% in methanol. Sub-fraction capable of bleaching the DPPH colour was subjected to further fractionation using a gravimetric column. The sub-fraction was loaded into a column using silica gel 230-400 mesh (0.040-0.063 mm) and isocratically eluted with n-hexane: acetone (2:2 v/v) and with petroleum ether: ethyl acetate (1:1 v/v), 5 mL of each elute was then collected.

2.10 Isolate purity identification

The purity of the isolate was identified using two methods, melting point and thin-layer chromatography (TLC). The melting point of the crystal was identified using the Buchi Melting Point B-450. Temperatures were recorded at the time the crystal began to melt and the temperature at which the crystal turns into liquid. The test was repeated by measuring the temperature of $\pm 10^\circ\text{C}$ below the melting point obtained, ramped at $1^\circ\text{C}/\text{min}$. For the purity test using TLC, the isolate was eluted using three eluent systems with a mobile phase with different polarity index, acetone: ethyl acetate: chloroform with a ratio of 4: 3: 3 v/v. The compound was considered pure if it had a single spot on the TLC

system. The spot detection was performed visually using UV₂₅₄ and UV₃₆₆.

2.11 Isolate identification

The chemical structure of the isolate was determined using FTIR spectroscopic method (FTIR 100 PERKIN ELMER) and gas chromatography-mass spectrometry (GCMS-QP2010S SHIMADZU, Japan).

3. Results and discussion

Antioxidant activities of *Nephelium lappaceum* L. rind were evaluated using radical scavenging DPPH and ABTS, ferric reducing activity power (FRAP) and metal chelating activity, which can be differentiated as Hydrogen Atom Transfer (HAT) and Single Electron Transfer (SET) (Prior et al., 2005; Rohman et al., 2020).

3.1 Antiradical scavenging activities

DPPH is a synthetic radical which is stable and soluble in polar organic solvents such as methanol and ethanol. The principle of the DPPH radical scavenging test was based on the reduction of radical DPPH (violet in colour) into non-radical DPPH (colourless). Thus, the antiradical compounds were able to reduce DPPH radical intensity at wavelength 517 nm, with the absorptivity molar decreased from 9660 M⁻¹cm⁻¹ (radical DPPH) into 1640 M⁻¹cm⁻¹ (non-radical DPPH). When antiradical such as phenolics and flavonoids donate their hydrogen radical into DPPH radical, the colour of radical DPPH decreases. The proposed reaction between antiradical compounds (antioxidants) with DPPH radical could be seen in Figure 1. The parameter used for describing DPPH radical activity was IC₅₀, the concentration of the samples necessary to cause 50% scavenging of DPPH radical, calculated from a linear regression equation (Rohman et al., 2016). The higher the antiradical

activities, the lower the IC₅₀ values.

Table 1 compile the IC₅₀ values of methanol extract and its fraction of *Nephelium lappaceum* L. rind using DPPH radical with vitamin C (ascorbic acid) as the positive control as it has the strongest anti-radical activities (the lowest IC₅₀). In addition, ethyl acetate fractions showed antiradical activities among extract and fractions evaluated followed by methanol and chloroform (which are not statistically different or P > 0.05) and petroleum ether. It was in accordance with the study conducted by Mistriyani et al. (2021). The antioxidant activity of the ethyl acetate fraction of *Nephelium lappaceum* L. peel was the lowest compared to other fractions (22.6 µg/mL). Samples with IC₅₀ values of 10-50 µg/mL were considered to have strong antioxidant activity. This can be explained that the presence of compounds capable of donating hydrogen radicals present in the EA (ethyl acetate) fraction was effective to decolorize radical DPPH. The antiradical activity of samples measured by the ABTS method using Trolox (6-hydroxy-2,5,7,8-tetramethylchromane-2-carboxylic acid) as a positive control was compiled in Table 2. Among extracts, fractions and positive control evaluated, Trolox exhibited the highest antiradical activities using ABTS radical followed by methanol extract, petroleum ether, ethyl acetate and chloroform fractions.




Figure 1. Reaction Between Radical 2,2'-Diphenyl-1-Picrylhydrazil (DPPH) with Antioxidants (AH) into Non-Radical DPPH Causing the Discoloration of DPPH (Alam et al., 2013)

Table 1. IC₅₀ values of methanol extract and its fraction of rambutan rind using DPPH radical

Extract or fraction	IC ₅₀ values (µg/mL)			Average±SD (µg/mL)
	Replicate 1	Replicate 2	Replicate 3	
Vitamin C	3.43	3.29	3.26	3.33±0.09
Petroleum ether	49.31	47.35	49.25	48.64±1.12
Methanol	49.37	49.15	48.9	49.14±0.23
Chloroform	49.53	49.82	49.96	49.77±0.22
Ethyl acetate	45.65	45.64	45.87	45.72±0.13

Table 2. IC₅₀ values of methanol extract and its fraction of rambutan rind using ABTS radical

Extract or fraction	IC ₅₀ values (µg/mL)			Average±SD (µg/mL)
	Replicate 1	Replicate 2	Replicate 3	
Trolox	3.43	3.29	3.26	3.33±0.09
Petroleum ether	34.591	34.609	34.515	34.57±0.05
Methanol	27.57	27.17	27.17	27.39±0.17
Chloroform	45.06	45.48	45.55	45.36±0.27
Ethyl acetate	38.18	37.95	37.74	39.93±3.61

3.2 Reducing power activities

The reducing power of Fe(III) into Fe(II), known as ferric reducing activity power (FRAP), differs from radical scavenging activities because there are no free radicals, but involved the reduction of ferric ion (Fe^{3+}) from potassium ferricyanide into Ferro ion (Fe^{2+}). The Ferro ion can be monitored by measuring the intensity of Prussian blue colour at a wavelength of 700 nm, and the higher the absorbance at 700 nm indicated the higher the reduction power. The antioxidant activity based on the reducing power of Fe^{3+} into Fe^{2+} was expressed as mg equivalent of vitamin C in one (1) gram sample. The linear equation describing the relationship between concentration of vitamin C (x-axis) and absorbance of Fe^{2+} due to reduction of Fe^{3+} with vitamin C (y-axis) was: $y = 10.771x + 14.878$ ($R^2 = 0.9891$). Table 3 compiled the reducing power of extract and fractions (calculated as mg equivalent vitamin C/g sample). The methanol extract exhibited the strongest reducing power with a FRAP value of 14.446 ± 0.161 mg equivalent vitamin C/gram sample followed by ethyl acetate fraction, chloroform fraction, and petroleum ether fraction. This indicated that reducing compounds present in methanol extract were active and may be present in a high amount.

3.3 Metal chelating activities

The metal chelating activity of extract and fractions was performed in a slightly acidic medium (pH 6.0). The phenolic compounds can bind to Fe^{2+} and the remaining Fe^{2+} could react with ferrozine to form blue-coloured complexes which can be monitored spectrophotometrically at 562 nm. The absorbance of this complex could be reduced by antioxidants such as phenolic compounds, due to its capability to bind to metal (Fe^{2+}). Therefore, any compounds capable of reducing the complex Fe^{2+} -ferrozine could be considered an antioxidant through the mechanism of metal chelating. As a positive control, ethylene diamine tetraacetic (EDTA) was used, as a consequence, the metal chelating activity of extract and fractions were expressed as mg Na. EDTA/gram sample. Table 3 displays the metal chelating activity of methanol extract and its fractions. The chloroform extract revealed the highest metal activity compared to other fractions and methanol extract. Previous research also reported high metal chelating activity of chloroform fraction of *Nephelium lappaceum* L. peel, however, the highest metal chelating activity was found in the water fraction (Mistriyani et al., 2021).

Table 3. The reducing power of Fe^{3+} into Fe^{2+} by extract and fractions of rambutan rind (calculated as mg equivalent vitamin C/g sample)

Samples	Reducing power (Mean±SD, as mg vitamin C equivalent/g sample)	Metal chelating activity (Mean±SD, as mg Na EDTA equivalent/g sample)
Methanol extract	14.446±0.161	250.463±1.062
Petroleum ether fraction	18.796±0.161	284.075±0.0024
Chloroform fraction	34.478±0.245	332.753±0.695
Ethyl acetate fraction	47.636±0.161	200.692±0.0034

3.4 Phenolics and flavonoid contents

Due to its capability to provide hydrogen radicals, reducing Fe^{3+} and binding metals catalyzing oxidation reactions, phenolic and flavonoids contents were correlated with these antioxidants. Total phenolic contents were determined using Folin-Ciocalteu (F-C) reagent and gallic acid was used as standard, therefore phenolic contents were expressed as mg gallic acid equivalent/gram sample (mg GAE/g). The linear regression describing the relationship between gallic acid (x-axis) and its absorbance after reaction with F-C reagent (y-axis) was expressed as $y = 1.338x - 0.0068$ ($R^2 = 0.998$). In addition, flavonoid contents were determined after being reacted with $NaNO_2$, $AlCl_3$ and NaOH to form a red-coloured complex which can be measured spectrophotometrically at 510 nm. Rutin was used as standard during quantitative analysis of flavonoids, therefore, the flavonoid contents were expressed as mg rutin equivalent/g sample (mg RE/g). The linear regression describing the correlation between rutin (x-axis) and its absorbance (y-axis) was expressed as $y = 0.1438x - 0.0365$ ($R^2 = 0.995$). Table 4 compiled the phenolic and flavonoid contents of methanol extract and its fractions. Methanol extract has the highest phenolic contents accounting for $32.39 \pm 2.37\%$ compared to other methanol fractions, while the highest flavonoid contents were found in ethyl acetate fraction accounting for 78.51 ± 0.579 mg RE/g.

Table 4. The phenolic and flavonoid contents of methanol extract and its fractions of rambutan rind

Samples	Phenolics contents (Mean±SD, mg gallic acid equivalent/g)	Flavonoid contents (Mean±SD, as mg rutin equivalent/g)
Methanol extract	21.36±2.25	96.36±0.894
Petroleum ether fraction	27.11±1.23	93.06±0.579
Chloroform fraction	32.32±0.79	55.05±3.900
Ethyl acetate fraction	32.39±2.37	78.51±0.579

The phenolics and flavonoid contents were then correlated with antiradical activities using DPPH and

Table 5. The correlation between antioxidant activities with phenolics and flavonoid contents of methanol extract and its fraction of rambutan rind

Antioxidant activity tests (y-axis)	Its correlation with phenolics contents (x-axis)		Its correlation with flavonoid contents	
	Equation	R ²	Equation	R ²
DPPH radical scavenging	y = 0.056x + 46.732	0.0269	y = -0.0139x + 49.437	0.0211
ABTS radical scavenging	y = 0.6245x + 5.7764	0.7794	y = -0.3722x + 66.109	0.8916
Ferric reducing activity power	y = 0.3048x + 19.511	0.7813	y = -0.505x + 69.6	0.3899
Metal chelating activity	y = 9.9119x - 13.464	0.8726	y = -0.1712x + 126.46	0.2567

ABTS radicals, ferric reducing activity power, and metal chelating activity. The correlation between antioxidant activities with phenolics and flavonoid contents was expressed as linear regression with certain coefficient determination (R²) values, as compiled in Table 5. The R² value indicated the quantitative contribution of one variable (phenolics and flavonoids) toward antioxidant activities. Based on R² values, phenolic contents contributed toward metal chelating activities compared to other antioxidant activities with an R² value of 0.8726. This value indicated that 87.26% of metal chelating activity came from phenolic contents. In addition, flavonoid contents contributed to ABTS radical scavenging activity with an R² value of 0.8916 which indicated that 89.16% of ABTS radical activity was coming from flavonoid contents.

3.5 Isolation and identification of active compound

The chloroform fraction was then fractionated to sub-fractions to obtain isolates with good antiradical activities. Isolate 1 has been isolated from chloroform fraction and subjected to a purity test using TLC and melting point test. In addition, structure identification was performed using infrared spectroscopy and mass spectrometry. The purity test performed by TLC using three different solvent systems indicated that isolate 1 was TLC pure because only one spot was observed. Melting point analysis showed that isolate 1 had a sharp melting point of 62–64°C. Based on TLC and melting point results, isolate 1 can be considered pure and can be continued to be identified. Identification of isolate 1 using FTIR spectroscopy resulted in IR spectra with several peaks. The peak at 2925 cm⁻¹ corresponded to the stretching vibration of C-H. The peak at 1741 cm⁻¹ originated from the stretching vibration of the carbonyl group of C=O, while the peak at 1458 cm⁻¹ was corresponding to the stretching vibration of C=C (benzene), while peaks at 989cm⁻¹ and 722 cm⁻¹ came from C-H bending vibration. Using mass spectrometry, the molecular ion (M⁺) appeared at m/z of 390 amu with a base peak at m/z 279 amu. The fragment ions appeared at m/z of 279 (M-CHCH₃C₂H₅O(C₂H₅)₂ (CH₂)₃OCH₃), 167, 149(M-CH₂CH₃(CH₂)₂C(O)CHCH₃CH), 132, 113, 93,83,71,57,43. Based on IR and mass spectra, the compound was tentatively identified as 1,2-benzenedicarboxylic acid, bis(2-ethylhexyl) ester (CAS)

bis(2-ethylhexyl) phthalate (Figure 2). This compound can be found in some plant extracts such as *Podophyllum hexandrum* rhizome, the stem of *Hugonia mystax* L., and endophytic fungi and it has been reported to have strong antioxidant activity (Li et al., 2012; Vimalavady and Kadavul, 2013; Govindappa et al., 2014).




Figure 2. The Chemical Structure of Bis(2-Ethylhexyl) Ester (CAS) Bis (2-Ethylhexyl) Phthalate

4. Conclusion

Nephelium lappaceum L. (rambutan) rind showed good antioxidant activities determined using either DPPH or ABTS method. The chloroform fraction demonstrated IC₅₀ of 49.77±0.22 for DPPH and 45.36±0.27 for ABTS. Moreover, it also demonstrated good reducing power (34.478±0.245 as mg vitamin C equivalent/g sample) and metal chelating capacity (332.753±0.695 as mg Na EDTA equivalent/g sample). The content of phenolic and flavonoid compounds was correlated to the antioxidant and metal chelating activities. Identification from the most active fraction found that the compound of 1,2-benzenedicarboxylic acid is the active compound responsible for antioxidant activities. This result could be further explored the potential use of rambutan rind compound as a food supplement of antioxidant activities.

Conflict of interest

The authors declare no conflict of interest.

Acknowledgements

The authors acknowledged to Kemendikbud-ristekdikti, Republik Indonesia. The publication of this article was supported by UAD Professorship Program (with a letter of agreement for the implementation of the Professorship Program Number: R3/3/SP-UAD/II/2022).

References

- Alam, M.N., Bristi, N.J. and Rafiquzzaman, M. (2013). Review on in vivo and in vitro methods evaluation of antioxidant activity. *Saudi Pharmaceutical Journal*, 21(2), 143–152. <https://doi.org/10.1016/j.jsps.2012.05.002>
- Chingsuwanrote, P., Muangnoi, C., Parengam, K. and Tuntipopipat, S. (2016). Antioxidant and anti-inflammatory activities of durian and rambutan pulp extract. *International Food Research Journal*, 23(3), 939–947.
- Fidrianny, I., Fikayuniar, L. and Insanu, M. (2015). Antioxidant activities of various seed extracts from four varieties of rambutan (*Nephelium lappaceum*) using 2,2-diphenyl-1-picrylhydrazyl and 2,2'-azinobis (3-ethyl-benzothiazoline-6-sulfonic acid) assays. *Asian Journal of Pharmaceutical and Clinical Research*, 8(5), 227–231.
- Govindappa, M., Vinay, P.S. and Channabasava, R. (2014). Chemical composition of methanol extract of endophytic fungi, *Alternaria* sp. of *Tebeuia argentea* and their antimicrobial and antioxidant activity. *International Journal of Biological & Pharmaceutical Research*, 5(11), 861-869.
- Hernández, C., Ascacio-Valdés, J., De la Garza, H., Wong-Paz, J., Aguilar, C.N., Martínez-Ávila, G.C. and Aguilera-Carbó, A. (2017). Polyphenolic content, in vitro antioxidant activity and chemical composition of extract from *Nephelium lappaceum* L. (Mexican rambutan) husk. *Asian Pacific Journal of Tropical Medicine*, 10(12), 1201–1205. <https://doi.org/10.1016/j.apjtm.2017.10.030>
- Iman Kamaludin, N.H., Mun, L.S. and Sa'adi, R.A. (2016). Evaluation of antioxidant activity of some tropical fruit peel extracts: Extraction conditions optimization of rambutan peel extract. *ARPJN Journal of Engineering and Applied Sciences*, 11(3), 1623–1631.
- Khalil, A.A., Khan, M.R., Shabbir, M.A. and Rahman, K.U. (2018). In vitro antioxidant activity and punicalagin content quantification of pomegranate peel obtained as agro-waste after juice extraction. *Pakistan Journal of Agricultural Sciences*, 55(1), 197–201. <https://doi.org/10.21162/PAKJAS/18.5663>
- Kovarovič, J., Bystrická, J., Micová, M., Harangozo, L., Miššik, J. and Hegedüsová, A. (2019). The influence of variety on the content of total polyphenols and antioxidant activity in leek (*Allium porrum* L.). *Journal of Microbiology, Biotechnology and Food Sciences*, 8(4), 1072–1075. <https://doi.org/10.15414/jmbfs.2019.8.4.1072-1075>
- Li, M., Zhou, L., Yang, D., Li, T. and Li, W. (2012). Biochemical composition and antioxidant capacity of extracts from *Podophyllum hexandrum* rhizome. *BMC Complementary Alternative Medicine*, 12, 263-267. <https://doi.org/10.1186/1472-6882-12-263>
- Mistriyani, Riyanto, S. and Rohman, A. (2018). Antioxidant activities of Rambutan (*Nephelium lappaceum* L.) peel in vitro. *Food Research*, 2(1), 119-123. [https://doi.org/10.26656/fr.2017.2\(1\).150](https://doi.org/10.26656/fr.2017.2(1).150)
- Mistriyani, M., Riyanto, S., Windarsih, A. and Rohman, A. (2021). Antioxidant Activities and Identification of an Active Compound from Rambutan (*Nephelium lappaceum* L.) Peel. *Indonesian Journal of Chemistry*, 21, 259-267. <https://doi.org/10.22146/ijc.50421>
- Muhtadi, M., Haryoto, H., Sujono, T.A. and Suhendi, A. (2016). Antidiabetic and antihypercholesterolemia activities of rambutan (*Nephelium lappaceum* L.) and durian (*Durio zibethinus* Murr.) fruit peel extracts. *Journal of Applied Pharmaceutical Science*, 6(4), 2231–3354. <https://doi.org/10.7324/JAPS.2016.60427>
- Palanisamy, U., Cheng, H.M., Masilamani, T., Subramaniam, T., Ling, L.T. and Radhakrishnan, A.K. (2008). Rind of the rambutan, *Nephelium lappaceum*, a potential source of natural antioxidants. *Food Chemistry*, 109(1), 54–63. <https://doi.org/10.1016/j.foodchem.2007.12.018>
- Palanisamy, U., Manaharan, T., Teng, L.L., Radhakrishnan, A.K.C., Subramaniam, T. and Masilamani, T. (2011). Rambutan rind in the management of hyperglycemia. *Food Research International*, 44(7), 2278–2282. <https://doi.org/10.1016/j.foodres.2011.01.048>
- Pavithra, S. and Banu, N. (2017). Free Radical Scavenging Activity and Total Antioxidant Capacity of Tin Chlorophyllin from *Morinda citrifolia* L. *Research Journal of Pharmacy and Technology*, 10 (2), 453-455. <https://doi.org/10.5958/0974-360x.2017.00091.9>
- Prior, R.L., Wu, X. and Schaich, K. (2005). Standardized methods for the determination of antioxidant capacity and phenolics in foods and dietary supplements. *Journal of Agricultural and Food Chemistry*, 53(10), 4290–4302. <https://doi.org/10.1021/jf0502698>
- Rohman, A., Widodo, H., Lukitaningsih, E., Windarsih, A., Rafi, M. and Nurrulhidayah, A.F. (2020). Review on in vitro antioxidant activities of curcuma species commonly used as herbal components in Indonesia. *Food Research*, 4(2), 286–293. [https://doi.org/10.26656/fr.2017.4\(2\).163](https://doi.org/10.26656/fr.2017.4(2).163)
- Rohman, A., Riyanto, S. and Nugroho, A.E. (2016).

- Antiradical Activities of Rambutan Peel: Study from Two Cultivars. *Research Journal of Phytochemistry*, 11(1), 42–47. <https://doi.org/10.3923/rjphyto.2017.42.47>
- Rohman, A., Riyanto, S., Yuniarti, N., Saputra, W.R., Utami, R. and Mulatsih, W. (2010). Antioxidant activity, total phenolic, and total flavonoid of extracts and fractions of red fruit (*Pandanus conoideus* Lam). *International Food Research Journal*, 17(1), 97–106.
- Setyawati, A., Dewi, A.K., Atho'illah, M.F., Lestari, U. and Lestari, S.R. (2015). The effect of rambutan (*Nephelium lappaceum* L.) peel extract on lipid peroxidation in liver of obese rats. *KnE Life Sciences*, 2(1), 326. <https://doi.org/10.18520/cls.v2i1.167>
- Soeng, S., Evacuasiany, E., Widowati, W. and Fauziah, N. (2015). Antioxidant and hypoglycemic activities of extract and fractions of rambutan seeds (*Nephelium lappaceum* L.). *Biomedical Engineering*, 1(1), 13–18.
- Thinkratok, A., Supkamonseni, N. and Srisawat, R. (2014). Inhibitory potential of the rambutan rind extract and tannin against alpha-amylase and alpha-glucosidase activities *in vitro*, presented at the International Conference on Food, Biological and Medical Sciences (FBMS-2014), January 28-29, p. 44–48. Bangkok, Thailand. <https://doi.org/10.15242/iicbe.c0114582>
- Thitilertdecha, N. and Rakariyatham, N. (2011). Phenolic content and free radical scavenging activities in rambutan during fruit maturation. *Scientia Horticulturae*, 129(2), 247–252. <https://doi.org/10.1016/j.scienta.2011.03.041>
- Thitilertdecha, N., Teerawutgulrag, A. and Rakariyatham, N. (2008). Antioxidant and antibacterial activities of *Nephelium lappaceum* L. extracts. *LWT - Food Science and Technology*, 41(10), 2029–2035. <https://doi.org/10.1016/j.lwt.2008.01.017>
- Vimalavady, A. and Kadavul, K. (2013). Phytocomponents identified on the various extracts of stem of *Hugonia mystax* L. (Linaceae). *European Journal of Experimental Biology*, 3(1), 73-80.
- White, P.A.S., Oliveira, R.C.M., Oliveira, A.P., Serafini, M.R., Araújo, A.A.S., Gelain, D.P. and Santos, M.R.V. (2014). Antioxidant activity and mechanisms of action of natural compounds isolated from lichens: A systematic review. *Molecules*, 19(9), 14496–14527. <https://doi.org/10.3390/molecules190914496>
- Widowati, W., Maesaroh, M., Fauziah, N., Erawijantari, P.P. and Sandra, F. (2015). Free Radical scavenging and alpha/beta-glucosidases inhibitory activities of rambutan (*Nephelium lappaceum* L.) peel extract. *The Indonesian Biomedical Journal*, 7, 157-162. <https://doi.org/10.18585/inabj.v7i3.180>
- Wong, F.C., Yong, A.L., Ting, E.P.S., Khoo, S.C., Ong, H.C. and Chai, T.T. (2014). Antioxidant, metal chelating, anti-glucosidase activities and phytochemical analysis of selected tropical medicinal plants. *Iranian Journal of Pharmaceutical Research*, 13(4), 1407–1413. <https://doi.org/10.22037/ijpr.2014.1588>
- Zhuang, Y., Ma, Q., Guo, Y. and Sun, L. (2017). Protective effects of rambutan (*Nephelium lappaceum*) peel phenolics on H₂O₂-induced oxidative damages in HepG2 cells and D-galactose-induced aging mice. *Food and Chemical Toxicology*, 108(Part B), 554–562. <https://doi.org/10.1016/j.fct.2017.01.022>

ORIGINALITY REPORT

19%

SIMILARITY INDEX

13%

INTERNET SOURCES

16%

PUBLICATIONS

%

STUDENT PAPERS

PRIMARY SOURCES

- 1 Mahfuz Elmastaş, Ibrahim Dermirtas, Omer Isildak, Hassan Y. Aboul - Enein. " Antioxidant Activity of S - Carvone Isolated from Spearmint () ", Journal of Liquid Chromatography & Related Technologies, 2007
Publication 1%
- 2 arrow.tudublin.ie
Internet Source 1%
- 3 Daljit Singh Arora, Priyanka Chandra. "In Vitro Antioxidant Potential of Some Soil Fungi: Screening of Functional Compounds and their Purification from Penicillium citrinum", Applied Biochemistry and Biotechnology, 2011
Publication 1%
- 4 innovareacademics.in
Internet Source 1%
- 5 openaccesspub.org
Internet Source 1%
- 6 psasir.upm.edu.my
Internet Source 1%

7	repository.um.edu.my Internet Source	1 %
8	iicbe.org Internet Source	<1 %
9	scholar.sun.ac.za Internet Source	<1 %
10	Lei Zhao, Huaqing Yang, Meili Xu, Xin Wang, Chengtao Wang, Yunhe Lian, Arshad Mehmood, Haochen Dai. "Stevia residue extract ameliorates oxidative stress in d-galactose-induced aging mice via Akt/Nrf2/HO-1 pathway", <i>Journal of Functional Foods</i> , 2019 Publication	<1 %
11	journal.umpalangkaraya.ac.id Internet Source	<1 %
12	www.biotech-asia.org Internet Source	<1 %
13	Sugama, T.. "Poly(itaconic acid)-modified chitosan coatings for mitigating corrosion of aluminum substrates", <i>Progress in Organic Coatings</i> , 200005 Publication	<1 %
14	Laela Hayu Nurani, Florentinus Dika Octa Riswanto, Anjar Windarsih, Citra Ariani Edityaningrum, Any Guntarti, Abdul Rohman.	<1 %

"Use of chromatographic-based techniques and chemometrics for halal authentication of food products: A review", International Journal of Food Properties, 2022

Publication

15

farfar.pharmacy.bg.ac.rs

Internet Source

<1 %

16

ouci.dntb.gov.ua

Internet Source

<1 %

17

"Fruit Oils: Chemistry and Functionality", Springer Science and Business Media LLC, 2019

Publication

<1 %

18

impactfactor.org

Internet Source

<1 %

19

jbcs.s bq.org.br

Internet Source

<1 %

20

moam.info

Internet Source

<1 %

21

scholars.cityu.edu.hk

Internet Source

<1 %

22

www.iss.it

Internet Source

<1 %

23

Nont Thitilertdecha, Phanuphong Chaiwut, Nisakorn Saewan. "In vitro antioxidant potential of Nephelium lappaceum L. rind

<1 %

extracts and geraniin on human epidermal keratinocytes", Biocatalysis and Agricultural Biotechnology, 2020

Publication

24

MOHAMMAD ARFAN. "ANTIOXIDANT ACTIVITY OF EXTRACTS OF MALLOWUS PHILIPPINENSIS FRUIT AND BARK", Journal of Food Lipids, 9/2007

Publication

25

[dokumen.pub](#)

Internet Source

26

[repositorio.uam.es](#)

Internet Source

27

"Antioxidant activity, total phenolic and flavonoid contents of selected medicinal plants used for liver diseases and its classification with chemometrics", Journal of Applied Pharmaceutical Science, 2019

Publication

28

Abdul Rohman. "Physico-chemical Properties and Biological Activities of Rambutan (Nephelium lappaceum L.) Fruit", Research Journal of Phytochemistry, 2017

Publication

29

Kátia Pereira Santos, Martha Dalila Sedano-Partida, Lucimar Barbosa Motta, Inês Cordeiro, Claudia M. Furlan. "Antioxidant

<1 %

<1 %

<1 %

<1 %

<1 %

<1 %

activity of flavonoids from *Croton sphaerogynus* Baill.", *Brazilian Journal of Botany*, 2016

Publication

30

Maliheh Safavi, Mahroo Seyed Jafari Olia, Mohammad Haji Abolhasani, Mohsen Amini, Mehran Kianirad. "Optimization of the culture medium and characterization of antioxidant compounds of a marine isolated microalga as a promising source in aquaculture feed", *Biocatalysis and Agricultural Biotechnology*, 2021

Publication

31

Yongliang Zhuang, Qingyu Ma, Yan Guo, Liping Sun. "Protective effects of rambutan (*Nephelium lappaceum*) peel phenolics on H₂O₂-induced oxidative damages in HepG2 cells and d-galactose-induced aging mice", *Food and Chemical Toxicology*, 2017

Publication

32

journals.ashs.org

Internet Source

33

wjtcn.net

Internet Source

34

A. Rohman, Dwi Larasati Setyaningrum, Sugeng Riyanto. "FTIR Spectroscopy Combined with Partial Least Square for Analysis of Red Fruit Oil in Ternary Mixture

<1 %

<1 %

<1 %

<1 %

<1 %

System", International Journal of Spectroscopy, 2014

Publication

35

Dergipark.Org.Tr

Internet Source

<1 %

36

Muthukumar Abinaya, Baskaralingam Vaseeharan, Mani Divya, Sekar Vijayakumar et al. "Structural characterization of Bacillus licheniformis Dahb1 exopolysaccharide—antimicrobial potential and larvicidal activity on malaria and Zika virus mosquito vectors", Environmental Science and Pollution Research, 2018

Publication

<1 %

37

S. Badami, K.P. Channabasavaraj. " . Antioxidant Activity of Thirteen Medicinal Plants of India's Western Ghats ", Pharmaceutical Biology, 2008

Publication

<1 %

38

Safia Gacemi, Khedidja Benarous, Santiago Imperial, Mohamed Yousfi. "Lepidine B & E as New Target Inhibitors from Lepidium Sativum Seeds Against Four Enzymes of the Pathogen Candida albicans: In Vitro and In Silico Studies", Endocrine, Metabolic & Immune Disorders - Drug Targets, 2020

Publication

<1 %

39

Internet Source

<1 %

40

clinphytoscience.springeropen.com

Internet Source

<1 %

41

japsonline.com

Internet Source

<1 %

42

Castagnoli, Neal, Kay Castagnoli, Géraldine Magnin, Simon Kuttub, and Jackie Shang. "STUDIES ON THE OXIDATION OF 1,4-DISUBSTITUTED-1,2,3,6-TETRAHYDROPYRIDINES", *Drug Metabolism Reviews*, 2002.

Publication

<1 %

43

D Pandiangan, P Y Lamlean, P F Maningkas, N Nainggolan, A J A Unitly. " Antioxidant and Anticancer Activity Tests of "Pasote" Leaf Water Extracts (L.) by In Vitro Method in Leukemia Cancer Cells ", *Journal of Physics: Conference Series*, 2020

Publication

<1 %

44

Hefa Mangzira Kemung, Loh Teng-Hern Tan, Kok-Gan Chan, Hooi-Leng Ser, Jodi Woan-Fei Law, Learn-Han Lee, Bey-Hing Goh. " Antioxidant Activities of sp. strain MUSC 14 from Mangrove Forest Soil in Malaysia ", *BioMed Research International*, 2020

Publication

<1 %

45 Siti Nur Aqilah Abdul Malek, Hasnah Haron, Wan Aida Wan Mustafa, Suzana Shahar. "Physicochemical Properties, Total Phenolic and Antioxidant Activity of Mixed Tropical Fruit Juice, TP 3 in 1TM", Journal of Agricultural Science, 2017
Publication

46 www.isnff-jfb.com
Internet Source

47 ALBAYRAK, SEVIL, and AHMET AKSOY. "EVALUATION OF ANTIOXIDANT AND ANTIMICROBIAL ACTIVITIES OF TWO ENDEMIC ANTHEMIS SPECIES IN TURKEY : BIOACTIVITIES OF TWO ENDEMIC ANTHEMIS SPECIES", Journal of Food Biochemistry, 2012.
Publication

48 Abdul Rohman, Yaakob B. Che Man, Sugeng Riyanto. "Authentication Analysis of Red Fruit (Pandanus Conoideus Lam) Oil Using FTIR Spectroscopy in Combination with Chemometrics", Phytochemical Analysis, 2011
Publication

49 Chung-Weng Phang, Sri Nurestri Abd Malek, Halijah Ibrahim. "Antioxidant potential, cytotoxic activity and total phenolic content of Alpinia pahangensis rhizomes", BMC Complementary and Alternative Medicine, 2013

50

Jia Ling Tsong, Lucky Poh Wah Goh, Jualang Azlan Gansau, Siew-Eng How. "Review of Nephelium lappaceum and Nephelium ramboutan-ake: A High Potential Supplement", *Molecules*, 2021

Publication

<1 %

51

Khonkarn, R.. "Investigation of fruit peel extracts as sources for compounds with antioxidant and antiproliferative activities against human cell lines", *Food and Chemical Toxicology*, 201008/09

Publication

<1 %

52

T. K. Lim. "Edible Medicinal And Non-Medicinal Plants", Springer Science and Business Media LLC, 2012

Publication

<1 %

53

Tang, Ning, and Hong Zhuang. "Evaluation of Antioxidant Activities of Zein Protein Fractions : Antioxidant peptides obtained from zein...", *Journal of Food Science*, 2014.

Publication

<1 %

54

documents.mx

Internet Source

<1 %

55

dspace.alquds.edu

Internet Source

<1 %

56

ijpsr.com

<1 %

57

pubag.nal.usda.gov

Internet Source

<1 %

58

www.jmbfs.org

Internet Source

<1 %

59

www.sciencepub.net

Internet Source

<1 %

60

Geórgea V. Souza, Alex S. Simas, Amanda L. Bastos-Pereira, Gisele R. A. Frois et al.

"Antinociceptive Activity of the Ethanolic Extract, Fractions, and Aggregatin D Isolated from *Sinningia aggregata* Tubers", PLOS ONE, 2015

Publication

<1 %

61

Nayara Macêdo Peixoto Araujo, Henrique Silvano Arruda, David Roger Paixão Marques, Williara Queiroz de Oliveira et al. "Functional and nutritional properties of selected Amazon fruits: A review", Food Research International, 2021

Publication

<1 %

62

Silva, E.M.. "Antioxidant activities and polyphenolic contents of fifteen selected plant species from the Amazonian region", Food Chemistry, 2007

Publication

<1 %

63

Walid Kriaa, Hamadi Fetoui, Mohamed Makni, Najiba Zeghal, Nour-Eddine Drira. " Phenolic Contents and Antioxidant Activities of Date Palm (L.) Leaves ", International Journal of Food Properties, 2012

Publication

<1 %

64

Zohra Dhouafli, Hichem Ben Jannet, Borhane Mahjoub, Manuela Leri et al. " 1,2,4-trihydroxynaphthalene-2- - β -D-glucopyranoside: A new powerful antioxidant and inhibitor of A β aggregation isolated from the leaves of ", Natural Product Research, 2017

Publication

<1 %

65

Iacovino, Rosa, Filomena Rapuano, Jolanda Caso, Agostino Russo, Margherita Lavorgna, Chiara Russo, Marina Isidori, Luigi Russo, Gaetano Malgieri, and Carla Isernia. " β -Cyclodextrin Inclusion Complex to Improve Physicochemical Properties of Pipemidic Acid: Characterization and Bioactivity Evaluation", International Journal of Molecular Sciences, 2013.

Publication

<1 %

66

Siti Fatonah Fatonah, Sri Catur Setyawatiningsih, Sujarwati Sujarwati, Murniati Murniati et al. "Pemanfaatan Tanaman Pekarangan Untuk Pengobatan

<1 %

Herbal", JPPM (Jurnal Pengabdian dan Pemberdayaan Masyarakat), 2020

Publication

67

Wipawee Chanthasri, Nuntitporn Puangkeaw, Nongluk Kunworarath, Patcharawalai Jaisamut et al. "Antioxidant capacities and total phenolic contents of 20 polyherbal remedies used as tonics by folk healers in Phatthalung and Songkhla provinces, Thailand", BMC Complementary and Alternative Medicine, 2018

Publication

<1 %

68

www.imrfjournals.com

Internet Source

<1 %

Exclude quotes Off

Exclude matches < 3 words

Exclude bibliography On

artikel

GRADEMARK REPORT

FINAL GRADE

/100

GENERAL COMMENTS

Instructor

PAGE 1

PAGE 2

PAGE 3

PAGE 4

PAGE 5

PAGE 6

PAGE 7

PAGE 8
