

Sitti Nur Djannah, Lina Handayani, Rochana Ruliyandari, Intan Wahyuni Tukiyo

by Sitti Nur Djannah, Lina Handayani, Rochana Ruliy, Analysis Of Risky
Food Consumption Behavior Among

Submission date: 29-May-2023 10:01AM (UTC+0700)

Submission ID: 2104174340

File name: 23065-45132-1-RV_VERSION_22_Mei_2023.docx (345.41K)

Word count: 5872

Character count: 33922

Analysis of risky food consumption behavior among adolescents in Yogyakarta

Sitti Nur Djannah¹, Lina Handayani¹, Rochana Ruliyandari¹, Intan Wahyuni Tukiyo¹

¹Faculty of Public Health, Ahmad Dahlan University, Yogyakarta

Article Info

Article history:

Received month dd, yyyy

Revised month dd, yyyy

Accepted month dd, yyyy

Keywords:

Dental and oral diseases,
Non-communicable diseases,
Risky food behavior,
Students

ABSTRACT

Non-communicable diseases (NCD) and oral disease problems can arise in all age groups, including teenagers. This is triggered by risky behavior due to a bad lifestyle, uncontrolled eating patterns and risky foods, lack of physical activity, smoking, and consuming alcohol. This study aimed to determine risky behavior and the incidence of NCD as well as dental and mouth problems in students at the Yogyakarta campus which can trigger NCD and dental and mouth problems. This research is quantitative descriptive-analytic research with a total sample of 165 students at campuses in Yogyakarta. Data was collected through an electronic questionnaire (google form) and the data was analyzed descriptively. The results showed that students' risky behavior sequentially, namely consuming excessive sweet foods (87.27%), salty foods (87.88%), oily foods (82.42%), foods containing seasonings or MSG (80%), excessive consumption of instant noodles (57.58%), lack of fruit and vegetable consumption 145 (84.85%) and lack of physical activity for 30 minutes (87.27 %). All students had no symptoms or doctor's diagnosis of NCD disease, but dental and oral health in the last 12 months had a high incidence which required attention. It was concluded that the most dominant risk behavior for non-communicable disease risk sequentially was poor diet related to food consumption and lack of physical activity. All students had no symptoms or no doctor's diagnosis, but students experienced oral health within 12 months. Maintaining a healthy diet, increasing physical activity, and maintaining good oral hygiene can reduce the risk of NCD and dental and oral diseases.

3

This is an open access article under the [CC BY-SA](https://creativecommons.org/licenses/by-sa/4.0/) license.



Corresponding Author:

Sitti Nur Djannah

Faculty of Public Health, Ahmad Dahlan University, Yogyakarta, Indonesia

Email: sitti.nurdjannah@ikm.uad.ac.id

1. INTRODUCTION

Adolescence is a transitional age between children and adults with a high curiosity about their surroundings. According to UNICEF data for 2021 it shows that 17% of the 270,203,917 total population in Indonesia are teenagers of productive age, as many as 46 million people are teenagers with an age range of 10 to 19 years where the proportion of women as much as 48% and men 52% [1]. According to Piaget's theory, suggests that adolescence is the age when individuals integrate with adult society, the age when children no longer feel below the level of older people but are at the same level, at least in solving problems [2].

The World Health Organization (WHO) in 2018 reported that around 75% of the causes of death in the world are non-communicable diseases with 80% occurring in middle and low-income countries. Most occur due to heart and blood vessel disease by 35%, cancer 12%, chronic respiratory disease 6%, diabetes 6%, and 15% caused by other non-communicable diseases, with risk factors for non-communicable diseases 80% due to lifestyle such as lack of fruit and vegetable consumption, smoking, lack of physical activity, and obesity [3]. The 2018 Riskesdas results show the prevalence of non-communicable diseases has increased from 2013. The prevalence of cancer in 2018 has increased from 1.4% to 1.8%; stroke prevalence increased from 7% to 10.9%; and chronic kidney disease increased from 2% to 3.8%. Based on blood sugar examination, diabetes mellitus

increased from 6.9% to 8.5%; and blood pressure measurement results, hypertension rose from 25.8% to 34.1%. The increase in the prevalence of non-communicable diseases is related to unbalanced lifestyles and eating patterns, including smoking, consumption of alcoholic beverages, physical activity, and insufficient consumption of fruits and vegetables [4], [5].

As much as 60% of mortality in all age groups is a negative impact of the development of non-communicable diseases such as cardiovascular disease, diabetes mellitus, cancer and chronic obstructive pulmonary disease. Risk factors for the incidence of non-communicable diseases 80% are due to a bad lifestyle such as lack of consumption of fruits and vegetables, smoking, lack of physical activity, and obesity [3], [6]. The description of the health status of the adult population in the future is determined by the health status of today's adolescents. Adolescents have the opportunity to obtain optimal health status, but it is not always in line with the health behavior of adolescents. Tobacco use, drug abuse, including alcohol begins at this age. Risky behavior during adolescence greatly influences the incidence of chronic disease in the following decades. As risk factors are smoking behavior, alcohol, diet, physical activity, traffic behavior and emotional distress [7].

Risky behavior, especially diet, physical activity, in the future will have a negative impact on the body such as non-communicable diseases. A good diet is needed to fulfill optimal nutritional needs for adolescents so that they are able to prevent or minimize the risk of various non-communicable diseases (NCDs) that appear in old age such as diabetes mellitus, heart disease and stroke and are able to prevent anemia and stunting. Especially for women, if risky behavior continues into adulthood and marriage it will have an impact on pregnancy, namely babies are susceptible to disease and can cause stunting in children [8], [9]. According to IFRC (2022) NCDs are diseases that are not spread through infection or through other people, but are usually caused by unhealthy behavior. Non-communicable diseases are a major cause of death worldwide and present a major threat to health and development, especially in low- and middle-income countries with high risk of diseases such as cardiovascular disease, cancer, diabetes and chronic respiratory disease [10]. WHO (2014) describes that non-communicable diseases require a long period of time or are commonly referred to as chronic in their development [11]. Non-communicable diseases are a health problem that causes high morbidity, disability and death rates, as well as creates a large health financing burden Data and Information Center [12].

Adolescents are an age group that is very vulnerable to bad health habits/behaviors related to lifestyle, health promotion and the environment. The results of a preliminary study using the interview method with adolescents at a Yogyakarta campus resulted in the preferred risky food, namely junk food, they stated that it was easy to get junk food, the good taste of junk food made them like it. Another behavior is physical activity that is difficult to do consistently. Even though they are teenagers studying in the field of public health who are later expected to become health cadres who are able to provide education to the community in maintaining health and improving health. Based on this description, the purpose of this study is to analyze health behavior that is at risk of developing non-communicable diseases in students of one of the tertiary institutions in Yogyakarta, Indonesia.

2. METHOD

This research is research with quantitative descriptive analytic method to describe the health behavior of students at Yogyakarta university. An overview of student health behavior was taken using a survey on health behaviors that referred to questions in the Indonesian Basic Health Research called RISKESDAS. The population of this study were all students in semester 1 of Yogyakarta university, with accidental sampling technique and a final total sample of 165 respondents. Data collection was carried out using an electronic questionnaire (google form) which was then distributed directly to students consisting of 2 parts, namely the characteristics of the respondents and the core questions totaling 57 questions regarding the health conditions and behavior of students [13]. The questionnaire consisted of questions regarding health behavior and daily habits. Behavior of consuming risky foods, carbonated drinks, smoking and alcohol consumption: 1) risky (≥ 1 time per day and $\geq 1-6$ times a week); 2) not at risk (≤ 3 times per month), fruit and vegetable consumption behavior: 1) at risk (not consuming and/or consuming 1-2 servings); 2) not at risk (3-4 servings and 3 servings of vegetables, 2 servings of fruit). The data obtained was then subjected to quantitative descriptive analysis based on risky and non-risk health behaviors and habits in college students. This study has gone through the UNY Ethics Committee's ethics for review process.

3. RESULTS AND DISCUSSION

3.1. Results

1. Characteristics of Respondents

The following are the characteristics of the respondents, namely students at a university in Yogyakarta shown on Table 1.

Table 1. Distribution of Student Characteristics at Yogyakarta Higher Education

Characteristics	N	%
Gender		
Man	28	16.97
Woman	137	83.03
Age		
17 years	2	1.21
18 years	86	52.12
19 years old	55	33.33
20 years	17	10.30
21 years	5	3.03
Body Mass Index (BMI)		
Normal	29	17.58
Obesity I	4	2.42
Obesity II	3	1.82
Overweight	2	1.21
Underweight	127	76.97
Total	165	100

Based on Table 1, it is known that the majority of students are women (83.03%). The age range was dominated by students aged 18 years (52.12%), then from the calculation of the body mass index most of the students were underweight (76.97%).

2. Student Health Behavior

Student health behavior, including risk eating behavior, smoking behavior, alcohol consumption, and exercise habits as follows can shown on Tabel 2.

Table 2. Health Behavior towards Non-Communicable Diseases in Students at a University in Yogyakarta

Behavioral risk factors	No Risk		Risky	
		%		%
Sweet Eating Behavior	21	12.73	144	87.27
Salty Eating Behavior	20	12.12	145	87.88
Greasy/fried eating behavior	29	17.58	136	82.42
Burning Eating Behavior	86	52.12	79	47.88
Eating Behavior with Seasonings	33	20	132	80
Behavior of Eating Instant Noodles	70	42.42	95	57.58
Fruit and Vegetable Eating Behavior	25	15.15	140	84.85
Drinking behavior of carbonated/carbonated drinks	91	55.15	74	44.85
Smoking history in the last month	148	89.70	17	10.30
Habit of consuming alcohol	164	99.39	1	0.61
Habit of Doing Exercise for 30 Minutes	21	12.73	144	87.27

Based on Table 2, it shows that health behaviors that pose a risk to students are excess food and drink consumption behavior and lack of physical activity. Student eating and drinking behavior is dominated by risky behavior that can trigger NCD. Students have risky behavior in consuming sweet foods (87.27%), as well as consuming salty foods (87.88%) students, consuming oily foods such as fried foods (82.42%), consuming food containing seasonings or MSG (80%), excessive eating behavior of instant noodles (57.58%), lack of fruit and vegetable consumption 140 (84.85%). As well as having

less exercise habits (87.27%). Furthermore, student behavior that is not at risk is a history of not smoking in the last one month (89.70%) and the habit of not consuming alcohol (99.39%).

3. Incidence of Non-communicable Diseases and Dental and Mouth Problems

The incidence of Non-Communicable Diseases in students during the last 1 year is as follows show on Table 3.

Table 3. Incidence of Non-Communicable Diseases and Dental and Oral Problems in Students at the Yogyakarta Campus

Non-Communicable Diseases	Yes	%	No	%
Asthma	3	1.82	162	98.18
Cancer	0	0	165	100
Diabetes mellitus	0	0	165	100
Heart disease	2	1.21	163	98.79
Hypertension	3	1.82	162	98.18
Strokes	0	0	165	100
Chronic Renal Failure	0	0	165	100
Joint Disease	1	0.61	164	99.39
Dental caries	61	36.97	104	63.03
Missing Teeth due to Extraction	25	15.15	140	84.85
Rocking Teeth	20	12.12	145	87.88
Swollen gums and or boils (abscess)	25	15.15	140	84.85
Gums Bleeding Easily	25	15.15	140	84.85
Recurrent thrush	24	14.55	141	85.45

Based on Table 3 the incidence of non-communicable diseases in college students is dominated by students who have no symptoms or no doctor's diagnosis. However, the results showed that students had problems with oral and dental health, including dental caries (36.97%), tooth extraction (15.15%), swollen gums and/or abscesses (15.15%), gums that bleed easily (15.15%), recurrent canker sores (14.55%) and loose teeth (12.12%).

3.2. Discussion

Based on the results of the study, it was shown that the behavior of consuming food and drink, exercise for at least 30 minutes can trigger the risk of non-communicable diseases in adolescents. So it can be said that students have a poor diet and lack of daily physical activity. According to previous studies, poor eating patterns are prone to the emergence of non-communicable diseases such as diabetes mellitus, hypertension, and heart disease. Also supported by less physical activity can increase the risk of developing NCD from an early age [14], [15]. Previous research said that one of the non-communicable diseases that often appears asymptomatic and can appear since adolescence and whose prevalence has increased in recent years is hypertension. Many people do not realize this, that hypertension that occurs in adolescence will continue into adulthood and increase the risk of morbidity and mortality. Excessive consumption of sodium or foods with excessive levels of salt (salty) has a 14.752 greater risk of developing hypertension [16].

Non-Communicable Diseases (NCD) are diseases that cannot be transmitted so they are considered not to threaten the condition of others. Until now NCD is still a health problem in developing countries and industrial countries. Based on the WHO report, NCD which is often found in the South Asia region and contributes to very high incidence and death rates, namely heart (cardiovascular) disease, DM, cancer, chronic obstructive respiratory disease and diseases due to accidents [17]. Risky behavior adopted by teenagers due to unhealthy associations and information that is not directed [18]. Adolescents are a vulnerable group because they are in a period of development and a period of searching for identity and tend to engage in risky behavior. Risk factors in adolescents include increased blood pressure, blood sugar, body mass index or obesity, unhealthy eating patterns, lack of physical activity, and smoking and alcohol. The increase in these risk factors will have an impact on increasing the proportion of non-communicable diseases in adolescents, including diabetes mellitus and hypertension [6], [19].

The results of the same study conducted on Halu Oleo University students showed that apart from smoking and consuming alcohol, risk factors for lack of physical activity (56.3%), and eating patterns based on frequency of eating/drinking trigger non-communicable diseases for frequent consumption of types of food contains sugar (72.1%), contains sodium (salt) (35.0%), contains high fat (55.5%), while the type of drink containing caffeine with habit is a trigger factor for non-communicable diseases, from risk factors This gave

rise to diseases such as obesity (21.8%), and hypertension (13.9%) [20]. Lifestyle behaviors at risk for NCDs with the greatest proportion in adolescents are insufficient consumption of fruits and vegetables, excessive consumption of sodium, excessive consumption of fat, lack of physical activity and smoking [21]–[23]. Consuming excessive risky foods such as excessive junk food, consumption of sweet foods, processed foods with preservatives has a close relationship with the incidence of non-communicable diseases will trigger the emergence of nutritional disorders at a young age which can increase the risk of non-communicable diseases [24]–[26].

Poor eating patterns such as junk food can affect adolescent nutrition including overweight and obesity [12], because the high content of junk food will be salt, fat, sugar, and calories, but little nutrition and fiber can increase the risk of several diseases, such as obesity, diabetes, hypertension, and blood lipid disorders or dyslipidemia [26], [27]. So maintaining a good diet and increasing the intensity of physical activity, especially when the age has entered > 40 years and a healthy lifestyle is very important. This is due to the trend of increasing prevalence of non-communicable diseases caused by the epidemiological transition, which begins with the dominance of communicable diseases and ends with the dominance of non-communicable diseases [28].

In addition to poor food consumption patterns, risk factors for lack of physical activity also greatly affect the incidence of NCDs in adolescence. One of the negative impacts of lack of physical activity is the emergence of diabetes mellitus and obesity or overweight [23]. According to the Law of the Republic of Indonesia Number 36 of 2009 concerning Health article 141 states that it is necessary to improve the quality of nutrition through food consumption patterns and physical activity in accordance with the General Guidelines for Balanced Nutrition (PUGS) in order to achieve good nutritional status so that the degree of public health increases. Health is a human right. Research conducted on high school students in Semarang City showed the same results, namely students who did light physical activity as a risk factor for developing hypertension at high school age with a risk of 10.074 times. Mild physical activity in adolescents occurs due to increased insulin levels so that the body will become hungry quickly [16]. Apart from hypertension and obesity, lack of physical activity will cause other health problems such as stress, sleep disturbances, slowed metabolism, and osteoporosis [29], [30].

Increasing the intensity of physical activity, namely 30–45 minutes/day, is very important because it is one of the strategies for managing and preventing hypertension. Regular physical activity can reduce stiffness in the blood vessels and will increase the endurance of the heart and lungs so that it can lower blood pressure [15]. Doing strenuous physical activity or moderate physical activity that cannot control their eating patterns after doing physical activity, so that the incidence of diabetes mellitus does not affect if you have done heavy or moderate physical activity [30]. Physical activity significantly reduces systolic and diastolic blood pressure and can prevent hypertension and other non-communicable diseases. The recommended length of physical activity to prevent hypertension is 130 minutes, with a frequency of five or more days a week [31].

Lifestyles that are at risk increase the risk of non-communicable diseases in adolescents. One of them is by carrying out CERDIK Activities which consist of health check activities accompanied by providing knowledge about diabetes mellitus, non-smoking behavior, physical activity, reducing consumption of Sugar, Salt, Fat, adequate rest, and managing stress. In line with research on adolescents in Ambon who showed good attitudes and knowledge in realizing CERDIK activities, namely frequent physical activity (sports) [32], [33]. Based on the priority indicators of the Non-Communicable Disease Prevention And Control Program in the form of the implementation of an integrated Non-Communicable Disease Fostered Post Program (Posbindu NCD) in each village to carry out early detection of risk factors for non-communicable diseases, control of smoking consumption, and faulty health services for productive age one target is every citizen aged 15 years and over in a village or sub-district, in this case, are youth and adults [34][6], [34]. Primary prevention efforts for NCD can be started from childhood and adolescence to learn to behave in the right lifestyle. The number of adolescents, 24.01 percent of Indonesia's population, can become agents of change in good lifestyle behavior in preventing the risk of non-communicable diseases (NCD) in adulthood. Knowledge and proper and adequate health-nutrition knowledge and behavior from schools, residences and the surrounding environment such as health education about risky foods and the impact on health through banner media is useful in efforts to control degenerative diseases and prevent NCDs. Furthermore, when they enter the productive group, Indonesia has healthy human resources with high productivity [21], [35].

The results showed that diseases related to dental and oral problems occurred in adolescents, especially students, namely dental caries (36.97%), teeth extracted (15.15%), swollen gums and/or abscesses (15.15%), gums bleed easily (15.15%), recurrent canker sores (14.55%) and loose teeth (12.12%). Dental health efforts need to be reviewed from environmental aspects, knowledge, education, awareness and handling of dental health including prevention and treatment. This is in accordance with research in Ternate, most students from the city of Ternate in the city of Manado ignore the overall condition of dental health. Dental care is considered not very important, even though it has enormous benefits in supporting health and appearance. Apart from that, fear treatment that takes a long time and costs are quite high which allows the

Paper's should be the fewest possible that accurately describe ... (First Author)

risk of caries to students from the city of Ternate. Maintaining oral and dental hygiene is very important, some oral and dental problems occur because we don't maintain dental and oral hygiene. Awareness of maintaining oral hygiene is very necessary and is a medicine to prevent caries [36].

The 2013 Indonesian Basic Health Research obtained a DMF-T score in Indonesia of 4.6 and a DMF-T in South Sumatra of 5.3. Meanwhile, the results of the 2018 Riskesdas found that the score for damaged/cavity teeth in Indonesia was 45.3%, the score for missing teeth being removed/dated itself was 19.0%. And the score for teeth that have been filled because of cavities in Indonesia is 4.1%. Based on Indonesian Basic Health Research (2018) the proportion of habitual consumption of sweets ≥ 1 time per day in Indonesia is around 40.1%. The proportion of consumption of sweet drinks ≥ 1 time per day in Indonesia is around 61.27%. Incorrect eating patterns and some community behaviors as well as respondents who prefer sweet, less fibrous and easily sticky foods and drinks which cause dental caries, as well as the behavior of frequently consuming sweet-tasting snacks such as chocolate, milk, ice cream, candy, sweet cakes, and sweet chips, more than 2-3 times a day. This type of food is highly cariogenic and has the potential to cause dental caries [37]. One of the risk factors for the emergence of non-communicable diseases is smoking, where smoking habits can also cause failure to benefit from oral health care, and more dynamic tooth decay, which causes pain and even tooth loss [38]. In line with previous research which said that non-communicable diseases also have a relationship related to oral health, found a negative relationship between oral health knowledge and poor Self-Rated Oral Health (SROH), this study found a positive relationship, namely the possibility of an epidemiological transition from communicable diseases to non-communicable diseases [39]. In addition, overall, 13.6% of participants reported poor Self-Rated Oral Health, and 78.5% on average or poor SROH, oral conditions (teeth loss, cavities, bleeding gums, and teeth sensitive to heat or cold), knowledge better oral health, dental care, and skipping breakfast were associated with poor Self-Rated Oral Health [40].

Oral diseases have been associated with major conditions like five types of cancer, diabetes mellitus, cardiovascular diseases, depression, neurodegenerative conditions, rheumatic diseases, inflammatory bowel disease, gastric helicobacter pylori, obesity, and asthma and there are several risk factors that oral diseases share with these conditions (such as sugar consumption, smoking), which suggests that modeling the health and economic benefits to promote oral health can be useful [41], [42]. Additionally, in another study, school-based interventions are found to be particularly effective at reducing the burden of oral diseases among primary school children including skills-based education, teacher training, access to oral health services, and parental involvement. Primary school-based interventions to improve oral health need further research to be proven effective [43]. From a study conducted in Saudi Arabia on dental students showed that the existence of public dental health and preventive dentistry courses had a significant and positive impact on the knowledge, attitudes and behavior of undergraduate students. After an analysis using the HU-DBI score shows that male students do not show a significant relationship and have a low HU-DBI score on oral health behavior, this is related to the large number of students who smoke tobacco, drink alcohol, and use bad internet. Additionally, dental students' average HU-DBI scores are weakly correlated with the economic rankings of the countries where they live/study [44].

According to research in the Longhua district of Shenzhen, 27.6% of participants had never been to a dentist before, along with knowledge and education. Moderate and severe caries risk factors include Angle Class II malocclusion, brushing sometimes or never, flossing, having experienced a toothache within the past 12 months, a high level of knowledge attitude, a moderate level of knowledge attitude, and a high level of knowledge attitude. so it is necessary to increase dental and oral health education as well as good dental and oral hygiene habits [45]. Oral disease is caused by a variety of modifiable risk factors common to many non-communicable diseases (NCDs), including sugar consumption, tobacco use, alcohol use and poor hygiene, as well as underlying social and commercial factors [46]. Global Oral Health Status Report (2022) reported that non-communicable diseases such as mouth disease, heart disease, cancer, diabetes, chronic respiratory disease and mental illness are generally caused by risk factors. All forms of tobacco use, harmful use of alcohol, unhealthy diet, physical activity, environmental pollution [47]. Most oral diseases and conditions share modifiable risk factors such as tobacco use, alcohol consumption, and an unhealthy diet high in free sugars which are common in the 4 major NCDs (cardiovascular disease, cancer, chronic respiratory disease, and diabetes). In addition, diabetes has been reciprocally associated with the development and progression of periodontal disease. There is also a causal relationship between high consumption of sugar and diabetes, obesity and dental caries [46], [48].

4. CONCLUSION

Risky health behaviors in college students that can lead to non-communicable diseases and dental and mouth problems, respectively, are excessive consumption of salty foods (87.65%), lack of physical activity (87.65%), consumption of sweet foods (87.06%), lack of consumption of fruits and vegetables (85.29%), oily foods (81.76%), foods containing seasonings or MSG (80.59%), and excessive consumption of instant noodles

(57.65%). NCD's events, such as hypertension, diabetes mellitus, heart disease, chronic kidney failure where all students have no symptoms or no doctor's diagnosis results. However, students have problems with oral and dental health, including dental caries (36.97%), teeth extracted (15.15%), swollen gums and/or abscesses (15.15%), gums bleed easily (15.15%), recurrent canker sores (14.55%) and loose teeth (12.12%) in the last 12 months. Based on this research, we find that poor diet related to food consumption and lack of physical activity is the most important risk behavior according to NCD in students at Yogyakarta University. Furthermore, oral disease problems show no symptoms or diagnosis from doctors, but they had experienced oral health within 12 months. Therefore, from this study, we suggest that all students have to maintain with a healthy diet, and increase physical activity also oral hygiene, this behavior can reduce the risk of NCD and oral diseases.

ACKNOWLEDGEMENTS

Author thanks all parties and respondents who support this research.

REFERENCES

- [1] UNICEF (United Nations Children's Fund), "2021 Youth Profile," *unicef*, pp. 1–9, 2021, [Online]. Available: https://www.unicef.org/indonesia/media/9546/file/Youth_Profile.pdf.
- [2] EB Hurlock, *Developmental Psychology An Approach Throughout the Life Span*, Print 5. Jakarta: Erlangga Publisher, 2002.
- [3] WHO, "Noncommunicable Disease Country Profile 2018," Geneva, 2018.
- [4] Riskesdas, "2018 Riskesdas Report," *Health Research and Development Agency*, p.s. 198, 2018, [Online]. Available: http://labdata.litbang.kemkes.go.id/images/download/report/RKD/2018/Laporan_Nasional_RKD2018_FINAL.pdf.
- [5] Ministry of Health, "RISKESDAS 2013," 2013. doi: 10.1126/science.127.3309.1275.
- [6] Ministry of Health and Ministry of Health of the Republic of Indonesia, "Manual of non-communicable disease management," p. 2, 2019.
- [7] S. Isfandari and DB Lolong, "Analysis of risk factors and health status of Indonesian youth in the next decade Risk factors and health status of Indonesia young adults: Indonesia picture of the next decade," *Bul. Researcher. Health.*, vol. 42, no. 2, pp. 122–130, 2014.
- [8] P2PTM RI, "Unbalanced Nutrition When Adolescents Are at Risk of Increasing Non-Communicable Diseases," <https://p2ptm.kemkes.go.id/kotalgizi-tak-sebalance-saat-remaja-berisiko-level-disease-ntak-menular>, 2022. .
- [9] N. Akseret *et al.*, "Non-communicable diseases among adolescents: current status, determinants, interventions and policies," *BMC Public Health*, vol. 20, no. 1, pp. 1–21, 2020, doi: 10.1186/s12889-020-09988-5.
- [10] IFRC, "Non-Communicable Diseases," [https://www.ifrc.org/our-work/health-and-care/community-health/non-communicable-diseases#:~:text=Non%2Dcommunicable%20diseases%20are%20diseases,%2D%20and%20middle%2Dincome%20countries,2022. .](https://www.ifrc.org/our-work/health-and-care/community-health/non-communicable-diseases#:~:text=Non%2Dcommunicable%20diseases%20are%20diseases,%2D%20and%20middle%2Dincome%20countries,2022.)
- [11] WHO, "Global Status report on non-communicable Disease 2014," Geneva, 2014.
- [12] RI Ministry of Health, "Indonesian Health Profile 2015," Center for Data and Information, Indonesia, 2015.
- [13] H. Nawawi, *Social Field Research Methods*. Yogyakarta: Gajah Mada University Press, 2015.
- [14] E. Citizen and Nida Nabilah Nur, "Risk Factors of Behavioral Non-Communicable Diseases," *majority*, vol. 5, no. 2, pp. 88–94, 2016, [Online]. Available: <http://joke.kedokteran.unila.ac.id/index.php/majority/article/view/1082>.
- [15] N. Suryani, Noviana, and O. Libri, "The Relationship between Nutritional Status, Physical Activity, Consumption of Fruits and Vegetables with Hypertension in the Internal Medicine Polyclinic at Idaman Hospital, Banjarbaru City," *J. Health. Indonesia.*, vol. 10, no. 2, pp. 100–107, 2020.
- [16] BF Kurnianingtyas, Suyatno, and MIMI Kartasurya, "Risk Factors for Hypertension in High School Students in Semarang City in 2016," *J. Health. Mass.*, vol. 5, no. 2, pp. 70–77, 2017, doi: <https://doi.org/10.14710/jkm.v5i2.16372>.
- [17] Irwan, *Epidemiology of Non-Communicable Diseases*. Yogyakarta: Deepublish, 2016.
- [18] PS Hidayangsih, "Reproductive Health Problems And Risk Behavior Among Adolescence," *J. Health. R.*, vol. 5, no. 2, pp. 89–101, 2014, doi: <https://dx.doi.org/10.22435/kespro.v5i2.3886.89-101>.
- [19] RI Ministry of Health, *General Guidelines for Non-Communicable Diseases Integrated Development Post*. Jakarta, Indonesia: Indonesian Ministry of Health, 2020.
- [20] NA Antimas, H. Lestari, and JR Afa, "Survey of Risk Factors for Non-Communicable Diseases in Halu Oleo University Students in 2017," *J. Ilm. Mhs. healthy. Mass.*, vol. 2, no. 6, pp. 1–13, 2017, doi:

Paper's should be the fewest possible that accurately describe ... (First Author)










- <http://dx.doi.org/10.37887/jimkesmas.v2i6.2878>.
- [21] Salimar, B. Setyawati, and R. Rachmawati, "Adolescent Lifestyle Behavior at Risk Related to Non-Communicable Diseases in Indonesia," *Researcher. Nutrition and Food*, vol. 4, no. 1, pp. 11–20, 2021, doi: <https://doi.org/10.22435/pgm.v4i1.3337>.
- [22] S. Suryanti, S. Sudarman, and Aswadi, "Relationship between Lifestyle and Diet with the Incidence of Diabetes Mellitus at Bhayangkara Hospital, Makassar City," *J. Promot. Prev.*, vol. 4, no. 1, pp. 1–9, 2021, doi: [10.47650/jpp.v4i1.246](https://doi.org/10.47650/jpp.v4i1.246).
- [23] N. Widyasari, "Relationship of Respondent's Characteristic with The Risk of Diabetes Mellitus and Dyslipidemia at Tanah Kalikedinding," *J. Berk. Epidemiol.*, vol. 5, no. 1, pp. 130–141, 2017, doi: [10.20473/jbe.v5i1](https://doi.org/10.20473/jbe.v5i1).
- [24] N. Fadhilah, "Consumption of Foods Risk Factors Causing Non Infectious," *J. Health.*, vol. Volume VII, pp. 102–107, 2019, doi: <https://doi.org/10.52657/jik.v8i2.1059>.
- [25] MD Izhar and M. Syukri, "Contribution of Age and Risky Food Consumptions Towards Heart Disease in Jambi Province," *J. Ilm. healthy.*, vol. 4, no. 1, pp. 25–37, 2022, doi: [10.36590/if.v4i1.212](https://doi.org/10.36590/if.v4i1.212).
- [26] NU Tanjung, AP Amira, N. Muthmainah, and S. Rahma, "Junk Food and its Relation to Overnutrition in Adolescents," *J. Ilm. healthy. Mass.*, vol. 14, no. 3, pp. 133–140, 2022, doi: <https://doi.org/10.52022/jikm.v14i3.343>.
- [27] H. Yuningrum, H. Trisnowati, and NN Rosdewi, "Risk Factors for Non-Communicable Diseases (PTM) in Adolescents: Case Studies in Public and Private High Schools in the City of Yogyakarta," *J. Formal (Scientific Forum) Respati Public Health*, vol. 6, no. 1, p. 41, 2021, doi: [10.35842/formil.v6i1.343](https://doi.org/10.35842/formil.v6i1.343).
- [28] U. Kalsum, O. Lesmana, and DR Pertiwi, "Patterns of Non-Communicable Diseases and Their Risk Factors in the Anak Dalam Tribe in Nyogan Village, Jambi Province," *Health Media. Mass. Indonesia.*, vol. 15, no. 4, p. 338, 2019, doi: [10.30597/mkmi.v15i4.7062](https://doi.org/10.30597/mkmi.v15i4.7062).
- [29] W. Widagdo and I. Yulita, *Get to know the risk factors for non-communicable diseases (PTM) and their control*, no. 47. South Jakarta, 2018.
- [30] Cicilia L, WP Kaunang, FLFG Langi, and Langi LFG, "the relationship between physical activity and the incidence of diabetes mellitus in outpatients at the Bitung City General Hospital," *J. HEALTH*, vol. 7, no. 5, pp. 1–6, 2018.
- [31] M. Cristanto, M. Saptiningsih, and MY Indriarini, "Relationship between Physical Activity and Prevention of Hypertension in Young Adults: Literature Review," *J. Friends of Nursing*, vol. 3, no. 01, pp. 53–65, 2021, doi: [10.32938/jsk.v3i01.937](https://doi.org/10.32938/jsk.v3i01.937).
- [32] H. Hariawan, M. Tidore, and GZ Rahakbau, "Non-Communicable Disease Prevention Behavior in Ambon Adolescents," *J. Integrated Nursing*, vol. 2, no. 1, pp. 15–21, 2020, doi: <https://doi.org/10.32807/jkt.v2i1.46>.
- [33] Republic of Indonesia Ministry of Health, "Germas Creating Healthy Indonesia," www.Kemendes.go.id, 2016.
- [34] R. Sekarrini, RE Kurniawan, NA Makrifatullah, N. Rosar, Y. Triana, and K. Kunci, "Description of Risk Factors for Non-Communicable Diseases in Umban Sari Village, Rumbai District, Pekanbaru Using a Stepwise Who Approach," *J. Ilm. Multi Discipline Indonesia.*, vol. 1, no. 8, pp. 163–173, 2022, doi: <https://doi.org/10.32670/ht.v1i8.1929>.
- [35] A. Hidayah, FA Simamora, FTS Siregar, H. Oktasia, and K. Sihotang, "Health education about risk food and impact on health," *J. Servants. Mass. Aufa.*, vol. 4, no. 2, pp. 46–48, 2022, doi: <https://doi.org/10.51933/jpma.v4i2.846>.
- [36] Radiah, C. Mintjelungan, and NW Mariati, "Description of Caries Status and Patterns of Dental and Oral Health Care for Students from Ternate in Manado," *e-GEAR*, vol. 1, no. 1, pp. 45–51, 2013, doi: [10.35790/eg.1.1.2013.1929](https://doi.org/10.35790/eg.1.1.2013.1929).
- [37] SA Hamid, R. Kundre, and Y. Bataha, "The relationship between diet and dental caries in grade IV children aged 8-9 years at SD Negeri 126 Manado, Environment 1 Kleak, Malalayang District, Manado City, North Sulawesi Province," *Nursing e-Journal (eKp)*, vol. 5, no. 2, pp. 1–6, 2017, doi: <https://doi.org/10.35790/jkp.v5i2.17870>.
- [38] D. Olczak-Kowalczyk, J. Tomczyk, D. Gozdowski, and U. Kaczmarek, "Cigarette smoking as an oral health risk behavior in adolescents: A cross-sectional study among Polish youths," *Anthropol. Rev.*, vol. 83, no. 1, pp. 53–64, 2020, doi: [10.2478/anre-2020-0007](https://doi.org/10.2478/anre-2020-0007).
- [39] KCSS Htun and K. Peltzer, "Oral health-related quality of life among community dwelling middle-aged and older adults in an urban area in Magway region, Myanmar," *Nagoya J. Med. sci.*, vol. 81, no. 1, pp. 103–112, 2019, doi: [10.18999/nagjms.81.1.103](https://doi.org/10.18999/nagjms.81.1.103).
- [40] I. Jayasvasti, KCSS Htun, and K. Peltzer, "Self-rated oral health status and social and health determinants among 35-65 year-old persons in one region in Myanmar: A cross-sectional study," *Clin.*

- Cosmetics. Investig. Dent.*, vol. 11, p. 339–348, 2019, doi: 10.2147/CCIDE.S227957.
- [41] E. Goldberg, J. Eberhard, A. Bauman, and BJ Smith, “Mass media campaigns for the promotion of oral health: a scoping review,” *BMC Oral Health*, vol. 22, no. 1, pp. 1–19, 2022, doi: 10.1186/s12903-022-02212-3.
- [42] J. Botelho *et al.*, “An umbrella review of the evidence linking oral health and systemic non-communicable diseases,” *Nat. Commun.*, vol. 13, no. 7614, pp. 1–11, 2022, doi: 10.1038/s41467-022-35337-8.
- [43] P. Akera, SE Kennedy, R. Lingam, MJ Obwolo, AE Schutte, and R. Richmond, “Effectiveness of primary school-based interventions in improving oral health of children in low- and middle-income countries: a systematic review and meta-analysis,” *BMC Oral Health*, vol. 22, no. 264, pp. 1–21, 2022, doi: 10.1186/s12903-022-02291-2.
- [44] A. Riad *et al.*, “Oral Health-Related Knowledge, Attitudes and Behaviors of Arab Dental Students: Multi-National Cross-Sectional Study and Literature Analysis 2000–2020,” *Int. J. Environ. Res. Public Health*, vol. 19, no. 3, 2022, doi: 10.3390/ijerph19031658.
- [45] J. Heet *et al.*, “Socio-demographic factors, dental status, oral health knowledge and attitude, and health-related behaviors in dental visits among 12-year-old Shenzhen adolescents: a multilevel analysis,” *BMC Oral Health*, vol. 22, no. 1, pp. 1–11, 2022, doi: 10.1186/s12903-022-02110-8.
- [46] WHO, “Oral Health,” <https://www.who.int/news-room/fact-sheets/detail/oral-health>, 2022. .
- [47] WHO, *Global oral health status report*, vol. 57, no. 2. World Health Organization, 2022.
- [48] CZ Wuet *et al.*, “Epidemiologic relationship between periodontitis and type 2 diabetes mellitus,” *BMC Oral Health*, vol. 20, no. 1, pp. 1–15, 2020, doi: 10.1186/s12903-020-01180-w.




BIOGRAPHIES OF AUTHORS

The recommended number of authors is at least 2. One of them as a corresponding author.

Please attach clear photo (3x4 cm) and vita. Example of biographies of authors:

	<p>Sitti Nur Djannah   is a full senior lecturer and head of the Master of Public Health study program at the Faculty of Public Health, Ahmad Dahlan University, Yogyakarta. My areas of interest are health promotion and behavioral social sciences. She can be contacted at email: sitti.nurdjannah@ikm.uad.ac.id</p>
	<p>Lina Handayani   is a full senior lecturer of Faculty Of Public Health At Ahmad Dahlan University In Yogyakarta, Indonesia. My interest in health education and promotion. She can be contacted at email: lina.handayani@ikm.uad.ac.id</p>
	<p>Rochana Ruliyandari   is a full senior lecturer of Faculty Of Public Health At Ahmad Dahlan University In Yogyakarta, Indonesia. My interest in Health education and promotion. My research Interest in Elderly Health, Elderly Health Promotion, Management of Elderly Health Services, Leadership Learning Organization. She can be contacted at email: rochana.ruliyandari@ikm.uad.ac.id</p>

Paper's should be the fewest possible that accurately describe ... (First Author)

 A portrait photograph of a woman with dark hair, wearing a black hijab and a dark blazer over a white collared shirt. The background is a solid red color.	<p>Intan Wahyuni Tukiyo   is a Researcher Assistant and freshgraduate from Master Of Tropical Medicine At Gadjah Mada University. My research interest in public health, epidemiology, tropical disease and tropical infectious disease. She can be contacted at email: intanwahyunitukiyo@gmail.com</p>
--	--

Sitti Nur Djannah, Lina Handayani, Rochana Ruliyandari, Intan Wahyuni Tukiyo

ORIGINALITY REPORT

13%

SIMILARITY INDEX

13%

INTERNET SOURCES

9%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	Submitted to Universitas Indonesia Student Paper	2%
2	www.researchgate.net Internet Source	2%
3	ijphs.iaescore.com Internet Source	2%
4	Submitted to Mahidol University Student Paper	1%
5	www.moh.gov.jm Internet Source	1%
6	en.tatoli.tl Internet Source	1%
7	Agus Salim Burhanuddin, Katmini, Yuly Peristiowati. "Prevention Program Performance Improvement Strategy and Control of Non-Communicable Diseases at Blega Health Center, Bangkalan District", Journal for Quality in Public Health, 2023 Publication	1%

8	lppm.ub.ac.id Internet Source	1 %
9	download.atlantis-press.com Internet Source	1 %
10	www.ijstm.inarah.co.id Internet Source	1 %
11	bmcoralhealth.biomedcentral.com Internet Source	1 %
12	www.science.gov Internet Source	1 %
13	repo.stikmuhptk.ac.id Internet Source	1 %
14	Submitted to Rwanda Defence Forces Command and Staff College Student Paper	1 %
15	www.who.int Internet Source	1 %

Exclude quotes On

Exclude bibliography On

Exclude matches < 1%