

The relationship between drug therapy adherence and the results of blood pressure targets achievement in hypertensive patients with dyslipidemia complications at Government Hospital in Yogyakarta

Submission date: 18-Sep-2023 01:00PM (UTC+0700)
by Andriana Sari

Submission ID: 2169309576

File name: Pharmaciana_vol_10_no_3_2020.pdf (479.46K)

Word count: 4004

Character count: 21732

The relationship between drug therapy adherence and the results of blood pressure targets achievement in hypertensive patients with dyslipidemia complications at Government Hospital in Yogyakarta

Andriana Sari*, Faridah Baroroh, Fitriani Puji Astuti

Faculty of Pharmacy, Universitas Ahmad Dahlan
Jl. Prof. Dr. Soepomo, Janturan, Yogyakarta, Indonesia

Submitted: 13-05-2020

Reviewed: 15-06-2020

Accepted: 30-11-2020

ABSTRACT

Hypertension is a chronic disease that affects most people. Almost every people have certain specific factors that can cause this disease to increase. One of which is inadequate physical activity so that cholesterol levels remain and tend to get higher, which is a risk factor for heart and blood vessel disorders. Patients with this disease and having dyslipidemia complications must treat with appropriate therapy until reaching success. The patient's adherence determines the success of the treatment. Therefore the purpose of this research is to assess the relationship. Its objects were hypertensive patients at the Government Hospital in Yogyakarta. The specialty of the hypertensive patients having dyslipidemia complications. The study uses non-experimental with a cross-sectional study. The sample was selected based on the purposive sampling technique in December 2018-February 2019. The measurement of therapy adherence used MARS questionnaire and the achievement of therapy targets through secondary data on patient blood pressure. Chi-square test was chosen to determine the relationship between 2 variables, which are therapy adherence and achieving blood pressure targets. The sample in this study were 47 respondents, of which 53.2% of the patients were <60 years old, in which 51.1% of them were female. The results showed that patients who were adherent to therapy were 25.5%, and those who were not were 74.5%. The results of achieving the blood pressure target are known to be 68.1%. The data analysis results showed that the one-way hypothesis in Fisher's test obtaining p-value = 0.171 ($p > 0.05$) was no relationship between adherence and target achievement. Therefore the results of this study concluded that there is no relationship between drug therapy adherence and blood pressure targets achievement.

Keywords: hypertensive, dyslipidemia, adherence, blood pressure target achievement, MARS

*Corresponding author:

Andriana Sari

Faculty of Pharmacy, Universitas Ahmad Dahlan

Jl. Prof. Dr. Soepomo, Janturan, Yogyakarta, Indonesia

Email: andriana@pharm.uad.ac.id

INTRODUCTION

Hypertension is a disease that affects many people in the world. The disease presents an increase of more than 140 mmHg at systolic blood pressure and more than 90 mmHg at diastolic (Weber et al., 2014). The prevalence of hypertension in Indonesia occurs at age > 18 years as much as 25.8% (Kesehatan, 2013). In Yogyakarta, most of this disease is 26.7% (Anonim, 2015), and is the second of the ten most common conditions in Bantul district (Dinkes, 2014).

The increase of blood pressure causes the heart to work faster, and blood does not flow properly, so there is a possibility that there will be a decrease in the function of several organs in the body. The factors causing this kind of occurrence in each individual can be so varied that it is not easy to determine. Some of the most common causes are eating high salted foods, energy-dense foods, fatty foods, and a lack of fruit and vegetables. These all can increase the risk of increased blood pressure and cardiovascular disease (Ozemek et al., 2018). World health organization (Day, 2013) estimates that high cholesterol levels (hypercholesterolemia) can lead to stroke (20%) and heart problems (> 50%). And to prevent this occurrence, drug therapy must be successful.

The patient's adherence to taking the drug influencing the therapy's success, so that blood pressure targets achieved and prevent disease complications. Patients with hypertension who do not adhere to taking medication according to the rules of use that have been conveyed by health workers cause uncontrolled blood pressure and lead to death (Gousel-Wood et al., 2009). Therefore it is necessary to do findings in drug therapy adherence and the achievement of blood pressure targets in hypertensive patients with dyslipidemia complications at the Government Hospital in Yogyakarta.

MATERIALS AND METHOD

The non-experimental findings with a cross-sectional study were chosen. Its primary data is MARS questionnaire to determine patient adherence to drug therapy, and its secondary data from medical records is blood pressure form to assess the achievement of blood pressure targets. The study samples were hypertensive patients with dyslipidemia complications having the inclusion or exclusion criteria and was undergoing outpatient care at the Government Hospital in Yogyakarta. The inclusion criteria were male or female patients, aged ≥ 18 years, with antihypertensive therapy for at least three months. The exclusion criteria were pregnant women with pre-eclampsia, kidney and liver disorders, incomplete medical record data, and questionnaires.

The initial findings procedure was by applying for a finding permit and obtaining ethical approval number 011805083 from the Findings Ethics Committee of Ahmad Dahlan University. The selection of respondents is based on diagnosis by doctors according to inclusion and exclusion criteria. The respondents have explained the study to be carried out and were asked for approval by filling out the consent form. They must answer MARS questionnaire to measure adherence in taking drugs after the last three months of consumption. The next step was checking the questionnaire results and collecting blood pressure data as much as 1x measurement after 3 months of taking the drug through medical record data. Univariate analysis was used to determine the percentage of respondent characteristics, including age, gender, education level, occupation, habits (exercise, diet, smoking, coffee and alcohol consumption), disease complications, medications received by patients, adherence, and blood pressure. Bivariate analysis using Chi-square was used to determine the relationship between 2 variables: therapy adherence and achievement of blood pressure targets (significance value $p < 0.05$).

RESULT AND DISCUSSION

Patient characteristics data

The hypertensive patients' characteristics with dyslipidemia complications in Government Hospital in Yogyakarta are presented in Table 1. The results showed that patients aged <60 years were 53.2%. (Kishore et al., 2016) says that the potential for hypertension will be more significant at the

age of over 35 years (21.8%) with an odds ratio of 3.60 (range ratio 2.11-6.15) and p-value = 0.01. It means that people over 35 years old are more at risk of developing hypertension.

Table 1. Chi-square analysis on the relationship between patient characteristics and therapy adherence, and chi-square analysis on the relationship between characteristics and blood pressure targets achievement of hypertensive patients with dyslipidemia complications at the Government Hospital in Yogyakarta in the period December 2018-February 2109

Category	Patient characteristics		Chi-square Analysis (p-value)	
	Number of patients (n=47)	Percentage (100%)	Therapy adherence	Achievement of Therapy Target
Age				
< 60 years	25	53.2	0.057	0.495
≥ 60 years	22	46.8		
Sex				
Man	23	48.9	0.285	0.206
Woman	24	51.1		
Occupation				
Unemployed	8	17.0	0.203	0.327
Employed	39	83.0		
Education				
Did not complete basic education	6	12.7	0.324	0.469
Completed basic education	41	87.3		
Lifestyle				
Exercise	: Yes	32	0.099	0.303
	: No	15		
Diet	: Yes	31	0.227	0.238
	: No	16		
Habit	: Good	29	0.038*	0.320
	: Bad	18		
Types of complications				
Dyslipidemia/CAD+DM	35	74.5	0.481	0.281
Dyslipidemia+other complications	12	25.5		

Note: CAD= coronary artery disease, DM= Diabetes Mellitus

*significance <0.05 means that there is a significant association between patient characteristics based on lifestyle and drug therapy adherence

The research was dominated by women (51,1%). Women usually get high blood pressure after menopause. Findings on this have been conducted on thousands of women in America. The results showed that higher cholesterol levels in middle-aged women were a potent cause of hypertension. In contrast, women with high HDL were found to have a decreased risk of hypertension slightly (Nikolov et al., 2015). The results showed that the majority of patients were workers (83.0%). Low physical activity in office workers or people who sit more than 5 hours a day, working with relying on automatic machines, are more at risk of suffering hypertension (Bosu, 2017). The findings data shows that most of the patients' latest education has graduated from primary education so they may better understand drug use instructions to improve therapy adherence and achieve blood pressure targets.

The results showed that the lifestyle of patients who did physical activity/sports was 68.1%. Physical activity or sports are part of the risk of hypertension (Hardati and Ahmad, 2013). The results also showed that patients who did a diet were 66.0%. A finding in Africa showed that excess fat consumed is the cause of hypertension (Ramirez et al., 2010). So the findings result is divided into two categories of habits, which are good and less.

Good habits are defined when the patient does not consume alcohol, tea, coffee, or smoke. Bad habits are determined when the patient drinks alcohol, tea, coffee, and smokes, or just one of these. The results showed that the patients had good habits as much as 61.7%. Drinking coffee can increase blood pressure briefly but then return to normal. There is no significant relationship between coffee drinking habits and increased blood pressure in adults (Bistara and Kartini, 2018).

The findings showed that most complications suffered by patients were hypertension with dyslipidemia complications (CAD) and Diabetes Mellitus (74.5%). They were also had other diseases such as OA, IHD, asthma, cholelithiasis, CHF, BPH, HHD, hyperuricemia, GERD, and cephalgia. Based on patients' characteristics toward therapy adherence, it concluded that there was no significant relationship ($p > 0.05$) on each aspect unless the factors of lifestyle in the form of customs patients do not consume alcohol, tea, coffee, or smoke obtained p -value = 0.038. The relationship between patient characteristics and therapeutic targets' achievement also showed no significant relation ($p > 0.05$).

Drug profile

The results of the study on antihypertensive and antidyplipidemic drugs obtained by patients are presented in Table 2.

Table 2. Antihypertensive and antidyplipidemic drugs received by hypertensive patients with dyslipidemia complications at the Government Hospital in Yogyakarta for the period December 2018-February 2019

Antihypertensive Therapy	Antidyplipidemic Therapy			Total (%)
	Statin	Fibrate	Combination statin+fibrat	
CCB (amlodipine)	5 (10.7%)	3 (6.4%)	1 (2.1%)	47 (100)
ARB (irbesartan, candesartan)	4 (8.5%)	2 (4.3%)	2 (4.3%)	
Beta Bloker (bisoprolol)	0 (0%)	1 (2.1%)	0 (0%)	
ARB+Beta Bloker (valsartan/candesartan+ bisoprolol)	1 (2.1%)	4 (8.5%)	0 (0%)	
ARB+CCB (irbesartan/candesartan/ valsartan+ amlodipine)	8 (17.0%)	5 (10.7%)	5 (10.7%)	
ARB+Diuretic (candesartan+ furosemide)	0 (0%)	1 (2.1%)	0 (0%)	
ARB+ARB (valsartan+candesartan)	0 (0%)	1 (2.1%)	0 (0%)	
ARB+CCB+ Beta Bloker (valsartan+ amlodipine+ bisoprolol)	0 (0%)	1 (2.1%)	0 (0%)	
ARB+Diuretic+Beta Bloker (candesartan+furosemide+ bisoprolol)	1 (2.1%)	0 (0%)	0 (0%)	
ARB+ARB+ Beta Bloker (candesartan+ irbesartan+ bisoprolol)	1 (2.1%)	0 (0%)	0 (0%)	
ARB+Diuretic (candesartan,amlodipine, furosemide)	0 (0%)	1 (2.1%)	0 (0%)	

Note: CCB: Calcium Channel Blocker, ARB: Angiotensin Receptor Blocker,

Data on antihypertensive drug therapy were accepted from the findings. Those are single therapy, a combination of two drugs, and an assortment of three drugs from various antihypertensive groups. This available antihypertensive therapy is used in patients with stage I hypertension, and two or three combination therapy is used in patients with stage II hypertension. (Bakris et al., 2003) Antihypertensive and antidyplipidemic drugs, which were often used by patients, were CCB combination with ARB and statin (17%). (Last et al., 2017) suggest that statin therapy should be prescribed for secondary prevention in patients with known ASCVD (atherosclerotic cardiovascular disease), unless contraindicated. The more significant amount of drugs used by the patient may affect adherence in taking drugs.

Therapeutic adherence

Therapeutic adherence in the research was measured based on the results of filling out MARS questionnaire. Measurement of patient adherence to therapy was carried out once after consuming the doctor's drug for three months. To measure the consumption of antihypertensive and anti-cholesterol drugs, they were given two MARS questionnaires. Patients are said to be adherent in taking drugs if the total score from the measurement of antihypertensive and anti-cholesterol therapy is known to have each score = 25. It is said to be non-adherent if the score of the two measures or one of them is a score <25. Data based on patient therapy are presented in Table 3.

Table 3. Therapy adherence of hypertensive patients with dyslipidemia complications at Government Hospital in Yogyakarta for period December 2018-February 2019

Therapy Adherence	Numbers of Respondent (n=47)	Percentage (100%)	P-Value
Adherent	12	25,5	14
Non-adherent	35	74,5	0,000*

*p-value <0.05 means that there are significant differences in patients who are adherent and non-adherent

The results obtained 12 patients (25.5%) who adhere to drug therapy, antihypertensive, and anti-cholesterol drugs. The finding's results were following (Hazwan and Pinatih, 2017), where the level of adherence to hypertension drugs is relatively high for about 30%. Adherence to drug therapy affects the patient's blood pressure because it can be controlled. Patients who reach their target blood pressure are considered to have high levels of adherence. From Maan Withney's difference test, groups of patients who adhered and did not adhere to taking the drug toward the score obtained a value of $p = 0.000$. It is showed that there is a significant difference in scores between patients who are adherent and non-adherent in taking the drug. Uncontrolled hypertension for a long time can lead to heart attacks, enlarged hearts, and heart failure. Hypertension causes the blood vessels' muscle wall to weaken, thus developing a bulge (aneurysm), which is thought to be a blockage and rupture of the blood vessel. The high pressure in the blood vessels can leak or bleed, resulting in a stroke. Hypertension is also known to give rise to kidney failure, vision problems, and cognitive impairment (WHO, 2015).

In Table 4, some patients stated that they forgot to take drugs. No one reminds them about the time, the amount of drug that must be consumed, and because of being busy. Inadvertently forgetting to take a drug is also because the patient feels that their body condition has improved.

Table 4. Duration of taking hypertension drugs on the MARS questionnaire in hypertensive patients with dyslipidemia complications at Government Hospital in Yogyakarta in the period December 2018-February 2019

Questions	Selection of questions					Total
	Always	Often	Occasionally	Rarely	Never	
Fail to take drug	0 (0%)	4 (8,5%)	11 (23,4%)	16 (34,0%)	16 (34,0%)	
Change the dosage	0(0%)	0 (0%)	3 (6,4%)	3 (6,4%)	41 (87,2%)	
Stop to take drug	0 (0%)	1 (2,1%)	10 (21,3%)	11 (23,4%)	25 (53,2%)	47
Decided not to adhere the dosage	0 (0%)	2 (4,3%)	3 (6,4%)	4 (8,5%)	38 (80,9%)	(100%)
Take less than the rule	0 (0%)	1 (2,1%)	4 (8,5%)	5 (10,6%)	37 (78,7%)	

Achievement of therapeutic targets

Table 5. Achievement of therapy targets in hypertensive patients with dyslipidemia complications at the Government Hospital in Yogyakarta for the period December 2018-February 2019

Therapy Achievement	Numbers of Respondents n=47	Percentage (100%)	P-Value
Achieved	32	68,1	1 0,000
Not Achieved	15	31,9	

* p-value <0.05 means that there is a significant difference between patients who have achieved and not achieved blood pressure target

The results in Table 5. showed that 68.1% of patients suffering from hypertension with dyslipidemia complications at the Government Hospital in Yogyakarta achieved the blood pressure target as the therapy, according to JNC 8. The findings showed that the blood pressure gets achievement of hypertensive patients without diabetes complications at age <60 years old was <140/90 mmHg, at age ≥60, years old was <150/90 mmHg, and whose diabetes complications <140/90 mmHg (Dennison-himmelfarb, 2014). The blood pressure of patients with an exact lower limit of 140/90 mmHg or 150/90 mmHg is included in the unachieved category. The factor that causes blood pressure target to be unachieved is non-adherent with the drug therapy. The Mann Withney's different test results showed the value of p = 0.000, so this indicates a significant difference in blood pressure results between patients who achieved the target and was unachieved.

Chi-square analysis: the relationship between therapy adherent towards blood pressure target achievement

A Chi-square test was used to analyze the relationship between therapeutic adherence and the result of blood pressure target achievement. The Findings selected Fisher's trial because the 2x2 table did not meet the Chi-square requirements, in which more than 20% of cells had a smaller expected than 5 (Dahlan, 2014). The one-way hypothesis obtained a significance value of p = 0.171, which means that the required conditions were unfulfilled, so there was no relationship between patient adherence to the blood pressure target. The Odds Ratio value (CI 95%) was 2.955 (0.559 - 15.630). The findings showed that therapeutic adherent has a risk of 2,955 times to affect blood pressure compared with non-adherents. The patient's blood pressure probably influences it at the target limit, categorized as unachieved by the findings. Another factor that may have influenced the results could also be because the patient was classified as adhering if all five questions were met with a total score of 25, so it is necessary to include a limit value in future findings. Chi-square test analysis of the relationship between therapy adherence with the results of blood pressure target can be seen in Table 6.

Table 6. Chi-square test analysis of the relationship between therapy adherence with the results of blood pressure target in Government Hospital in Yogyakarta for the period December 2018-February 2019

Therapy Adherent	Therapy of Target Achievement		Total	p-value	OR (CI 95%)
	Achieved	Unachieved			
Adhere	10 (21.3%)	2 (4.2%)	47	0.171	2.955 (0.559 - 15.630)
Not Adhere	22 (46.8%)	13 (27.7%)	(100%)		

Note: p-value (sig> 0.05) means no relationship between therapy adherence and target achievement.

7 The results showed that most of the respondents had diabetes mellitus (74.5%) as a comorbid disease. The chi-square statistical test results on the relationship between patients' characteristics related to comorbid and therapeutic adherence obtained a p-value of 0.481 (> 0.05). The findings conclude that there is no comorbid influence on therapy adherence. The chi-square statistical test results for the relationship between patients' characteristics related to comorbid and the achievement of blood pressure targets obtained a p-value of 0.281 (> 0.05). Table 1 shows that comorbid does not affect blood pressure targets achievement as presented. The findings were different from (Cahyani, 2018). It concluded a significant relationship between adherence to taking antihypertensive drugs and blood pressure (p-value = 0.005), with hypertensive patients with diabetes mellitus complications considering that this patient has a high adherence level. Comorbidities may directly influence the complexity of the treatment, the amount of drug consumed, the cost, and the patient's motivation. In a study conducted, other factors, such as exercise, diet, tea consumption, coffee, and alcohol, also affect not achieving therapeutic targets. The findings lack the data collection that was carried out once when the patient had taken the drug for three months, so that the progress in achieving the target each month was not known. Also, the sample size was limited, only using hypertensive patients with dyslipidemia and other complications.

CONCLUSION

The research concluded that 25.5% of patients adhere to take antihypertensive and antidyslipidemic consumption, and 68.1% of patients achieved their blood pressure. There is no relationship between drug therapy adherence to the results of blood pressure targets achievement in hypertensive patients with dyslipidemia complications at the Government Hospital in Yogyakarta at the period December 2018-February 2019 ($p=0.171$).

15

ACKNOWLEDGEMENT

The authors thank Lembaga Penelitian dan Pengabdian kepada Masyarakat Universitas Ahmad Dahlan Yogyakarta.

REFERENCES

- Anonim. (2015). *Kementerian kesehatan republik indonesia*.
- Bakris, G., Cushman, W. C., Green, L. A., & Izzo, J. L. (2003). The Seventh Report of the Joint National Committee on Prevention, Detection, Evaluation, and Treatment of High Blood Pressure: The JNC 7 Report, 289(May 21), 2560–2572. <https://doi.org/10.1001/jama.289.19.2560>
- Bistara, D. ., & Kartini, Y. (2018). Hubungan Kebiasaan Mengonsumsi Kopi dengan Tekanan Darah pada Dewasa Muda. *Jurnal Kesehatan Vokasional*, 3(1), 24–27.
- Bosu, W. K. (2017). systematic review, 9716(August). <https://doi.org/10.3402/gha.v8.26227>
- Cahyani, F. M. (2018). Hubungan Kepatuhan Minum Obat Antihipertensi Terhadap Tercapainya Target Terapi Pasien Hipertensi di Puskesmas Wirobrajan Yogyakarta Melengkapi data primer yang diperoleh dari wawancara langsung dengan data pada rekam medik pasien hipertensi yang menjalani pengobatan rawat jalan di Puskesmas Wirobrajan Yogyakarta ., 1(2), 10–16.
- Dahlan, M. . (2014). *Statistik untuk Kedokteran dan Kesehatan Edisi 6*.
- Day, W. H. (2013). A global brief on Hyper - tension World Health Day 2013.
- Dennison-himmelfarb, C., Handler, J., & Lackland, D. T. (2014). 2014 Evidence-Based Guideline for the Management of High Blood Pressure in Adults Report From the Panel Members Appointed to the Eighth Joint National Committee (JNC 8), 1097(5), 507–520. <https://doi.org/10.1001/jama.2013.284427>
- Dinkes. (2014). *District Health Account*.
- Hardati, A. T., & Ahmad, R. A. (2013). Aktivitas fisik dan kejadian hipertensi pada pekerja : analisis data Riskesdas 2013.

- Hazwan, A., & Pinatih, G. N. . (2017). Gambaran karakteristik penderita hipertensi dan tingkat kepatuhan minum obat di wilayah kerja puskesmas Kintamani I, 8(2), 130–134. <https://doi.org/10.1556/ism.v8i2.127>
- Indonesian Ministry of Health. (2015). *Indonesian Health Statistics 2014. Kementerian Kesehatan Republik Indonesia* (Vol. 51). <https://doi.org/10.1037/0022-3514.51.6.1173>
- Kesehatan, K. (2013). Riset kesehatan dasar. Badan penelitian dan pengembangan kesehatan kementerian kesehatan RI.
- Kishore, J., Gupta, N., Kohli, C., & Kumar, N. (2016). Prevalence of Hypertension and Determination of Its Risk Factors in Rural Delhi, 2016.
- Krousel-Wood, M., Islam, T., Webber, L. S., Re, R. N., Morisky, D. E., & Muntner, P. (2009). New medication adherence scale versus pharmacy fill rates in seniors with hypertension. *American Journal of Managed Care*, 15(1), 59–66. [https://doi.org/11024 \[pii\]](https://doi.org/11024 [pii])
- Last, A. R., College, M., Valley, F., Medicine, F., Program, R., Ference, W. J. D., ... Program, R. (2017). Risk Reduction in Adults.
- Nikolov, P., Nikolova, J., Orbecova, M., Deneva T., Vladimirova, L., A. (2015). Flow Mediated Vasodilation and Some Biomarkers of Endotelial Activation in Pre-Hypertensive Object. *The West Indian Medical Journal*, 1–12.
- Ozemek, C., Laddu, D. R., Arena, R., & Lavie, C. J. (2018). The role of diet for prevention and management of hypertension, 1–6. <https://doi.org/10.1097/HCO.0000000000000532>
- Ramirez, S. S. De, Enquobahrie, D. A., Nyadzi, G., Mjungu, D., & Magombo, F. (2010). Prevalence and correlates of hypertension : a cross-sectional study among rural populations in sub-Saharan Africa, 786–795. <https://doi.org/10.1038/jhh.2010.14>
- Weber, M. A., Schiffrin, E. L., White, W. B., Mann, S., Lindholm, L. H., Kenerson, J. G., ... Ram, C. V. S. (2014). Clinical Practice Guidelines for the Management of Hypertension in the Community A Statement by the American Society of Hypertension and the International Society of Hypertension, 16(1). <https://doi.org/10.1111/jch.12237>
- WHO. (2015). Q & As on hypertension, (September).

The relationship between drug therapy adherence and the results of blood pressure targets achievement in hypertensive patients with dyslipidemia complications at Government Hospital in Yogyakarta

ORIGINALITY REPORT

11%

SIMILARITY INDEX

9%

INTERNET SOURCES

5%

PUBLICATIONS

3%

STUDENT PAPERS

PRIMARY SOURCES

1	staff.ui.ac.id Internet Source	1%
2	scholarworks.boisestate.edu Internet Source	1%
3	jurnal.ugm.ac.id Internet Source	1%
4	Submitted to Universitas Muhammadiyah Purwokerto Student Paper	1%
5	Intan Maeilani Rahayu, Iwan Shalahuddin, Kurniawan Yudianto. "An Overview of Blood Pressure Based on Affecting Hypertension Factors in Elderly Stage; Description Study at Garut Elderly Social Rehabilitation Service Unit", Jurnal Aisyah : Jurnal Ilmu Kesehatan, 2020 Publication	1%

6	www.science.gov Internet Source	1 %
7	janh.candle.or.id Internet Source	1 %
8	Submitted to Academic Library Consortium Student Paper	<1 %
9	Lingga Kusuma Wardani, Febriana Ina Lemba, Joel Rey Ugsang Acob. "Risk Factor Analysis in Community with Consuming Habits Caffeinated Drinks", Journal Of Nursing Practice, 2022 Publication	<1 %
10	www.aafp.org Internet Source	<1 %
11	f1000research.com Internet Source	<1 %
12	Moser, M.. "From JNC I to JNC 7-What Have We Learned?", Progress in Cardiovascular Diseases, 200603/04 Publication	<1 %
13	www.coursehero.com Internet Source	<1 %
14	repository.binausadabali.ac.id Internet Source	<1 %
15	eprints.untirta.ac.id Internet Source	<1 %

<1 %

16

P. O. Okunrotifa. "A COMPARISON OF THE RESPONSES OF NIGERIAN PUPILS TO TWO SETS OF PROGRAMMED INSTRUCTION MATERIALS IN GEOGRAPHY", Programmed Learning and Educational Technology, 2006

Publication

<1 %

17

Steven G. Chrysant. "The debate over the optimal blood pressure treatment target of less than 130/80 mmHg", Postgraduate Medicine, 2022

Publication

<1 %

18

repository.ubn.ru.nl

Internet Source

<1 %

19

univendspace.univen.ac.za

Internet Source

<1 %

20

Ahmet Yurteri, Numan Mercan, Mehmet Kılıç, Ahmet Temiz, Fatih Doğar, Duran Topak, Ahmet Yıldırım. "Rotational alignment after intramedullary nailing in trochanteric fractures: Lateral decubitus versus supine position with fracture table. Does operative position matter?", Research Square Platform LLC, 2023

Publication

<1 %

21 B J Que, C Y Ihalauw, F de Lima. "Prevalence of hypertension and efforts to develop a monitoring system for hypertension in community living in the coastal areas of Noloth and Itawaka Village, Saparua Island, Central Maluku District", IOP Conference Series: Earth and Environmental Science, 2019
Publication

22 Santhra Segaran Balan, Rusliza Basir, Yoke Keong Yong, Noor Aishah Mohammed Izham et al. "36th Malaysian Society of Pharmacology and Physiology Annual Scientific Meeting 2023", Life Sciences, Medicine and Biomedicine, 2023
Publication

23 [baixardoc.com](https://www.baixardoc.com)
Internet Source

24 www.medarhive.ru
Internet Source

25 www.researchgate.net
Internet Source

26 www.texilajournal.com
Internet Source

27 Adnan Adnan, Dyah Aryani Perwitasari, Uly Adhie Mulyani. "Reliability and Validaty of St George Respiratory Questionnaire (SGRQ)

into Indonesian Version", International Journal of Public Health Science (IJPHS), 2014 Publication

Exclude quotes On

Exclude bibliography On

Exclude matches Off