

# HASIL CEK\_60160955

*by* Tekim 60160955

---

**Submission date:** 19-Aug-2023 10:26AM (UTC+0700)

**Submission ID:** 2147831314

**File name:** Teknik\_Kimia\_60160955.docx (328.52K)

**Word count:** 2571

**Character count:** 14422

1 **EFFECT OF PINEAPPLE PEEL EXTRACTS ON ORGANIC**  
2 **FERTILIZER NUTRIENTS AND GROWTH OF CAYENNE PEPPER**  
3 **(*CAPSICUM ANNUUM*)**

4  
5 LUKHI MULIA SHITOPHYTA<sup>1\*</sup>, RATIH DEA LARASATI<sup>1</sup>, MAHARANI NUR BELA PERTIWI<sup>1</sup>, MEILYA  
6 SUZAN TRIYASTUTI<sup>2</sup>

7 <sup>1</sup>Department of Chemical Engineering, Faculty of Industrial Technology, Universitas Ahmad Dahlan,  
8 Yogyakarta, 55191, Indonesia

9 <sup>2</sup>Department of Fishery Product Processing Techniques, Politeknik Kelautan dan Perikanan Bitung, Sulawesi  
10 Utara 95526, Indonesia

11 \*Corresponding author: [lukhi.mulia@che.uad.ac.id](mailto:lukhi.mulia@che.uad.ac.id)

12  
13 **Abstract**

14 Pineapple peel extract can be used for manufacturing organic fertilizer due to its high content of carbohydrates,  
15 protein, and sugar. The study aimed to find the effect of pineapple peel extract on fertilizer nutrients and the  
16 growth of cayenne pepper. The experiment was performed in 5 L of plastic containers for 15 days. The result  
17 showed that pineapple peel extract significantly affected fertilizer nutrients and the growth of cayenne pepper (p  
18 < 0.05). The extract of 2500 mL generated the highest height of the plant and the greatest number of leaves. The  
19 extract. The extract of 2000 mL produced low potassium and nitrogen. Lack of potassium and nitrogen could  
20 reduce plant height and change leaf color.

21 **Keywords: cayenne pepper; organic fertilizer; pineapple peel; potassium**

22  
23 **Introduction**

24 Biomass waste comprises rich nutrients and high organic material, which can be put to beneficial use as  
25 fertilizer sources (Chew et al., 2019). Organic fertilizers are naturally obtainable mineral resources containing  
26 adequate amounts of vital plant nutrients. Organic fertilizer can be derived from animal excrement, plant matter  
27 (peanut shells, straw), poultry litter, fish manure, blood meal and cottonseed meal (Shaji et al., 2021). The function  
28 of organic fertilizers affects soil qualities such as soil microbial activity, soil organic material household, soil salt  
29 content and soil pH (Möller & Schultheiß, 2015). Moreover, organic fertilizers can increase yield, soil bulk density  
30 and essential soil minerals such as nitrogen, phosphorus, potassium, calcium and magnesium (Oyetunde-Uzman  
31 et al., 2021).

32 The application of organic fertilizers may not only enhance soil characteristics but may also assist in  
33 climate protection by increasing carbon (C) sequestration in soils. Increasing organic material by sequestering C  
34 in soils may diminish atmospheric CO<sub>2</sub> emissions and lead to continuing losses in the form of CH<sub>4</sub> and N<sub>2</sub>O  
35 emissions by affording substrates for microbial activity and microbial oxygen consumption (Zhang et al., 2016).  
36 Many studies have revealed that applying organic fertilizers to the soil can enhance microbe communities and  
37 increase yield (Ma et al., 2023; Shang et al., 2020; Ye et al., 2022). However, no study has not investigated the  
38 combination of pineapple peels and tofu liquid waste in organic fertilizer and its effect on the growth and yield of  
39 cayenne pepper.

40 Pineapple peels have contents of 81.72% water, 17.53% carbohydrates, 4.41% protein and 13.65%  
41 reducing sugar (Sutikarini et al., 2023). Pineapple peels can be used as the raw material of organic fertilizer  
42 because it has a high content of carbohydrates, sugars, and proteins (Suryani et al., 2022). The previous study who  
43 was conducted by Alasa et al., (2021) also reported that adding organic fertilizer from pineapple peels enhanced  
44 soil fertility, increase plant yield, and improve growth parameters such as sprouting time, plant height, leaf area  
and the number of leaves per branch. Tofu liquid waste contains proteins, amino acids, and plentiful nutrients

45 such as N, P, K, Ca, Mg and Fe. Protein will be released as an N compound by microbes. The N compounds will  
46 be absorbed by the plant roots (Chaorlina<sup>10</sup> al., 2021). Fauziah & Idris (2022) show that application of tofu liquid  
47 waste significantly affected chlorophyll, height, number of flowers, pod length, and fruit weight of beans. Samah  
48 & Candra (2022) also show that organic fertilizer from tofu liquid waste influenced the height, number of leaves  
49 and stem diameter of mustard green. Therefore, many studies have revealed the effect of organic fertilizer on plant  
50 growth. The study aimed to discover the influence of pineapple peel extract combined with tofu liquid waste on  
51 fertilizer nutrients (N, P, and K) and the growth of cayenne pepper (height, number of leaves, and color).

52

## 53 **Materials and Methods**

### 54 **Preparation of pineapple peels extracts**

55 2000 gram of pineapple peels were chopped then mashed using a blender. Mashed pineapple peels were strained  
56 to collect the pineapple peels extracts then stored in tight-closed bottles.

### 57 **Preparation bio activator**

58 1500 mL of tofu liquid waste was mixed with pineapple peels extract at different volumes of 1500 mL, 2000  
59 mL, and 2500 mL. 200 mL of Effective Microorganism- 4 (EM-4) was added into the mixture and poured into 5  
60 L of bottles then stirred well.

### 61 **Production of organic fertilizer**

62 Leaves and vegetable wastes were shredded using vegetable waste shredder. 500 gram of leaves and 1200 gram  
63 of vegetable wastes were introduced into 5 L of plastic drum containers then added bio activator and water, and  
64 stirred thoroughly. The containers were sealed tightly. The fermentation was carried out for 15 days.

### 65 **Analysis of fertilizer**

66 The nutrients of fertilizer were analyzed according to the standard methods. The total N content was determined  
67 by the Kjeldahl method. The total P and K contents were measured by the perchloric acid digestion method and  
68 spectrophotometry procedures (Shang et al., 2020).

### 69 **Statistical analysis**

70 The significance of the results was performed by analysis of variance (ANOVA) using Microsoft Excel 2019. A  
71 p-value less than 0.05 denotes significant influence, while a p-value higher than 0.05 signifies insignificant  
72 impact.

## 73 **Results and Discussions**

### 74 **Effect of pineapple peels on fertilizer contents**

75 The effect of pineapple peels on fertilizer content was investigated in different volumes of 1500 mL, 2000 mL  
76 and 2500 mL. Table 1 presents fertilizer contents at various volumes of pineapple peel extracts.

77 **Table 1. Nutrients of fertilizer at various volumes of pineapple peel extracts**

Pineapple peel extracts (mL)	Nutrient content		
	N	P	K
1500	9.395% ± 0.0002	3.645% ± 0.0005	1.710% ± 0.003
2000	9.395% ± 0.0006	3.810% ± 0.0003	1.580% ± 0.004
2500	9.420% ± 0.0001	3.685% ± 0.0011	1.615% ± 0.004

78

79 The different volumes of extract provided the highest nutrient of each parameter. 2500 mL of extract obtained the  
80 highest N content. The highest P content was obtained by 2000 mL of extracts, while the highest K content was  
81 generated by 1500 mL of extracts. As seen in Table 1, volumes of pineapple peel extract affected the nutrient  
82 content of fertilizer. Statical analysis proved pineapple peels significantly affected the nutrient content of fertilizer  
83 (p < 0.05). Ramadhani & Nuraini (2018) reported that pineapple waste increased the availability of N, P, and K.

84 The total nitrogen contents provided the highest amount among total phosphorus and potassium contents, while  
85 the potassium content had the lowest amount. The results are similar to the previous study by Mustapha et al.  
86 (2021), where pineapple peel had the lowest potassium amount among nitrogen and phosphorus amount. Nitrogen  
87 plays a role in the photosynthesis of plants, and it is a constituent component of protein. Lack of nitrogen can lead  
88 to inadequacy in sunlight absorbance by the plant, resulting in a deficiency of nutrient uptake (Hamid et al., 2019).

89 Hariyanto et al., (2023) stated that nitrogen and phosphorus affected plant growth and yield. They reported that  
90 fertilizer containing nitrogen and phosphorus impacted the number of shoots, flowers and fruit sets on the dragon  
91 fruits' growth. Pineapple peel extracts generated the lowest potassium nutrient among nitrogen and phosphorus,  
92 signifying that fruit peels are essential sources of potassium for plant growth (Mustapha et al., 2021).

### 93 **Effect of pineapple peel extracts on the growth of cayenne peppers**

94 The application of pineapple peel extract was observed towards the growth of cayenne peppers (plant height,  
95 number of leaves and color of leaves). Figure 1 shows the growth of cayenne peppers in different volume extracts



96

97 Figure 1. Growth of cayenne peppers in different volume extracts: (A) 1500 mL; (B) 2000 mL; (C) 2500 mL

98

99 Treatment of 2500 mL extracts had the most significant number of leaves (8 leaves), followed by 1500 mL and  
100 2000 mL with the number of leaves of 6 and 5, respectively. The result showed that pineapple peel extracts  
101 significantly affected the number of leaves ( $p < 0.05$ ). This condition may occur due to richer nitrogen owned by  
102 2500 mL of extract. Increase in nitrogen concentration increased number of leaves. Nitrogen is part of the cause  
103 of the increasing size of cells and the rate of their fission, which results in a substantial increase in growth  
104 characteristics, and the number of leaves is a trait linked to the genotype principally, nevertheless, it is influenced  
105 by growth factors, specifically nitrogen fertilizer and its contents (Hashim et al., 2022).

106 Data on plant heights performs that the treatment of 2000 mL generated a height of 10 cm, while the treatment of  
107 1500 mL as well as the treatment of 2000 mL provided a height plant of 10.2 cm. The results happen since the  
108 extract of 2000 mL had the lowest potassium level. A deficiency of potassium reduces plant height and inhibits  
109 root and leaf respiration as well as photosynthesis (Fontana et al., 2020). The analysis result verified that pineapple  
110 peel extracts provided a significant effect on plant height ( $p < 0.05$ )

111 The use of 2500 mL and 1500 mL extracts gave leaves a color of light green, while 2000 mL of extract produced  
112 yellowish-green of leaves. Yue et al., (2022) revealed a correlation between a lack of nitrogen and changes in color  
113 and shape. The color of the leaves turned yellowish-green, the vein transformed into white, and the shape of the  
114 leaves was deformed unevenly. A previous study conducted by Yang et al. (2016) found that potassium deficiency  
115 turned cotton leaf color from green to yellow and red. The summary of the height of the plant, the color of the  
116 leaves, and the number of leaves can be seen in Table 2.

117 **Table 2. Effect of treatment of extract on the growth of cayenne peppers**

Pineapple peel extracts (mL)	Height of plant (cm)	Number of leaves	Color of leaves
1500	10.2	6	Lime green
2000	10	5	Yellowish-green
2500	10.2	8	Lime green

118

119 **Conclusion**

120 The level of pineapple peel extract significantly affects fertilizer nutrients ( $p < 0.05$ ). The treatment of pineapple  
 121 peel extract substantially influences cayenne pepper's growth, viz the number of leaves, the color of leaves and  
 122 the height of the plant ( $p < 0.05$ ). Pineapple peel extracts produces higher level of nitrogen compared to potassium  
 123 and phosphorus. The low potassium level reduces the plant's height, decreases the number of leaves, and changes  
 124 leaf color.

125 **References**

- 126 Alasa, J. J., Bashir, A. U., Mustapha, M., & Mohammed, B. (2021). Experimental study on the use of banana  
 127 and pineapple peel waste as biofertilizers, tested on hibiscus sabdariffa plant: promoting sustainable  
 128 agriculture and environmental sanitation. *Arid Zone Journal of Engineering, Technology & Environment*,  
 129 17(4), 547–554. <https://azojete.com.ng/index.php/azojete/article/view/537>
- 130 Chaorlina, A., Setyaningsih, M., & Faruq, H. (2021). The utilization of tofu waste water as an addition of  
 131 nutrition in hydroponic media to lettuce growth (*Lactuca sativa* L.). *IOP Conference Series: Earth and  
 132 Environmental Science*, 755(1). <https://doi.org/10.1088/1755-1315/755/1/012049>
- 133 Chew, K. W., Chia, S. R., Yen, H. W., Nomanbhay, S., Ho, Y. C., & Show, P. L. (2019). Transformation of  
 134 biomass waste into sustainable organic fertilizers. *Sustainability (Switzerland)*, 11(8).  
 135 <https://doi.org/10.3390/su11082266>
- 136 Fauziah, N., & Idris, M. (2022). The Effect Of Liquid Tofu Waste And Growing Media On The Growth And  
 137 Yield Of Long Beans (*Vigna sinensis* L.). *Jurnal Bioteknologi & Biosains Indonesia*, 9(October), 217–  
 138 226.
- 139 Fontana, J. E., Wang, G., Sun, R., Xue, H., Li, Q., Liu, J., Davis, K. E., Thornburg, T. E., Zhang, B., Zhang, Z.,  
 140 & Pan, X. (2020). Impact of potassium deficiency on cotton growth, development and potential  
 141 microRNA-mediated mechanism. *Plant Physiology and Biochemistry*, 153, 72–80.  
 142 <https://doi.org/10.1016/j.plaphy.2020.05.006>
- 143 Hamid, H. A., Pei Qi, L., Harun, H., Mohamed Sunar, N., Hanim Ahmad, F., & Suliza Muhamad, M. (2019).  
 144 Development of Organic Fertilizer from Food Waste by Composting in UTHM Campus Pagoh. *JDSE  
 145 Journal of Design for Sustainable and Environment*, 1(1), 1–6. <http://www.fazpublishing.com/jdse>
- 146 Hariyanto, B., Mayura, E., Muas, I., Jumjunidang, & Octrina, L. (2023). Effects of Nitrogen and Phosphorus  
 147 Fertilizer on Yield of Dragon Fruit (*Hylocereus polyrhizus*). *Journal of Applied Agricultural Science and  
 148 Technology*, 7(2), 172–185.
- 149 Hashim, A. A., Hussein, M., & Kareem, A. (2022). The Effect Of Different Levels Of Nitrogen Fertilizer On  
 150 Some Growth Characteristics Of Three Cultivars Of (Sorghum) Sorghum Bicolor.L. *Nat. Volatiles &  
 151 Essent. Oils*, 9(2), 397–405.
- 152 Ma, G., Cheng, S., He, W., Dong, Y., Qi, S., Tu, N., & Tao, W. (2023). Effects of Organic and Inorganic  
 153 Fertilizers on Soil Nutrient Conditions in Rice Fields with Varying Soil Fertility. *Land*, 12(5).  
 154 <https://doi.org/10.3390/land12051026>
- 155 Möller, K., & Schultheiß, U. (2015). Chemical characterization of commercial organic fertilizers. *Archives of  
 156 Agronomy and Soil Science*, 61(7), 989–1012. <https://doi.org/10.1080/03650340.2014.978763>
- 157 Mustapha, M., Alasa, J. J., Bashir, A. U., & Mohammed, B. (2021). Study on the Use of Banana and Pineapple  
 158 Peel Waste As Biofertilizers: Enhancing Soil Fertility, Promoting Sustainable Agriculture and  
 159 Environmental Sanitation. *Arid Zone Journal of Engineering, Technology & Environment*, 17(4), 569–  
 160 574. [www.azojete.com.ng](http://www.azojete.com.ng)
- 161 Oyetunde-Usman, Z., Ogunpaimo, O. R., Olagunju, K. O., Ambali, O. I., & Ashagidigbi, W. M. (2021).

- 162 Welfare Impact of Organic Fertilizer Adoption: Empirical Evidence From Nigeria. *Frontiers in*  
163 *Sustainable Food Systems*, 5(July), 1–17. <https://doi.org/10.3389/fsufs.2021.691667>
- 164 Ramadhani, W. S., & Nuraini, Y. (2018). The use of pineapple liquid waste and cow dung compost to improve  
165 the availability of soil N, P, and K and growth of pineapple plant in an Ultisol of Central Lampung.  
166 *Journal of Degraded and Mining Lands Management*, 6(1), 1457–1465. <https://doi.org/10.15243/jdmlm>.
- 167 Samah, E., & Candra, I. A. (2022). The effect of tofu liquid waste and organic mulch on the growth of cocoa  
168 (theobroma cacao L.) Nursery plants in a polybag. *Journal of Applied Agricultural Science and*  
169 *Technology*, 6(2), 171–179.
- 170 Shaji, H., Chandran, V., & Mathew, L. (2021). Organic fertilizers as a route to controlled release of nutrients. In  
171 *Controlled Release Fertilizers for Sustainable Agriculture*. Elsevier Inc. [https://doi.org/10.1016/b978-0-](https://doi.org/10.1016/b978-0-12-819555-0.00013-3)  
172 [12-819555-0.00013-3](https://doi.org/10.1016/b978-0-12-819555-0.00013-3)
- 173 Shang, L., Wan, L., Zhou, X., Li, S., & Li, X. (2020). Effects of organic fertilizer on soil nutrient status, enzyme  
174 activity, and bacterial community diversity in *Leymus chinensis* steppe in Inner Mongolia, China. *PLoS*  
175 *ONE*, 15(10 October), 1–18. <https://doi.org/10.1371/journal.pone.0240559>
- 176 Suryani, R., Masulili, A., Sutikarini, S., & Tamtomo, F. (2022). Utilization of Liquid Organic Fertilizer of  
177 Pineapple Waste to Improving Growth of Sweet Corn Plant in Red Yellow Podsolc Soil. *International*  
178 *Journal of Multi Discipline Science (IJ-MDS)*, 5(1), 30. <https://doi.org/10.26737/ij-mds.v5i1.3160>
- 179 Sutikarini, S., Masulili, A., Suryani, R., Setiawan, S., & Mulyadi, M. (2023). Characteristics of Pineapple Waste  
180 as Liquid Organic Fertilizer and Its Effect on Ultisol Soil Fertility. *International Journal of Multi*  
181 *Discipline Science (IJ-MDS)*, 6(1), 38. <https://doi.org/10.26737/ij-mds.v6i1.3754>
- 182 Yang, J. S., Hu, W., Zhao, W., Meng, Y., Chen, B., Wang, Y., & Zhou, Z. (2016). Soil Potassium Deficiency  
183 Reduces Cotton Fiber Strength by Accelerating and Shortening Fiber Development. *Scientific Reports*,  
184 6(June), 1–11. <https://doi.org/10.1038/srep28856>
- 185 Ye, S., Peng, B., & Liu, T. (2022). Effects of organic fertilizers on growth characteristics and fruit quality in  
186 Pear-jujube in the Loess Plateau. *Scientific Reports*, 12(1), 1–11. [https://doi.org/10.1038/s41598-022-](https://doi.org/10.1038/s41598-022-17342-5)  
187 [17342-5](https://doi.org/10.1038/s41598-022-17342-5)
- 188 Yue, H. F., Zhou, M., Hou, X. K., Yang, X., Zhao, H., Xu, M. S., Wang, L., Liu, H. M., Wuyun, T. N., Zhu, G.  
189 P., & Li, F. D. (2022). Effects of Nitrogen, Phosphorus and Potassium on Phenotype, Photosynthesis and  
190 Biomass Accumulation At Juvenile Phase of *Prunus Armeniaca* × *Sibirica*. *Pakistan Journal of Botany*,  
191 54(2), 577–588. [https://doi.org/10.30848/PJB2022-2\(11\)](https://doi.org/10.30848/PJB2022-2(11))
- 192 Zhang, M., Li, B., & Xiong, Z. Q. (2016). Effects of organic fertilizer on net global warming potential under an  
193 intensively managed vegetable field in southeastern China: A three-year field study. *Atmospheric*  
194 *Environment*, 145, 92–103. <https://doi.org/10.1016/j.atmosenv.2016.09.024>
- 195

# HASIL CEK\_60160955

## ORIGINALITY REPORT

8%

SIMILARITY INDEX

8%

INTERNET SOURCES

7%

PUBLICATIONS

2%

STUDENT PAPERS

## PRIMARY SOURCES

1	<a href="http://static.frontiersin.org">static.frontiersin.org</a> Internet Source	1%
2	<a href="http://journal.stkipsingkawang.ac.id">journal.stkipsingkawang.ac.id</a> Internet Source	1%
3	N F H Kamaruddin, S Adam, N Q Boll Kassim. "The effects of different pineapple wastes on soil chemical properties and growth and yield of Okra ( <i>Abelmoschus esculentus</i> L.)", IOP Conference Series: Earth and Environmental Science, 2023 Publication	1%
4	<a href="http://journals.plos.org">journals.plos.org</a> Internet Source	1%
5	<a href="http://www.jestr.org">www.jestr.org</a> Internet Source	1%
6	<a href="http://www.ornamentalhorticulture.com.br">www.ornamentalhorticulture.com.br</a> Internet Source	1%
7	<a href="http://www.researchgate.net">www.researchgate.net</a> Internet Source	1%

8

D Sartika, S Astuti, R Iswandari. "Inhibitory Study Of Cassava Leather Ethanol Extract As Natural Antimicrobial In Reducing Salmonella Sp. And Escherichia Coli On Contamination Chicken Meat (Gallus Domesticus)", Journal of Physics: Conference Series, 2021

Publication

1 %

9

Submitted to Universiti Malaysia Kelantan

Student Paper

1 %

10

[repository.uinsu.ac.id](https://repository.uinsu.ac.id)

Internet Source

1 %

Exclude quotes  On

Exclude matches  < 1%

Exclude bibliography  On