

PERCEIVED STRESS SCALE DENGAN MODEL RASCH

by Resti Diah Silviani

Submission date: 11-Oct-2023 01:06AM (UTC-0500)

Submission ID: 2192239983

File name: 17903-53892-1-PB_4.pdf (184.59K)

Word count: 1774

Character count: 9661

PERCEIVED STRESS SCALE DENGAN MODEL RASCH

Arif Budi Prasetya*, Diana Septi Purnama, Fajar Wahyu Prasetyo

*Correspondent Author

Arif Budi Prasetya
Universitas Negeri Yogyakarta
Jalan Colombo Yogyakarta Nomor 1,
Karang Malang, Caturtunggal, Depok,
Kabupaten Sleman, Daerah Istimewa
Yogyakarta
Indonesia
Email: arifbudi.2019@student.uny.ac.id

Diana Septi Purnama
Universitas Negeri Yogyakarta
Jalan Colombo Yogyakarta Nomor 1,
Karang Malang, Caturtunggal, Depok,
Kabupaten Sleman, Daerah Istimewa
Yogyakarta
Indonesia
Email: dianaseptipurnama@uny.ac.id

Fajar Wahyu Prasetyo
Universitas Negeri Yogyakarta
Jalan Colombo Yogyakarta Nomor 1,
Karang Malang, Caturtunggal, Depok,
Kabupaten Sleman, Daerah Istimewa
Yogyakarta
Indonesia
Email:
fajarwahyu.2019@student.uny.ac.id

ABSTRAK

The present study aimed to examine the validity and reliability of perceived stress scale using Rasch model. This quantitative study involved 195 students of guidance and counseling department in a university in Yogyakarta, Indonesia. The data were collected using Perceived stress scale and analyzed using Rasch model. The result of the study showed that perceived stress scale's person reliability score was acceptable (0.79) and item reliability score of 0.98. The person separation score was 1.93 (poor) while the item separation score was 7.17 (very good). Based on misfit test, each item of the scale was fit to Rasch model. TO conclude, the perceived stress scale was valid and reliable, thus can be used by university counselors to measure students' perceived stress level.

Keywords: perceived stress scale, rasch model, university student.

Halaman
48-51

INTRODUCTION

Every individual ideally should go through their developmental task according to their age. University students' developmental task, according to Danim (2013), is to achieve emotional autonomy and develop responsible behaviors. However, students often face hindrances in accomplishing their developmental tasks. Stress is one of the students' hindrances in fulfilling their developmental tasks (Kholidah & Alsa, 2012).

According to Shagsavarani, Abadi, & Kalkhoran (2015), stress refers to a depressed mental condition. Child Mind Institute (2015) reports that more than 17 million youths suffer from mental illness. One

of the illnesses is stress. Habeeb (2010) reports that the prevalence of university students' stress around the world reaches 71%. This is supported by Fitasari (2011) who reports that 71.6% of Indonesian university students suffer from stress.

University students' stress, according to Maslakhatur (2014), is affected by a range of factors, including the demand to graduate on time, GPA, and organization. Coping strategies are needed to cope with stress. Dexter (2018), Kwaah (2017), Bamuhair (2015) state that to cope with stress, an individual may meditate, listen to music, take exercise, and seek advice from others. These could be done in a special treatment, where the treatment is initiated by collecting

supporting data. Perceived stress scale could be used to collect students' data related to stress (Andreou et al., 2011; Lee 2012; Lesage et al., 2012; Leung et al., 2010; Siqueira et al., 2010). This is an instrument developed by Cohen (1994) to measure one's perceived stress. This scale could be used to collect initial data to design a treatment for solving university students' stress.

Studies on perceived stress scale (PSS) are carried out massively around the world, such as Sun (2019) in China, Manzar (2019) in Ethiopia, Khalili (2017) in Iran, Andreou (2011) in Greece. The result of these studies showed that PSS was valid and reliable to be used in these countries.

The present study aimed to examine the validity and reliability of the perceived stress scale. The analysis result showed that the perceived stress scale was valid and reliable, thus can be used by university counselors to measure students' perceived stress level.

METHODOLOGY

This quantitative study involved 195 students of the Guidance and Counseling Department of a university in Yogyakarta, Indonesia. They were recruited using simple random sampling technique. The data were collected using the Perceived stress scale and analyzed using Rasch model.

RESULT AND DISCUSSION

Based on the description mentioned earlier, we conclude that stress is a general problem occurs among university students, which is affected by various factors. Stress could be handled using treatment. Before conducting any treatment, it is necessary to have an adequate data as the basis so that the treatment given fits the condition. To obtain such data, a valid instrument is required.

Cohen (1994) validate the scale by involving subjects from various racial background, age, and gender. As Cohen (1994) did, the data collected in the present study involved respondents as presented in Table 1.

Table 1
Category and Number of Respondents

Category	Total
Gender	
Male	51
Female	144
Age	
18-22	162
23-27	30
>28	1
Unknown	2
Javanese	163
Sundanese	10
Kebo	1
Jawir	1
Komerling	1
Malay	2
Palembang	1
Sasak	3
Toraja	2
Javanese-Malay	1
Gayo	1
Dayak Taboyan	1
Batak-Malay	1
Bali	2
Hawa	1
Unknown	4

After that, misfit test was conducted to see whether or not the PSS fits to the Rasch model. This scale consisted of 10 items in Table 2.

Table 2
Item Score

No. Item	Value	Description
1	0.95	Fit
2	0.89	Fit
3	1.12	Fit
4	1.07	Fit
5	1.04	Fit
6	0.92	Fit
7	0.90	Fit
8	1.09	Fit
9	1.03	Fit
10	0.92	Fit

As shown in the table above, each item of the PSS fits the Rasch model since the score of each item is higher than 0.5 and less than 1.5.

The score of person reliability was 0.79, which was categorized as good, while the score of item reliability was 0.98, which was

1 categorized as very good. It was found that the person separation score was 1.93 (poor) and the item separation score was 7.17 (very good). The data on person and item criteria are presented in Table 3.

Table 3
Person and Item Criteria

Criteria	Value	Description
Person reliability	0.79	Fair
Item reliability	0.98	Very good
Person Separation	1.93	low
Item Separation	7.17	Very good

Based on the scores of person and item reliability, it could be concluded that the respondents' answer was consistent and the quality of the PSS item was very good. Based on the result of person and item separation, the PSS items lack of variation between the easy and difficult items. However, regarding the item reliability criteria, the respondents recruited had been varied, from low to high perceived stress level. The variance in data explained by measure was 47.5%, while the Unexplained variance data in contrast 1-5 of PCA of residuals was 14.7%. The explained and unexplained variance criteria are described in Table 4.

Table 4
Explained and Unexplained Variance Criteria

Criteria	Value
Variance in data explained by Measures	47.5%
Unexplained variance data in contrast 1-5 PCA of residuals	14.7%

As shown by the analysis result, the PSS contains factors beyond the perceived stress.

Regarding the function of the rating scale (using Winsteps), the result indicated a consistent increase in each item (from -4.09 to 4.03). In other words, the PSS's rating scale functions well. This is supported by equal distribution of the choice in the instrument. Each item serves its functions well.

1 Based on the description above, it could be said that the weakness of Cohen's perceived stress scale lies in its item separation and the presence of factors beyond perceived stress. However, based on the items of the scale, Cohen's PSS fits the Rasch Model.

CONCLUSION

Based on the result of analysis using Winsteps, Perceived Stress Scale was found to fit rasch model. The PSS could be used as a reference for other researchers who develop an instrument related to university students' stress in order to develop a scale with higher person separation value.

REFERENCES

- Andreou, E. Dkk. (2011). Perceived stress scale: reliability and validity study in greece. *International Journal of Environmental Research and Public Health*, 8, 3287-3298.
- Bamuhair, S. S. Dkk. (2015). Souce of stress and coping strategies among undergraduate medical students enrolled in a problem-based learningg curriculum. *Journal of Biomedical Education*, 2015(575139), 1-8.
- _____. (2015). *Children's mental health report*. Child Mind Institute.
- Cohen, S. (1994). *Perceived stress scale*. Mind Garden.
- Danim, S. (2013). *Perkembangan peserta didik*. Bandung: Alfabeta.
- Dexter, L. R. Katie, H. Rudecki, M. & Abraham, S. (2018). College students' stress coping behaviors and perception of stress-effects holistically. *International Journal of Studies in Nursing*, 3(2), 1-6.
- Fisher. W. P. (2007). RASCH measurment. *Rasch Measurement Trasaction*, 21(1), 1087-1096.
- Fitasari, I. N. (2011). *Faktor yang berhubungan dengan kejadian stres pada mahasiswa fakultas kesehatan masyarakat universitas airlangga*. Surabaya: Universitas Airlangga.
- Habeeb, K. A. (2010). Prevalence of stressors among female medical

- student. *Journal of Taibah University Medical Sciences*, 5(2), 110-119.
- Kholidah, E. N. & Alsa. A. (2012). Berfikir positif untuk menurunkan stres psikologis. *Jurnal Psikologi*, 39(1), 67-65.
- Khalili, R. Siranti nir. M. Ebadi, A. Tavalai, A. & Habibi, M. (2017). Validity and reliability of cohen 10-item perceived stress scale in patients with chronic headache: persian version. *Asian Journal of Psychiatry*, 26, 136-140.
- Kwaah, C. Y. (2017). Stress and coping strategies among distance education students at the university of cape coast, ghana. *Turkish Online Journal and Distance Education-TOJDE*, 18(3), 120-134.
- Lee, E. H. (2012). Review of the psychometric evidence of the perceived stress scale. *Asian nursing research*, 6(4), 121-127.
- Lesage, F. X., Berjot, S., & Deschamps, F. (2012). Psychometric properties of the french versions of the perceived stress scale. *International journal of occupational medicine and environmental health*, 25(2), 178-184.
- Leung, D. Y., Lam, T. H., & Chan, S. S. (2010). Three versions of perceived stress scale: validation in a sample of chinese cardiac patients who smoke. *BMC public health*, 10(1), 1-7.
- Manzar, M. D. Dkk. (2019). Psychometric properties of the perceived stress scale in ethiopian university students. *BMC Public Health*, 19(41), 1-8.
- Maslakhatun, U. (2015). *Problem-problem, strategi coping dan resiliensi mahasiswa bidikmisi fakultas ilmu pendidikan universitas negeri yogyakarta*. Yogyakarta: Universitas Negeri Yogyakarta.
- Shahsavarani, M. Abadi, E. A. M. & Kalkhoran, M. H. (2015). Stress: facts and theories through literature review. *International Journal of Medical Reviews*, 2(2), 230-241.
- Siqueira Reis, R., Ferreira Hino, A. A., & Romélio Rodriguez Añez, C. (2010). Perceived stress scale: reliability and validity study in brazil. *Journal of health psychology*, 15(1), 107-114.
- Sun, Y. Gao, L. Kan, Y. & Shi, B. (2018). The perceived stress scale-10 (PSS-10) is reliable and has construct validity in chinese patients with systemic lupus erythematosus. *Sage Journals*, 28(2), 149-155.

PERCEIVED STRESS SCALE DENGAN MODEL RASCH

ORIGINALITY REPORT

98%

SIMILARITY INDEX

98%

INTERNET SOURCES

7%

PUBLICATIONS

7%

STUDENT PAPERS

PRIMARY SOURCES

1

journal.uad.ac.id

Internet Source

98%

Exclude quotes On

Exclude matches Off

Exclude bibliography On