

Hasil Cek_B1

by Universitas Ahmad Dahlan Yogyakarta 23

Submission date: 01-Nov-2023 08:38AM (UTC+0700)


Submission ID: 2213721765

File name: B1._The_Determinants_of_Four_or_More_Antenatal_Care_Vi.pdf (109.1K)

Word count: 3898

Character count: 19776

The Determinants of Four or More Antenatal Care Visits Among Working Women in Indonesia

Hanifa M. Denny, BSPH, MPH, PhD¹ ,
Agung D. Laksono, BSPH, MPH, PhD²,
Ratu Matahari, BSPH, MPH³, and Bina Kurniawan, BSPH, MPH¹

11

Asia Pacific Journal of Public Health

1–6

© 2021 APJPH



Article reuse guidelines:

sagepub.com/journals-permissions

DOI: 10.1177/10105395211051237

journals.sagepub.com/home/aph



Abstract

This study aimed to analyze the determinants of 4 or more antenatal care (ANC) visits among working women in Indonesia. The researchers extracted data from the Indonesian Demographic and Health Survey 2017 and obtained a sample size of 8239 working women aged between 15 and 49 years. Women's residence, age, marital status, education level, parity, economic status, and health insurance were selected as the dependent variables. Binary logistic regression was used for the analysis. Older working women, married working women, educated working women, those in higher economic status, and those with health insurance were more likely to complete four or more of their ANC visits. The more children the working women had, the less likely they would complete their ANC visits. In conclusion, age, marital status, education, parity, economic status, and health insurance are the determinants for completing ANC visits among working women in Indonesia. At the same time, place of residence does not affect the frequency of ANC visits.

Keywords

antenatal care, women's health, parity, ANC, working women

What We Already Know

- The demographic area plays an important factor in contributing to 4 or more antenatal care (ANC) visits.
- Older pregnant women tend to have 4 or more ANC visits.
- The more educated the working woman was, the more likely she was to complete 4 or more ANC visits.

What This Article Adds

- Working women have a potential contribution to the success of 4 or more ANC visits.
- Less-educated working women living in rural areas need more attention to increase the coverage of 4 or more ANC visits.
- The determinants of 4 or more ANC visits among female workers in Indonesia augment the scarcity of information on reproductive health program implementation among working women in Indonesia.

Introduction

Maternal mortality rate (MMR) and infant mortality rate (IMR) are the main maternal and child health indicators. Reduction in maternal and infant deaths is the target of Sustainable Development Goals (SDGs).¹ In 2015, the Indonesian MMR had decreased from 359 deaths per 100 000 live births to 305 per 100 000 live births. However, this condition is still below the target SDGs, 70 deaths per 100 000 live births.² Meanwhile, according to the Indonesian Demographic and Health Survey (IDHS) 2017, the IMR in Indonesia is 15 deaths per 1000 live births.³

¹Faculty of Public Health, Diponegoro University, Semarang, Indonesia

²National Institute of Health Research and Development, The Indonesian Ministry of Health, Jakarta, Indonesia

³Faculty of Public Health, Ahmad Dahlan University, Yogyakarta, Indonesia

Corresponding Author:

Hanifa M. Denny, Faculty of Public Health, Diponegoro University, Jl. Prof. Sudarto, SH, No. 1269, Tembalang, Semarang, Central Java 50275, Indonesia.

Email: hanifadenny@live.undip.ac.id

Antenatal care (ANC) is defined as the access and use of health care during pregnancy. Its objective is to prevent malnutrition among women during pregnancy and reduce low birth weight and infant mortality. Antenatal care is part of the Indonesian programs on decent work to improve women workers' health status during pregnancy.^{4,5} Antenatal care involves health risk diagnoses during pregnancy and childbirth to minimize the adverse effects on women's reproductive health.² The World Health Organization recommends 4 visits during pregnancy to prevent malnutrition in mothers and reduce the risk of low birth weight.⁶ The ANC visit in Ethiopia was negatively affected by lower education, poverty, rural residence, and higher birth order. However, better ANC services contributed to the pregnant women having more visits.⁷

In Indonesia, complete ANC or at least 4 ANC visits during pregnancy are insufficient to reach the country's target.⁸ This phenomenon is similar to that in Pakistan, showing 65.8% to 79.8% access toward 4 ANC visits or ANC completeness.⁹ However, Indonesia has the highest percentage of ANC visits (86%) compared with Cambodia (83%) and Myanmar (47%).¹

Several factors related to ANC completeness include geographical conditions (living in rural/urban areas), sociodemographic conditions (economic status, low education, number of short birth spacing, giving birth without the assistance of health workers or skilled attendants), and cultural values adopted by an individual.^{8,10} A national survey in 2018 showed that the first ANC visit percentage was 96.1% in women aged between 10 and 54 years or increased by 1% compared with the coverage in 2013. Meanwhile, the four ANC visit percentage in 2018 was 74.1% or increased by 4.1% compared with 70% in 2013. Although the percentage of ANC visits has increased, the number is still below the national target of 76%.¹¹

Working women's reproductive health is crucial, as workplace hazards could affect their pregnancy. Some examples of the hazards include fatigue due to excessive working hours or standing for more than 7 hours, stress, and extreme exposure to chemicals, sounds, heat, and biological hazards such as viruses and vibrations that can cause miscarriage or reproductive outcome disorders.¹²

This study aimed to analyze determinants of 4 or more ANC visits among working women in Indonesia. This study is expected to provide some inputs for the upcoming policy and strategy to increase the coverage of 4 or more ANC visits among working women in Indonesia.

Methods

This study used secondary data from the IDHS 2017 as part of the IDHS program conducted by the Inner-City Fund (ICF). The authors obtained an ethical approval from the IDHS data utilization under the following category: Project complies with all of the requirements of 45 CFR 46, "Protection of Human Subjects" from the ICF IRB FWA00000845. Project Title: The

Demographic and Health Survey (DHS) Program (DHS-7), Project Number: 132989.0.000. The IDHS 2017 uses stratification and multistage random sampling methods. The sample consisted of 8239 working women who had a job at the interview time, were between 15 and 49 years old, and had given birth in the last five years.

Procedure

The researchers obtained permission to analyze the IDHS data upon registering the study on the following Web site: <https://dhsprogram.com/data/new-user-registration.cfm>. Subsequently, the respondents' identity was removed from the data set to ensure confidentiality.

Data Analysis

The data set includes information about working women's ANC visits to a health service facility or a midwife. The ANC procedure entails a 1-time visit during the first trimester, a 1-time visit during the 6 months of pregnancy, and 2 visits between 7 and 9 months. Antenatal care is considered complete when a pregnant woman completes 4 visits to a health service facility or a midwife.³ The independent variables for this study were categorical data of the respondents' residence, age, marital status, education level, parity, economic status, and health insurance coverage. A χ^2 test was used to detect the variables related to ANC utilization during pregnancy. Binary logistic regression was applied at the final stage due to the nature of the dependent variable. The SPSS V.22 software was operated in all stages of the statistical analysis.

Results

Working women from urban areas were higher in their 4 ANC visits (Table 1). Working women aged 30 to 34 years had the highest ANC completeness score. Married working women, women with secondary education, women who had 2 to 4 parities, women of higher economic status, and women who had health insurance coverage recorded higher in their 4 or more ANC visits.

Table 2 shows the result of the binary logistic regression of ANC visits among working women in Indonesia. There was no difference in the score of 4 or more ANC visits of women living in rural and urban areas.

Working women aged between 20 and 24 years tended to have 4 or more ANC visits compared with women aged 15 to 19 years. Meanwhile, working women aged 30 to 34 years were more likely to have 4 or more ANC visits than those aged between 15 and 19 years. Subsequently, working women aged 45 to 49 years were more apparent to have 4 or more ANC visits than those aged 15 to 19 years. The tendency to have 4 or more ANC visits increased with the respondents' age (see Table 2).

6
Table 1. Descriptive Statistics of the Respondent of Female Workers in Indonesia (n = 8239).

Variables	Categories	ANC visits		P value
		<4 (n = 923)	≥4 (n = 7316)	
Type of residence	Urban	33.6%	51.4%	<.001*
	Rural	66.4%	48.6%	
Age group, y	15-19	3.7%	1.2%	<.001*
	20-24	13.8%	12.1%	
	25-29	20.4%	25.6%	
	30-34	23.1%	27.3%	
	35-39	22.1%	21.6%	
	40-44	12.9%	10.0%	
Having a husband	No	8.5%	4.0%	<.001*
	Yes	91.5%	96.0%	
Education level	No education	6.9%	1.1%	<.001*
	Primary	40.1%	21.5%	
	Secondary	40.4%	51.0%	
	Higher	12.6%	26.4%	
Parity	<2	21.6%	30.9%	<.001*
	2-4	54.7%	61.9%	
	>4	23.7%	7.2%	
Wealth status:	Poorest	51.4%	22.3%	<.001*
	Poorer	18.7%	17.5%	
	Middle	14.4%	18.2%	
	Richer	8.7%	20.3%	
	Richest	6.8%	21.7%	
Health insurance coverage	No	44.4%	33.7%	<.001*
	Yes	55.6%	66.3%	

Abbreviation: ANC, antenatal care.

*P < .001.

Working women with husbands or partners were inclined to have 4 or more ANC visits than unmarried ones (see Table 2). It indicates that being married or having a partner is a predictive factor for respondents' 4 or more ANC visits.

The analysis showed that working women with primary education were more frequent to have 4 or more ANC visits than those with no educational attainment. Working women with secondary education were more eager to have 4 or more ANC visits than those with no educational attainment. The result also indicated that working women with higher education tended to have 4 or more ANC visits than those with no educational attainment. Thus, this information shows that working women's tendency to have 4 or more ANC visits increased with their education level (see Table 2.).

Working women with parity <2 were more likely to have 4 or more ANC visits than those with parity >4. Furthermore, working women with 2 to 4 parities were more prevalent to have 4 or more ANC visits than those with parity >4. It shows that more children contribute negatively to have 4 or more ANC visits (see Table 2.).

Women whose economic status was in the mediocre category were more likely to complete 4 or more ANC visits than impoverished women. In the meantime, working women with economic status in the middle category tended to complete 4 or more ANC visits than most impoverished women. The most prosperous woman had a higher possibility to complete 4 or more ANC visits than the poorest group. This analysis suggests that the likelihood of 4 or more ANC visits increased with an increase in economic status (see Table 2.).

Working women with health insurance coverage were more positive to have 4 or more ANC visits than those without health insurance (see Table 2). It reveals that health insurance among working women is a predictive factor for having 4 or more ANC visits.

Discussion

We found that older working women are more likely to complete their ANC visits during pregnancy. Age is one of the determinants for completing ANC, as shown in previous studies in Indonesia,¹⁰ Guinea-Bissau, Ethiopia, Ghana, and

6

Table 2. The Result of Binary Logistic Regression of ANC Visits Among Female Workers in Indonesia (n = 8,239).

Variables	Categories	P value	AOR	95% CI	
				Lower bound	Upper bound
Type of residence	Urban	.999	1.000	0.841	1.189
	Rural	—	—	—	—
Age group	15-19	—	—	—	—
	20-24	<.001*	2.533	1.601	4.007
	25-29	<.001*	3.944	2.477	6.278
	30-34	<.001*	4.466	2.765	7.213
	35-39	<.001*	4.830	2.958	7.889
	40-44	<.001*	5.179	3.085	8.692
Having a husband	40-44	<.001*	4.896	2.640	9.081
	No	—	—	—	—
Educational attainment:	Yes	<.001*	2.012	1.524	2.658
	No education	—	—	—	—
	Primary	<.001*	2.371	1.644	3.421
	Secondary	<.001*	3.658	2.513	5.324
	Higher	<.001*	3.147	2.062	4.803
Parity	<2	<.001*	4.789	3.525	6.507
	2-4	<.001*	2.949	2.362	3.682
	>4	—	—	—	—
Wealth status	Poorest	—	—	—	—
	Poorer	<.001*	1.623	1.325	1.988
	Middle	<.001*	2.015	1.603	2.532
	Richer	<.001*	3.324	2.516	4.392
	Richest	<.001*	4.177	3.023	5.771
Health insurance coverage	No	—	—	—	—
	Yes	<.001*	1.436	1.236	1.667

14

Abbreviations: ANC, antenatal care; AOR = adjusted odds ratio; CI = confidence interval.

*P < .001.

several other countries.^{8,13} The age factor is closely related to a woman and her partner's decision related to bearing a child or children, as working women tend to delay their first pregnancy and are more mature in terms of age during pregnancy than those who do not have an occupation.¹⁴

Being married is a predictive factor among working women to complete their ANC visits. Extramarital pregnancy is frowned upon in the Indonesian culture. This social context often leads to unmarried pregnant women hiding their pregnancies, resulting in lower ANC visits.^{10,15,16}

Better education might help working women understand their needs and demands and make them confident about making independent decisions.^{17,18} Education is closely related to power in sexual relationships, especially in a paternalistic social system. Women with better education might have more freedom to decide what is beneficial for their health. In Vietnam, inequality or fewer ANC visits occurred among minority groups, those in poor economic status, and low educational women.^{19,20}

The results show that the higher the parity, the less likely to complete the ANC requirements. Women who have been pregnant or have given birth before tend to neglect their need to seek help from health workers.^{8,14} This result contradicts a study in China that pregnant women tend to have fewer ANC visits during their first and second pregnancies.²¹

The higher the working women's economic status, the higher the possibility of them having ANC visits. A study in India reported that the wealthiest woman was 3.59 times more likely than the most impoverished household to complete all ANC visits.²² Studies reported that women's education and economic status influenced the effectiveness of the ANC visits.^{19,23}

Having health insurance coverage is a positive factor for working women to have 4 or more ANC visits. This result supports the Indonesian health insurance scheme's ultimate goal known as "BPJS" as the Indonesian universal health coverage.²⁴ Studies have confirmed that health insurance positively impacts access to health care services, whether run by the government or the private sector.^{25,26}

Several other studies show that maternal health service providers' factors also contribute to the frequency of ANC visits (eg, pregnant women's response to service innovation and the midwife's attitude as a maternal service provider).²⁷ A possible limitation of this study could be that the data focus only on working women as consumers of maternal health services.

Conclusion

Age group, marital status, education level, parity, economic status, and health insurance coverage play a significant role in having more ANC visits among working women in Indonesia. Those factors are the keys to enhance the future success of ANC visits among working women in Indonesia. However, the place of residence was not determinant of 4 or more visits among working women in Indonesia. The interpretation of this finding seems to be important and informative for readers. The factors found as determinants of 4 or more ANC visits in the analysis of this study would provide a clear direction for the government to modify its acceleration policy to increase the coverage of 4 or more ANC visits among working women in Indonesia. The findings in this study could be useful for other countries in other regions as these results are consistent and confirm previous studies from multiple countries.

Acknowledgments

We express our gratitude to ICF International for granting permission to the author to analyze the IDHS 2017. Diponegoro University will provide a financial award upon publishing this article.

1


Declaration of Conflicting Interests

The author(s) declared no potential conflicts of interest with respect to the research, authorship, and/or publication of this article.

Funding

The author(s) received no financial support for the research, authorship, and/or publication of this article.

ORCID iD

Hanifa M. Denny  <https://orcid.org/0000-0002-9359-3153>

References

- Sebayang SK, Efendi F, Astutik E. Women's empowerment and the use of antenatal care services: analysis of demographic health surveys in five Southeast Asian countries. *Women Health*. 2019;59(10):1155-1171. doi:10.1080/03630242.2019.1593282.
- Efendi F, Chen C-M, Kumiati A, Berliana SM. Determinants of utilization of antenatal care services among adolescent girls and young women in Indonesia. *Women Health*. 2017;57(5):614-629. doi:10.1080/03630242.2016.1181136.
- National Population and Family Planning Board (BKKBN), Statistics Indonesia (BPS), Ministry of Health (Kemenkes), and ICF. Maternal health care. In: *Indonesia Demographic and Health Survey 2017*. Jakarta, Indonesia: BKKBN, BPS, Kemenkes, and ICF; 2018. Accessed September 2, 2020. <https://dhsprogram.com/pubs/pdf/FR342/FR342.pdf>.
- Efendi F, Ni'mah AR, Hadisyatmana S, Kuswanto H, Lindayani L, Berliana SM. Determinants of facility-based childbirth in Indonesia. *Sci World J*. 2019;2019:1-7. doi:10.1155/2019/9694602.
- Titaley CR, Dibley MJ, Roberts CL. Factors associated with underutilization of antenatal care services in Indonesia: results of Indonesia Demographic and Health Survey 2002/2003 and 2007. *BMC Public Health*. 2010;10(1):485. doi:10.1186/1471-2458-10-485.
- Doku DT, Neupane S. Survival analysis of the association between antenatal care attendance and neonatal mortality in 57 low-and middle-income countries. *Int J Epidemiol*. 2017;46(5):1668-1677. doi:10.1093/ije/dyx125.
- Muchie KF. Quality of antenatal care services and completion of four or more antenatal care visits in Ethiopia: a finding based on a demographic and health survey. *BMC Pregnancy Childbirth*. 2017;17(1):300. doi:10.1186/s12884-017-1488-0.
- Jiwani SS, Amouzou-Aguirre A, Carvajal L, et al. Timing and number of antenatal care contacts in low and middle-income countries: analysis in the Countdown to 2030 priority countries. *J Glob Health*. 2020;10(1):010502-010512. doi:10.7189/jogh.10.010502.
- Tarar MA, Khan YN, Ullah MZ, Salik MH, Akhtar S, Sultan T. Knowledge and attitude; pregnancy and antenatal care among young agrarian & non-agrarian females in Faisalabad District, Pakistan. *Pak J Agric Sci*. 2019;56(1):261-273. doi:10.21162/PAKJAS/19.7730.
- Laksono AD, Rukmini R, Wulandari RD. Regional disparities in antenatal care utilization in Indonesia. *PLoS One*. 2020;15(2):e0224006. doi:10.1371/journal.pone.0224006.
- World Health Organization. *State of health inequality: Indonesia*. World Health Organization; 2017. <https://apps.who.int/iris/handle/10665/259685>
- Banerjee B. Physical hazards in employment and pregnancy outcome. *Indian J Community Med*. 2009;34(2):89-93. doi:10.4103/0970-0218.51224.
- Yaya S, Bishwajit G, Gunawardena N. Socioeconomic factors associated with the choice of delivery place among mothers: a population-based cross-sectional study in Guinea-Bissau. *BMJ Glob Health*. 2019;4(2):e001341. doi:10.1136/bmjgh-2018-001341.
- Reynolds CM, McMahon L, O'Malley EG, O'Brien O, Sheehan SR, Turner MJ. Maternal employment and pregnancy outcomes in a large European maternity hospital. *Eur J Obstet Gynecol Reprod Biol*. 2020;250:86-92. doi:10.1016/j.ejogrb.2020.04.005.
- Forbes F, Wynter K, Wade C, Zeleke BM, Fisher J. Male partner attendance at antenatal care and adherence to antenatal care guidelines: secondary analysis of 2011 Ethiopian demographic and health survey data. *BMC Pregnancy Childbirth*. 2018;18(1):145. doi:10.1186/s12884-018-1775-4.
- Wulandari RD, Laksono AD, Nantabah ZK. Effect of marital status on completeness of antenatal care visits among child-bearing age women in rural Indonesia. *Med Leg Update*. 2020;20(4):453-458. Accessed September 4, 2020. <http://www.ijop.net/index.php/mlu/article/download/1858/1628>.

17. Afrizal SH, Hidayanto AN, Handayani PW, et al. Evaluation of integrated antenatal care implementation in primary health care. *J Integr Care*. 2020;99-117. doi:10.1108/JICA-07-2019-0031.
18. Noh J-W, Kim Lee LJ, Akram N, Shahid F, Kwon YD, Stekelenburg J. Factors associated with the use of antenatal care in Sindh province, Pakistan: a population-based study. *PLoS One*. 2019;14(4):e0213987. doi:10.1371/journal.pone.0213987.
19. Kaplowitz ET, Fiori KP, Lauria ME, et al. Sexual relationship power and socio-demographic factors predicting contraceptive use, antenatal visits and sick child health service use in Northern Togo. *Matern Child Health J*. 2020;24(7):845-855. doi:10.1007/s10995-020-02948-w.
20. Kien VD, Jat TR, Phu TV, et al. Trends in socioeconomic inequalities in the use of antenatal care services by women aged 15 to 49 years in Vietnam. *Asia Pac J Public Health*. 2019;31(5):413-421. doi:10.1177/1010539519857305.
21. Jiang K, Liang L, Wang H, et al. Sociodemographic determinants of maternal health service use in rural China: a cross-sectional study. *Health Qual Life Outcomes*. 2020;18(1):1-8. doi:10.1186/s12955-020-01453-6.
22. Ghosh A, Ghosh R. Maternal health care in India: a reflection of 10 years of National Health Mission on the Indian maternal health scenario. *Sex Reprod Healthc*. 2020;25:100530. doi:10.1016/j.srhc.2020.100530.
23. Manyeh AK, Amu A, Williams J, Gyapong M. Factors associated with the timing of antenatal clinic attendance among first-time mothers in rural southern Ghana. *BMC Pregnancy Childbirth*. 2020;20(1):47. doi:10.1186/s12884-020-2738-0.
24. Nasution SK, Mahendradhata Y, Trisnantoro L. Can a national health insurance policy increase equity in the utilization of skilled birth attendants in Indonesia? a secondary analysis of the 2012 to 2016 National Socio-Economic Survey of Indonesia. *Asia Pac J Public Health*. 2020;32(1):19-26. doi:10.1177/1010539519892394.
25. Tilahun H, Atnafu DD, Asrade G, Minyihun A, Alemu YM. Factors for healthcare utilization and effect of mutual health insurance on healthcare utilization in rural communities of South Achefer Woreda, North West, Ethiopia. *Health Econ Rev*. 2018;8(1):1-7. doi:10.1186/s13561-018-0200-z.
26. Wu R, Li N, Erica A. The effects of private health insurance on universal health coverage objectives in China: a systematic literature review. *Int J Environ Res Public Health*. 2020;17(6):20491-20421. doi:10.3390/ijerph17062049.
27. Oscarsson MG, Stevenson-Ågren J. Midwives experiences of caring for immigrant women at antenatal care. *Sex Reprod Healthc*. 2020;24:100505. doi:10.1016/j.srhc.2020.100505.

Hasil Cek_B1

ORIGINALITY REPORT

19%

SIMILARITY INDEX

15%

INTERNET SOURCES

16%

PUBLICATIONS

8%

STUDENT PAPERS

PRIMARY SOURCES

1	cdr.lib.unc.edu Internet Source	2%
2	ecronicon.net Internet Source	2%
3	www.opmcm.gov.np Internet Source	2%
4	childinfo.org Internet Source	1%
5	Agung Dwi Laksono, Rukmini Rukmini, Ratna Dwi Wulandari. "Regional disparities in antenatal care utilization in Indonesia", PLOS ONE, 2020 Publication	1%
6	ijop.net Internet Source	1%
7	Ratna Dwi Wulandari, Agung Dwi Laksono, Nikmatur Rohmah. "Urban-rural disparities of antenatal care in South East Asia: a case study in the Philippines and Indonesia", BMC Public Health, 2021	1%

8	Submitted to Lira University Student Paper	1 %
9	www.sysrevpharm.org Internet Source	1 %
10	eprints.ners.unair.ac.id Internet Source	1 %
11	Submitted to Abdullah Gul University Student Paper	1 %
12	Agung Dwi Laksono, Ratna Dwi Wulandari, Ratu Matahari, Nikmatur Rohmah. "The choice of delivery place in Indonesia: Does home residential status matter?", Heliyon, 2023 Publication	1 %
13	Emmanuel Barankanira, Arnaud Iradukunda, Nestor Ntakaburimvo, Willy Ahishakiye, Jean Claude Nsavyimana, Emmanuel Nene Odjidja. "Spatial distribution and predictive factors of antenatal care in Burundi: A spatial and multilevel baseline analysis for the third burundian demographic and health survey", PLOS ONE, 2023 Publication	1 %
14	bmcpregnancychildbirth.biomedcentral.com Internet Source	1 %

15 Submitted to Badan PPSDM Kesehatan
Kementerian Kesehatan 1 %
Student Paper

16 www.mdpi.com 1 %
Internet Source

17 Submitted to Victoria University 1 %
Student Paper

Exclude quotes On

Exclude matches < 1%

Exclude bibliography On