

Development of Transdermal Patch Preparations Binahong Leaf Extract (*Anredera cordifolia*) (Ten.) Steenis) and Antihyperglycemia Test In Rats

by Galang Pergiawan Ramadhan

Submission date: 27-Mar-2023 08:51PM (UTC+0700)

Submission ID: 2048037246

File name: 24602-68621-2-ED.docx (234.72K)

Word count: 3335

Character count: 18545

17
**Development of Transdermal Patch Preparations Binahong Leaf Extract
(*Anredera cordifolia* (Ten.) Steenis) and Antihyperglycemia
Test In Rats**

Galang Pergiawan Ramadhan¹, Iis Wahyuningsih^{*1}, Wahyu Widyaningsih¹

¹Faculty of Pharmacy, Universitas Ahmad Dahlan, Jl. Prof. Dr. Soepomo, SH, Yogyakarta, Indonesia

Submitted : August 2022 Reviewed : 2022 Accepted 2022

ABSTRACT

Binahong leaves are plants that have great potential as antihyperglycemics, and have chemical content including triterpenoids, flavonoids, and saponins. However, the development of binahong leaves in pharmaceutical preparations is still very lacking. Objectives of this study are (1) knowing the transdermal patch formulation of binahong leaf extract (BLE) produces good physical properties. (2) the development of BLE transdermal patches with different extract concentrations may blood glucose lower (BGL) levels in glucose-induced mice. This study uses a single-layer patch because it has the advantage of being able to penetrate faster and is suitable for compounds that have solubility in polar solvents. The research began with the extraction of binahong leaves, then formulated a transdermal BLE patch using a formula that includes HPMC (polymer), glycerol (plasticizer), oleic acid (enhancer) and tween 80 (enhancer). Diabetic rats were divided into 4 groups of 6 rats, negative control group (glucose + placebo), positive control group (glucose+ insulin), FI group (glucose+ transdermal patch BLE 30%), and FII group (glucose+ transdermal patch BLE 47%). BGL measurements are carried out on the 0th, 3rd, 5th, and 7th days. The physical characterization carried out includes organoleptic, weight uniformity, thickness, pH, folding resistance, and moisture content, obtained FI and FII results that meet the requirements set out by each test standard. BGL from day 1 to day 7 decreased in each group, positive control (68.8±3.76), negative control (112.5±6.89), FI (81.0±4.04), and FII (72.33±5.12). Based on the statistical results of the T-test on the physical evaluation of BLE transdermal patch preparations of FI and FII there were significant differences in weight and thickness uniformity, but there were no significant differences in pH, folding power, and humidity tests. The administration of a 47% BLE transdermal patch provides a faster but insignificant BGL reduction effect compared to the 30% BLE FI transdermal patch.

Keywords: *Anredera cordifolia* (ten). Steenis, patch transdermal, physical characterization, antihyperglycemic.

***Corresponding author:**

Galang Pergiawan Ramadhan

Faculty of Pharmacy

Universitas Ahmad Dahlan

Jl. Prof. Dr. Soepomo, S.H, Warungboto, Umbulharjo, Yogyakarta, Indonesia

Email: iis.wahyuningsih@pharm.uad.ac.id

INTRODUCTION

Hyperglycemia is closely related to diabetes mellitus caused by a lack of insulin hormone, insulin deficiency will cause glucose to accumulate in the blood which then causes blood glucose levels to increase (ADA, 2010).

Binahong leaves (*Anredera cordifolia* (Ten.) Steenis) are a plant that is empirically proven to lower blood glucose levels in diabetic patients (Elisma, 2014). The main ingredients of binahong leaf extract (BLE) that can lower blood glucose levels are triterpenes, flavonoids, alkaloids, tannins, and saponins (Selawa, 2013). The optimal dose of binahong leaf extract as an antidiabetic is 400 mg/KgBW. Binahong leaf ethanol extract gel preparation at a concentration of 10% and 30% can lower blood glucose on the 8th day (Kintoko, 2015). However, the use of natural ingredients of binahong leaves is not accepted by the public, especially oral use so formulations are carried out using transdermal patch preparations. In addition, there have been no studies that have formulated binahong leaf extract into a transdermal patch preparation.

Transdermal patch preparations are preparations that are placed or attached to the skin to provide a certain dose of the drug and enter the bloodstream (Mane, 2021). Transdermal preparations can improve the administration of drugs through the skin for topical and systemic administration. Patch preparations are very suitable in hyperglycemic patients because the drugs given do not undergo first-pass metabolism so that drug absorption is more optimal, can be used long-term therapy or follow-up treatment, and is easily stopped in case of poisoning/allergies. The frequency of administration can be reduced to improve patient adherence (Mane, 2021). The purpose of this study is to know if the formulation of BLE patches meets the physical characteristics of good patches and to know if the difference in BLE concentrations in transdermal patch preparations can affect BGL-lowering activity.

MATERIALS AND METHOD

Materials

Fresh binahong leaves come from Allizzu herbal garden, Bantul district, Yogyakarta, aluminum foil, aquadest (technical), oleic acid (pharmaceuticals), oral disposable, plaster (medical), ethanol 70% (technical), ethanol 96% (technical), glycerol (pharmaceuticals), glucose (pharmaceuticals), HPMC (pharmaceuticals), insulin (pharmaceuticals), standard feed (hi-pro-vit), plastic wrap, tissue, tween 80 (pharmaceuticals), water for injection (technical).

Transdermal patch making

The transdermal patch formula of binahong leaf extract tested for antihyperglycemic activity can be seen in table I.

Table I. Transdermal patch formula of binahong leaf extract.

No	Material	Concentration	
1	Binahong leaf extract	5%	47%
2	HPMC (<i>Stabilizer</i>)	0.2g	0.2g
3	Glycerol (<i>Plasticizer</i>)	0.2g	0.2g
4	Oleic Acid (<i>Enhancer</i>)	0.1g	0.1g
5	Tween 80 (<i>surfaktan</i>)	0.06 g	0.06 g

BLE transdermal patches are created by mixing HPMC into glycerol as a mixture of 1. Oleic acid was mixed with tween 80 and then dissolved in 5 mL as a mixture of 2. BLE has dissolved in alcohol as much as 5 mL then added mixture 1 then mixture 2 and stirred until homogeneous, the mixture is poured into a mold then dried at room temperature for 30 minutes to evaporate the alcohol and ventilated at 50°C for 24 hours so

that a layer of film is obtained then wrapped in aluminium foil and then stored in a desiccator.

Test Characteristics of The Preparation

a. Organoleptic Test

Testing was carried out visually in the form of color, texture, and aroma (Kumar *et al.*, 2013).

b. Weight Uniformity Test

The five patch matrices of each formulation are weighed and averaged, and then calculated the standard deviation and relative standard deviation (Kumar *et al.*, 2013).

c. Patch thickness test

The five patch matrices of each formulation are measured in thickness using calipers. Calculated average patch matrix thickness, standard deviation and relative standard deviation (Kumar *et al.*, 2013).

d. pH Test

The five patch matrices of each formulation are tested using a special pH meter (pH meter surface) tool on the surface. Calculated the average value of the pH obtained (Kumar *et al.*, 2013).

e. Fold Durability Test (*Folding Endurance*)

It is done by means of one patch of each formulation folded in the same place several times until it looks visibly torn. The number of times a patch can be folded in the same place without tearing occurring is calculated as the value of the durability of the fold (Kumar *et al.*, 2013).

f. Moisture Absorption Test

The five patch matrices of each formulation are weighed and placed in a desiccator containing silica gel in them at room temperature for 24 hours. The patch matrix is then weighed and calculated the percentage of moisture absorption (Kumar *et al.*, 2013).

BGL Drop Activity Test

The number of experimental animals is based on the calculation formula of freed (n-1) (t-1) ≥ 15 . The test animals used in this study were 24 male white rats (*Rattus norvegicus*) weighing 200-300 grams which were divided into 4 groups, namely:

- Negative control group (Glucose + placebo)
- Positive control group (glucose + insulin)
- FI group (glucose + Patch BLE 30%)
- FII group (glucose + Patch BLE 47%)

The test animal was acclimatized for ± 7 days. Each group of test animals was satisfied for 10-18 hours and then measured the initial blood glucose level using a glucometer. Glucose administration is done orally for 8 days at a dose of 4.5 mg / kgBW. Mice are said to be diabetic if the blood glucose level is >124 mg/dL using a glucometer autocheck device (Rohma *et al.*, 2015). Then the rat's hair is shaved to the back with a size of 3x5 cm as a place for applying transdermal patch preparations. Each rat measured its blood glucose levels after 6 hours of treatment on days 1, 3, 5 and 7 (Rohma *et al.*, 2015).

Data Analysis

The data obtained in the form of patch physical characteristics using the t-test and the time of decrease in BGL of male white rats (*Rattus norvegicus*) was analyzed statistically using the one-way ANOVA test with a 95% confidence level. Sig. <0.05 Signification is established if $p < 0.05$.

Judul manuskrip (Penulis pertama)

RESULTS AND DISCUSSION

Characterization Of Transdermal Patches

The formulated BLEtransdermal patch is followed by physical tests including organoleptic tests, weight uniformity tests, patch thickness tests, ph tests, folding endurance tests, and moisture content tests.

1. Organoleptic

In organoleptic tests of transdermal patch preparations of binahong leaf extract include aroma, color, and texture. The organoleptic test results of each extract patch formula appear in figure 1.




Figure 1.(a) FI patch BLE 30%; (b) FI patch BLE47%

Based on the organoleptic test above FI (binahong extract 30%) and FII (binahong extract 500 g) have the same characteristics of odor (typical of extract) and texture (smooth, homogeneous, elastic surface), the difference only occurs in the color of the preparation where FI the amount of binahong extract is 30% brighter than FII with binahong extract 47%. This happens because binahong extract has a natural green-black color, the higher the amount of extract used to eat the darker the color of the preparation follows the natural color of the extract.

2. Uniformity of Weights

Measurements were made using 10 replications in each formula. The weight of each patch should not deviate from the standard deviation of <0.05 (Yadav, 2011). As seen in table II.

Table II. Uniformity of Weights of Transdermal Patch Preparations BLE

Replication	FI (30%)	FII (47%)
1	0.853	1.112
2	0.826	1.119
3	0.834	1.073
4	0.848	1.086
5	0.856	1.141
6	0.822	1.091
7	0.863	1.128
8	0.843	1.107
9	0.821	1.103
10	0.877	1.098
Average	0.8443	1.1058
SD	± 0.018	± 0.020
CV (%)	$\pm 2.20\%$	$\pm 1.83\%$

Significant, namely FI (SD ± 0.018 , CV $\pm 2.20\%$) and FII (SD ± 0.020 , CV $\pm 1.83\%$) are still in the stipulated provisions, namely the value of SD < 0.05 and CV $< 5\%$ (Baharudin & Maesaroh,

2020). This difference in patch weight affects the amount of active substance used so it affects the uniformity of the content. Thickness test.

A good patch thickness size is in the range of 0.5– 1.0 mm, if it is smaller it will make it difficult to use (Nuryanti et al., 2016), if the patch is too thick, it will be difficult to release the active substance from the patch (Shirsand et al., 2012). Testing the thickness of the patches in each formula is to measure the thickness one by one 3 replication patches (Nurahmanto, 2016), Patch thickness is measured at 5 different points using Vernier calipers (Baharudin & Maesaroh, 2020). The requirement for a good range of durable patches is CV <2% (Nurahmanto, 2016) and an elementary < score of 0.05 (Baharudin & Maesaroh, 2020). As shown in table III.

Table III. Thickness of Transdermal Patch Preparation BLE

Replication (mm)	FI (mm)	FII (mm)
1	0.52	0.92
2	0.54	0.94
3	0.52	0.98
Average	0.526	0.946
SD	±0.005	±0.14
CV (%)	±0.014	±1.52%

The value obtained from the patch thickness test shows that the patch results made met the patch thickness requirements.

3. pH

pH testing determines whether the transdermal patch preparation has a pH of the preparation used for the skin, which is 4.5-6.5. If the pH of the patch is too acidic and too alkaline it can cause the skin to become dry (Mappa et al., 2013). As shown in table IV.

Table IV. pH Transdermal Patch Preparation BLE

Formula	Replication (%)			Average (%)	SD±CV (%)
	1	2	3		
FI	6.1	6.4	6.2	6.23	0.152±2.45
FII	6.7	6.4	6.4	6.53	0.173±2.66

The test results of the two formulas are known to meet the pH requirements of transdermal preparations, namely 4.5-6.5 (Mappa et al., 2013).

4. Folding Endurance Test

This approval aims to see if the patches made have resistance to folds. Patches are said to meet the criteria when they withstand folds more than 300 times (Lakhani et al., 2015). As seen in table V.

Table V. Folding Power of Transdermal Patch Preparations BLE

Formula	Replication (%)			Rata-rata (%)	SD±CV (%)
	1	2	3		
FI	357	332	378	355.66	23.02±6.47
FII	322	346	362	343.33	20.13±5.86

It can be seen table VI above shows that the two patch formulations made are qualified for good patch fold resistance, which is > 300 times (Ningsi, 2015).

5. Moisture Absorption Test

The prepared patches are each weighed (initial weight) and stored in a desiccator containing silica gel at room temperature for 24 hours. The patch is then re-weighed (final weight) (Shabbir et al., 2017).

The required moisture content range is 1 - 10% (Kumar et al., 2013) in table VI.

Table VI. Moisture Preparations Transdermal Patch BLE

Formula	Replication(%)			Average (%) (x ±SD)
	1	2	3	
Formula I	2.5	2.78	2.10	2.54±2.5385
Formula II	2.9	3.24	2.53	2.85±0.3570

The results of testing percent moisture content show that formula I and Formula II have values that already meet the required range of 1 - 10% (Kumar et al., 2013).

Test The Activity of Lowering Blood Glucose Levels (BGL)

1. Hyperglycemia Rats

After 8 days of the administration, blood glucose measurements are carried out using an autocheck glucometer tool. Rats were said to be hyperglycemia if blood glucose levels exceeded BGL values >135 mg/dL or >124 mg/dL based on the maximum value of the glucometer autocheck tool (Wolfensohn, 2013). The results of 0-8 days of administration obtained seen in table VII

Table VII. BGL Data Day 0 - 8 Days Of Oral Glucose Induction

Formula	Blood glucose levels (mg/dL)	
	Day 0 (x ± SD)	Day 8 (x ± SD)
Positive control (insulin)	100.5 ± 8.09	171.83 ± 43.3
Negative control (placebo)	94.7 ± 14.13	163.33 ± 11.21
Trandermal patch extract (30%)	103.7 ± 4.41	166.33 ± 44.2
Trandermal patch extract (47%)	88.2 ± 3.65	162.50 ± 38.6

From table VII obtained after oral administration of 4.5 g / kgBB glucose, the blood glucose levels of rats in each group increased in value, namely ≥ 124 mg / dL on the 8th day. Oral administration of glucose for 8 days makes blood glucose levels rise, Insulin hormone in charge of introducing glucose into cells cannot work optimally because the amount of glucose is too much. So that there is an increase in glucose levels in the blood (Wolfensohn, 2013).

2. Binahong Leaf Transdermal Patch Preparation

Transdermal patching of binahong leaves is carried out once every 2 days. It is based on the function of the patch preparation which is to reduce the frequency of use and the ability of the HPMC polymer, namely the formula with a single polymer HPMC able to release excellent drugs, namely 99.626% in 48 hours (Mangilal et al., 2015).

Treatment for 7 days can be concluded for positive control to provide a significant reduction of 72.33 mg/dL FI 72.33 mg/dL and FII 68.8mg/dL. From the data above, it can be seen that the difference in the concentration of BLE extract used can provide a different decrease in BLS. The greater the concentration, the higher the percentage decrease in BLS. Although the FI gives a slightly lower decrease than the FII the decrease that occurs is within the normal limits of BLS. This signals FI is effective for lowering BGL in mice. Meanwhile, the negative control decrease was 112.5mg/dL. The low decrease in BGL in negative control compared to other groups is caused by the absence of substances that can help reduce BGL. However, there is still a decrease in BLS in the negative control, this is likely to occur due to the role of the body's metabolism from mice that can reduce BGL by converting it into energy (Supriyanto *et al.*, 2022).

Fig 1. Decreased BGL of Hyperglycemia Rats After Administration of Transdermal Patches of Bianahong Leaf Extract




Fig 1. Graph of decrease in blood glucose levels against time (day)

CONCLUSION

Results of **test** statistics on the physical evaluation of BLE transdermal patch preparations of 30% and 47% **there was a significant difference in** uniformity of weight and thickness, but there were no significant differences in the pH, folding power, and humidity tests. The administration of a 47% BLE transdermal patch gave a faster but not significant BGL reduction effect compared to a 30% BLE transdermal patch.

ACKNOWLEDGEMENT

Thanks to Mrs. Dr. apt. Iis Wahyuningsih, M.Si. who has guided and entrusted this research to me.

LIST OF REFERENCES

- Amjad, Ehteshamuddin, S. Chand, Hanifa, M. Sabreesh, and G. S Kumar. (2011). Formulation and evaluation of niosomes. *Indian Journal of Pharmaceutical Sciences*, 73(3), 323–328. <https://doi.org/10.4103/0250-474X.93515>
- Baharudin, A., & Maesaroh, I. (2020). Formulasi Sediaan Patch Transdermal dari Ekstrak Bonggol Pohon Pisang Ambon (*Musa paradisiaca* var. *sapientum*) Untuk Penyembuhan Luka Sayat. *Herbapharma: Journal of Herb Pharmacological*, 2(2), 55–62. <http://ojs.stikes-muhammadiyahku.ac.id/index.php/herbapharma/article/view/149>
- Elisma. (2014). Prosiding Seminar Nasional dan Workshop “Perkembangan Terkini Sains Farmasi dan Klinik IV” tahun 2014. 220–227.
- Lucia Hendriati, Angelica Kresnamurti, Teguh Widodo, Elizabeth Ivani, Merry Yulia. (2015). Efek hipoglikemik dan uji hipersensitivitas sediaan transdermal ekstrak angsa (*Pterocarpus indicus* Wild). *Jurnal Farmasi Sains Dan Terapan*, 2(2), 15–19.
- Hu, X., Wang, S., Xu, J., Wang, D. B., Chen, Y., & Yang, G. Z. (2014). Triterpenoid saponins from *Stauntonia chinensis* ameliorate insulin resistance via the AMP-activated protein kinase and IR/IRS-1/PI3K/Akt pathways in insulin-resistant HepG2 cells. *International Journal of Molecular Sciences*, 15(6), 10446–10458. <https://doi.org/10.3390/ijms150610446>
- Kintoko. (2015). The effectivity of ethanolic extract of binahong leaves (*Anredera cordifolia* (tenore) steen) gel in the management of diabetic wound healing in aloxan-induced rat

Judul manuskrip (Penulis pertama)

- models. *Jurnal Kedokteran Dan Kesehatan Indonesia*, 7(2), 52–58.
- K.P.Sampath Kumar, Debjit Bhowmik AndChiranjib.B,R.M.Chandira. (2013). Formulation and evaluation of transdermal patch of stavudine. *Dhaka University Journal of Pharmaceutical Sciences*, 12(1), 63–69. <https://doi.org/10.3329/dujps.v12i1.16302>
- Lakhani, P., Bahl, R., & Bafna, P. (2015). Transdermal Patches: Physiochemical and in-Vitro Evaluation Methods. 6(5). [https://doi.org/10.13040/IJPSR.0975-8232.6\(5\)](https://doi.org/10.13040/IJPSR.0975-8232.6(5)).
- Mangilal, T., Patnaik, K. S. K. R., Kumar, M. S., Thirupathi, L., Nagaraju, I., & Kumar, P. V. (2015). Formulation and evaluation of nitrendipine loaded transdermal patch using HPMC K-15 and Eudragit RS100 polymers. *International Journal of Pharma Research and Health Sciences*, 3(3), 708–719. http://www.pharmahealthsciences.net/pdfs/volume3-issue3/6_Vol3_Issue3_MS_15177.pdf
- Mappa, T., Edy, H. J., & Kojong, N. (2013). Formulasi Gel Ekstrak Daun Sasaladahan (*Peperomia pellucida* (L.) H.B.K) Dan Uji Efektivitasnya Terhadap Luka Bakar Pada Kelinci (*Oryctolagus cuniculus*). *Pharmacon*, 2(2), 49–56.
- Nasution, R. (2020). Uji Aktifitas Ekstrak Daun Binahong (*Anredera Cordifolia* (Ten .) Steenis) Terhadap Kadar Gula Darah Mencit Rani Ardiani * Haris Munandar Nasution Faisal Amin Tanjung * Fakultas Farmasi Universitas Muslim Nusantara (UMN) Al-Washliyah. *Jurnal Ilmiah Farmasi*, 484–490.
- Surya Ningsi, Naswina Putrianti, Dwi Wahyuni. (2015). Formulasi, Karakterisasi Dan Uji Penetrasi In Vitro Patch Ekstrak Biji Kopi Robusta (*Coffea Canephora*) Sebagai Sediaan Anti Selulit. *Jurusan Farmasi FIK Universitas Islam Negeri Alauddin Makassar*. 2(3), 6–8.
- Nurahmanto, D. (2016). Pengaruh Perbedaan Chemical Penetration Enhancer Pada Penetrasi Transdermal Patch Prometazin HCl. *Jurnal Ilmiah Manuntung*, 2(2), 208. <https://doi.org/10.51352/jim.v2i2.68>
- Nuryanti, Nugroho, A. K., and Martien, R. (2016). The Effect of Propylene glycol , Oleic Acid and Isopropilalcohol To The Transdermal Patch Losartan Potassium Formula. *Acta Pharmaciae Indonesia*, 4(1), 7–14.
- Rohma, S. C., Ulfa, E. U., & Holiday, D. (2015). Pengaruh Gel Binahong (*Anredera cordifolia* (Ten.) Steenis) terhadap Penyembuhan Luka Tikus Diabetes yang Diinduksi Aloksan. *E-Jurnal Pustaka Kesehatan*, 3(3), 414–418.
- Selawa, W. (2013). Antioksidan, Kandungan Flavonoid Dan Kapasitas [Anredera, Total Ekstrak Etanol Daun Binahong(*Anredera cordifolia* (Ten.) Steenis)]. *Jurnal Ilmiah Farmasi*.
- Shabbir, M., Ali, S., Raza, M., Sharif, A., Akhtar, M. F., Manan, A., Fazli, A. R., Younas, N., & Manzoor, I. (2017). Effect of hydrophilic and hydrophobic polymer on in vitro dissolution and permeation of bisoprolol fumarate through transdermal patch. *Acta Poloniae Pharmaceutica - Drug Research*, 74(1), 187–197.
- Shirsand, S., Suresh, S., Swamy, P., Keshavshetti, G., & Reddy, P. P. (2012). Formulation and optimization of mucoadhesive bilayer buccal tablets of atenolol using simplex design method. *International Journal of Pharmaceutical Investigation*, 2(1), 34. <https://doi.org/10.4103/2230-973x.96924>
- Supriyanto, H., Vellyana, D., & Stiawan, D. (2022). Pengaruh Aktivitas Fisik Jalan Kaki Terhadap Gula Darah Sewaktu Penderita Diabetes Melitus Tipe 2 Di Wilayah Kerja Puskesmas Kotadalam Pesawaran Tahun 2021. *Healthcare Nursing Journal*, 4(1), 194–205. <https://doi.org/10.35568/healthcare.v4i1.1844>
- Wolfensohn, S. (2013). *Handbook of Laboratory Animal Management and Welfare*, 4th Edition. <https://www.wiley.com/en-au/9780470655498>
- Yadav and Jain DK. (2011). Preparation And Characterization Of Microparticulate System Of Propranolol Hydrochloride, *JDDT. All Rights Reserved*. 1(2), 15–20.

Development of Transdermal Patch Preparations Binahong Leaf Extract (*Anredera cordifolia*) (Ten.) Steenis) and Antihyperglycemia Test In Rats

ORIGINALITY REPORT

10%

SIMILARITY INDEX

10%

INTERNET SOURCES

3%

PUBLICATIONS

5%

STUDENT PAPERS

PRIMARY SOURCES

1	eprints.uad.ac.id Internet Source	3%
2	Submitted to Universitas Sanata Dharma Student Paper	1%
3	Submitted to Surabaya University Student Paper	1%
4	journal.uad.ac.id Internet Source	1%
5	www.ciser.com.br Internet Source	<1%
6	e-prosiding.umnaw.ac.id Internet Source	<1%
7	Submitted to Jawaharlal Nehru University (JNU) Student Paper	<1%
8	vs1.doczz.cz Internet Source	<1%

9	www.scribd.com Internet Source	<1 %
10	bbrc.in Internet Source	<1 %
11	journal.uny.ac.id Internet Source	<1 %
12	www.science.gov Internet Source	<1 %
13	innovareacademics.in Internet Source	<1 %
14	www.scielo.br Internet Source	<1 %
15	"Taurine 11", Springer Science and Business Media LLC, 2019 Publication	<1 %
16	Hans-Peter Volz, Arthur Mackert, Rolf-Dieter Stieglitz, Bruno Müller-Oerlinghausen. "Diurnal Variations of Mood and Sleep Disturbances during Phototherapy in Major Depressive Disorder", Psychopathology, 1991 Publication	<1 %
17	jddtonline.info Internet Source	<1 %
18	www.journal.uad.ac.id Internet Source	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On