

Phytochemical Screening Analysis and Determination of Total Flavonoids and Total Phenolics Content of Ethanol Extract of Sungkai Leaf (*Penorema canescens* Jack) from Samarinda City

by Universitas Ahmad Dahlan Yogyakarta 24

Submission date: 04-Jan-2024 08:40AM (UTC+0700)

Submission ID: 2266554798

File name: phytochemical.pdf (874.65K)

Word count: 5603

Character count: 30697

DOI : 10.35311/jmpi.v9i2.360

Phytochemical Screening Analysis and Determination of Total Flavonoids and Total Phenolics Content of Ethanol Extract of Sungkai Leaf (*Penorema canescens* Jack) from Samarinda City

Muhammad Ma'ruf²⁵, Moch. Saiful Bachri²⁵, Laela Hayu Nurani²⁵

Faculty of Pharmacy, Universitas Ahmad Dahlan Yogyakarta, 55164, Indonesia

Citation: Ma'ruf, M., Bachri, M. S., & Nurani, L. H. (2023). Phytochemical Screening Analysis and Determination of Total Flavonoids and Total Phenolics Content of Ethanol Extract of Sungkai Leaf (*Penorema canescens* Jack) from Samarinda City. *Jurnal Mandala Pharmacon Indonesia*, 9(2), 262-272. <https://doi.org/10.35311/jmpi.v9i2.360>

Submitted: 16 July 2023

Accepted: 12 October 2023

Published: 23 December 2023

*Corresponding Author:

Muhammad Ma'ruf

Email:

muhammadmaruf2400@gmail.com



Jurnal Mandala Pharmacon Indonesia is licensed under a Creative Commons Attribution 4.0 International License

ABSTRACT

Sungkai is a plant that contains compounds such as flavonoids, tannins, phenolics, saponins, steroids, and terpenoids, which have the potential to develop native Indonesian herbal medicines to treat the prevention and treatment of various diseases. This research aimed to analyze phytochemical screening and the total flavonoid and phenolic content in sungkai leaves originating from Samarinda City. This research method was carried out by extracting sungkai leaves with 96% ethanol and evaporating them with a vacuum rotary evaporator. Sungkai leaf extract was analyzed for phytochemical screening using the tube method with color testing from various reagents and total phenolic and flavonoid content testing using the colorimetric method using a UV-Vis spectrophotometer. The results of the phytochemical screening showed that the ethanol extract of sungkai leaves from Samarinda City were positive for containing flavonoids, saponins, alkaloids, phenolics, and steroids. The results of determining the flavonoid and phenolic content of sungkai leaf ethanol extract were $18,210 \pm 0.271$ mg QE/g sample and $33,666 \pm 1.052$ mg GAE/g sample.

Keywords: Sungkai, Screening, Phytochemical, Flavonoid, Phenolic

ABSTRAK

Sungkai merupakan salah satu tumbuhan yang memiliki kandungan senyawa seperti flavonoid, tanin, fenolik, saponin, steroid, dan terpenoid yang memiliki potensi sebagai pengembang obat herbal asli Indonesia untuk menangani pencegahan dan pengobatan dari berbagai penyakit. Tujuan penelitian ini adalah untuk menganalisis skrining kimia serta analisis kandungan flavonoid dan fenolik total pada daun sungkai yang berasal dari Kota Samarinda. Metode penelitian ini dilakukan dengan mengekstrak daun sungkai dengan etanol 96% dan diuapkan dengan *vacuum rotary evaporator*. Ekstrak daun sungkai dianalisis skrining fitokimia menggunakan metode tabung dengan uji warna dari berbagai pereaksi dan uji kandungan fenolik total dan flavonoid total dengan metode kolorimetri menggunakan spektrofotometer UV-Vis. Hasil dari skrining fitokimia diperoleh bahwa ekstrak etanol daun sungkai yang berasal dari Kota Samarinda positif mengandung flavonoid, saponin, alkaloid, fenolik, dan steroid. Dan hasil penetapan kadar flavonoid total serta fenolik total ekstrak etanol daun sungkai berturut-turut sebesar $18,210 \pm 0,271$ mg QE/g sampel dan $33,666 \pm 1,052$ mg GAE/g sampel.

Kata Kunci: Sungkai, Skrining, Fitokimia, Flavonoid, Fenolik

INTRODUCTION

Indonesia is an archipelagic country which has various kinds of natural resources. One of the wealth owned by Indonesia is its flora and fauna. The wealth of flora owned by Indonesia is included in the category of medicinal plants. Indonesia has about 30,000 types of plants and 7,000 other types of plants

that can be used as medicinal plants (Jumi & Komalasari, 2017). One example of a plant that can be used as traditional medicine and whose efficacy has not been scientifically proven is sungkai (*Peronema canescens* Jack).

According to Adlis (2020), Sungkai leaves obtained from Padang Pariaman Regency with hexane, ethyl acetate, and

ethanol extracts of Sungkai leaves contain secondary metabolites, namely flavonoids, phenolics, saponins, steroids, and alkaloids. Plants that contain flavonoids can inhibit the activity of angiotensin-converting enzymes, reduce oxidative stress, and increase the relaxation of the blood vessel endothelium, which decreases blood pressure (Widiasari, 2018). A study conducted by Carolina et al., (2022), to giving sungkai leaf infusion to older adults in the Pahandut Palangka Raya Health Center area resulted in changes in blood pressure that reduced blood pressure in hypertensive older adults. So that Sungkai leaves can be efficacious in the treatment of hypertension.

Phytochemical screening is one way to identify a plant's secondary metabolite content. Phytochemical screening using the tube method was chosen because phytochemicals can determine the properties of active substances that cause toxic or beneficial effects on crude extracts if tested in a biological system (Shofi et al., 2020). Therefore, it is necessary to carry out further research on the ethanol extract of Sungkai leaves from Palaran District, Samarinda City, East Kalimantan, to determine the secondary metabolites contained in the form of flavonoids, saponins, alkaloids, phenolics, and steroids and to analysis of the total flavonoid and total phenolic content

METHOD

Tools

The tools used in this research were a vacuum rotary evaporator (Heidolph), glassware (Iwaki), analytical scales (Ohaus), micropipette (Socorex), UV-VIS spectrophotometer (Shimadzu UV-1800).

Materials

The materials used in this study were sungkai leaves (*Peronema canescens* Jack) from Palaran District, Samarinda City, East Kalimantan, Dragendorff peraction, Mayer peraction, solution of iron (III) chloride solution of 10%, ethanol 96%, quercetin, gallic acid, NaOH, and AlCl₃.

Identification of Sungkai Plants

The sample used in this study was carried out. Identification of plants was carried out to establish the truth of the material under investigation. Identification of Sungkai leaf plants is carried out at the UPT. Herbal Materia Medica Laboratory, Batu City, East Java, with letter number: 067/1382/102.20/2023. Sungkai leaves used are the older parts of the leaves and get enough sunlight. After washing with running water, the leaves are dried and protected from direct sunlight. The signs that the sungkai leaves are ready to be mashed have shown that if they are squeezed, they will break easily. Furthermore, sungkai leaves were mashed and sieved using sieve number 60.

Sample Preparation of Sungkai Leaf Ethanol Extract

The preparation of this extract follows the method used by Sadik (2021), macerating 1 kg of dried Sungkai leaf simplicia with 96% ethanol solvent with a ratio of 1:5 for 3 days. Then, the maceration results were evaporated using a vacuum rotary evaporator to obtain a thick ethanol extract of Sungkai leaves.

Phytochemical Screening Analysis

In the phytochemical screening analysis of the ethanol extract of sungkai leaves, tests were carried out, which included reaction tests on color changes and reaction tests on precipitation as follows:

1. Flavonoid Test

The Flavonoid test was carried out with Mg and 2N HCl reagents. A positive test for flavonoids is indicated by reddish-black, yellow, or orange precipitates (Febri Fatwami et al., 2023).

2. Saponin Test

The Saponin test was carried out with hot distilled water and 2N HCl. Stable foam indicates a positive saponin test, and the foam does not disappear when added with HCl 2 N (Larasati & Putri, 2023).

3. Alkaloid Test

The alkaloid test was carried out with Dragendorff reagent. A positive test for alkaloids is indicated by the presence of an

orange-to-yellow precipitate (Imran et al., 2023).

4. Phenolic Test

The phenolic test was carried out with a FeCl₃ reagent. A positive test for phenolic is indicated by the formation of a blue or dark blue-green color (Rifkia & Prabowo, 2020).

5. Steroid Test

The steroid test was carried out by reacting anhydrous acetic and sulfuric acid. A positive steroid test is indicated by the formation of a bluish-green color (Silvani et al., 2023).

Determination of Total Flavonoid Content by UV-Vis Spectrophotometry

In conducting a quantitative analysis of the total flavonoid content, follow the method from Ramadhani et al., (2022) with a few modifications. The operating time was determined with quercetin mother liquor for 60 minutes at a theoretical maximum wavelength of 415 nm. Then, the maximum wavelength of quercetin was determined and incubated according to the operating time obtained and then read at a wavelength of 350-500 nm. Comparison solutions were prepared with concentrations of 125, 100, 75, 50, and 25 µg/ml incubated according to the operating time and measured at the maximum wavelength according to the results of the previous measurements. Weigh 0.1 gram of ethanol extract from Sungkai leaves and then dissolve it with ethanol until the volume is 10 mL. Take 0.5 mL of the previous solution, then add 1.5 ml of ethanol, 0.1 mL of 10% AlCl₃, 0.1 ml of 1M sodium acetate, and 2.8 ml of distilled water. Then, it was incubated according to the operating time, and the absorbance was measured at the maximum wavelength of the previous measurement results. The sample absorbance measurement procedure was repeated 5 times to minimize errors, and the results obtained were expressed in mg QE/g simplicia.

Determination of Total Phenolic Content by UV-Vis Spectrophotometry

In conducting a quantitative analysis of total phenolic content, follow the method

from Ramadhani et al., (2022) with a few modifications. The operating time was determined with gallic acid mother liquor and then read at a wavelength of 730 nm every minute until the 90th minute. Then, the gallic acid wavelength was determined. Maximum wavelength measurements were made using visible spectrophotometry at 550-800 nm. Then, a series of concentration solutions were created with concentrations of 30, 40, 50, 60, 70, and 80 µg/ml and incubated according to the operating time that had been obtained. The absorbance was measured at the maximum wavelength that had been obtained previously. Weigh 0.2 grams of ethanol extract from Sungkai leaves and dissolve it with methanol p.a until the volume is 25 mL. Take 1 mL of the once-diluted solution and add 5 mL of Folin-Ciocalteu solution. And they added 4.0 mL of NaOH. Then, it was incubated according to the operating time, and the absorbance was measured at the maximum wavelength of the previous measurement results. The sample absorbance measurement procedure was repeated 5 times to minimize errors, and the results obtained were expressed in mg QE/g simplicia.

RESULTS AND DISCUSSION

Extraction of Sungkai Leaf Ethanol Extract

The yield of macerate obtained from the evaporation of the solvent using a vacuum rotary evaporator is 125 grams of thick extract. A vacuum rotary evaporator is used because it has the principle of evaporating solvent below its boiling point by lowering the pressure using a vacuum. The lower the pressure given, the faster the evaporation of the solvent obtained so that the sungkai viscous extract obtained was as much as 125 grams compared to the weight of the simplicial powder obtained.

$$\begin{aligned} \text{\% yield} &= \frac{\text{weight of viscous extract obtained}}{\text{weight of simplicia}} \times 100\% \\ &= \frac{125 \text{ grams}}{1000 \text{ grams}} \times 100\% \\ &= 12,5\% \end{aligned}$$

The percentage yield of the ethanol extract of Sungkai leaves obtained was higher than that obtained in a study conducted by (Fadlilaturrahmah et al., 2021), which was 7.28%. This happens because it can be influenced by the location where the Sungkai plant grows. The percentage yield calculation is carried out to compare the final product obtained from the extraction process to the raw materials used before the extraction process so that the amount of raw materials needed to get the desired extract can be

determined. The higher the yield percentage obtained, the more dissolved substances during the extraction process.

Phytochemical Screening Analysis

The results of the phytochemical screening analysis of the ethanol extract of Sungkai leaves (*Peronema canescens* Jack) originating from Samarinda City, East Kalimantan, can be seen in Table 1:

Table 1. Phytochemical Screening Analysis of Sungkai Leaf Ethanol Extract

No.	Test	Reactor	Results	Observation
1	Flavonoid	Mg dan HCl 2N	+	Yellow color formed
2	Saponin	Hot aquades and HCl 2N	+	Stable foam formed for 10 minutes and with the addition of 2N HCl the foam did not disappear
3	Alkaloid	Dragendorf	+	An orange precipitate formed
4	Fenolik	FeCl ₃	+	Formed a bluish-green color
5	Steroid	Acetic anhydrous acid and sulfuric acid	+	Formed a bluish-green color

Phytochemical testing on Sungkai leaf extract consisted of flavonoids, saponins, alkaloids, phenols, and steroids. The results of the phytochemical screening showed that the ethanol extract of Sungkai leaves from Palaran District, Samarinda City, East Kalimantan, positively contained flavonoids, saponins, alkaloids, phenols, and steroids.

The ethanol extract of Sungkai leaves contains polar secondary metabolites. The content of phenolic and flavonoid compounds, which contain hydroxy groups, makes it easier for the polar solvent to attract these two compounds into the ethanol solvent, which is also rich in hydroxy groups (Ramadhani et al., 2022).

Based on the phytochemical tests conducted by (Rahman et al., 2022), it was found that Sungkai leaves originating from Merangin Regency, Jambi Province, also contain secondary metabolites of tannins, phenols, flavonoids, saponins, and alkaloids. Differences in the location where Sungkai grows can cause differences in the presence of secondary metabolites contained in the

number of secondary metabolites contained in the Sungkai extract.

Determination of Total Flavonoid Content by UV-Vis Spectrophotometry

This research aimed to determine the total flavonoids content contained in the ethanol extract of Sungkai leaves using the UV-Vis Spectrophotometry method. The choice of this method is because it has the advantage that the process is quite simple, can be used to determine tiny quantities of substances, the results obtained are pretty fast and accurate, and the numbers that are read are recorded directly by the detector and printed in the form of digital numbers or graphs that have been regressed. The drawbacks of this method are that absorption is affected by the pH of the solution, temperature, the presence of interfering substances, and the cleanliness of the cuvette; it is used only in functional groups containing valence electrons with low excitation energy; the light used must be monochromatic (Rohmah et al., 2021).

Determination of the levels of flavonoids contained in the ethanol extract of Sungkai leaves was carried out using the UV-Vis spectrophotometry method because flavonoids have a conjugated chromophore group and an autochrome group that binds directly to the chromophore group (Susilowati & Sari, 2021). The use of the quercetin standard in the determination of flavonoid levels is because the quercetin compound has

an amount of around 60-75% of the flavonoids. According to (Lindawati et al., 2020), the structure of quercetin is a flavonoid compound in the flavonol group with a keto group on C-4 and a hydroxyl group on C-3 and C-5 atoms. Quercetin is a flavonoid compound that can react with AlCl_3 and form complex reactions, which can be seen in Figure 1:




Figure 1. Formation of quercetin complexes with AlCl_3 reagent (Susilowati & Sari, 2021)

In determining the levels of flavonoids, a colorimetric reaction occurs because the sample reacts with AlCl_3 in acidic media. The addition of AlCl_3 will form a complex response between the flavonoids and AlCl_3 this causes a shift in the wavelength to the visible direction, producing a more yellow color (Lindawati et al., 2020). Furthermore, adding sodium acetate in this experiment maintains the wavelength in the visible area (Asmorowati, 2019).

Operating time measurement is carried out before the measurement is carried out. This is because there is a possibility that the reaction of the complex compound that is formed could be better. Still, on the contrary, if the measurement is carried out after the operating time, there will be a possibility that the complex compound between AlCl_3 has been damaged (Suharyanto & Prima, 2020). This operating time measurement aims to determine the time when the quercetin uptake is stable (Damayanti et al., 2023). And obtained in the operating time measurement received a stable absorbance value at 28-32 minutes.

Determining the maximum wavelength helps determine the measurement wavelength of a complex compound between quercetin and AlCl_3 to provide optimal absorbance (Damayanti et al., 2023). The maximum wavelength in this study was obtained at 443.0 nm.

Determining the standard curve intends to recognize the relationship between the concentration of the solution and its absorbance value so that the attention of the test sample can be known (Hilma et al., 2021). The absorption results were read during the operating time for 30 minutes, with a maximum wavelength of 443.0 nm, which can be seen in Figure 2. The results of the measurements that have been carried out are the standard curve equation $Y = 0.0037x + 0.2491$ with an R-calculated value of 0.9932. From that day, according to the R table statistical test with the number of data (n) = 5 at the 95% confidence level, the value of the R table is 0.8783. The results show that the R count exceeds the R table. This indicates a solid significant relationship between the standard quercetin concentration and absorbance and the linear regression equation

(Asmorowati, 2019). With that, the normal curve of the linear regression equation that has been obtained is used to calculate the

levels of flavonoids contained in the ethanol extract of Sungkai leaves.




Figure 1. Standard curve graph on quercetin

In determining the levels of flavonoids expressed in QE (Quercetin Equivalent). QE is the equivalent number of milligrams of quercetin in 1 gram of sample (Hadi et al., 2023). The result of determining the total flavonoid content was 18.210 ± 0.271 mg QE/g,

which means that every gram of ethanol extract of Sungkai leaves has the same value as 18.210 mg of quercetin and which can be seen in Table 2:

Table 2. Results of Total Flavonoid Content in Sungkai Leaf Ethanol Extract

Replication	Sample weight	Absorbance	Rate (mg QE/g)	Average	SD	CV
1	0,1004	0,592	18,461	18,210	0,271	1,486%
2	0,1002	0,581	17,905			
3	0,1003	0,587	18,210			
4	0,1005	0,593	18,497			
5	0,1001	0,582	17,977			

These calculations found that the CV value was 1.468%, which fulfilled the homogeneity requirements because the value was <5% (Sari & Fitrianiingsih, 2020). In a study by Adlis Santoni et al., (2023) on the hexane extract of Sungkai leaves originating from Padang City, West Sumatera, a total flavonoid content of 28.572 mgGE/g sample was obtained.

The Flavonoid content possessed by the ethanol extract of Sungkai leaves has oxidation-reducing properties, which have the potential as antioxidants. This is caused by the presence of hydroxy groups found in the

structure of flavonoids. Flavonoids are indispensable in various biological uses, traditional pharmaceutical preparations, synthetic drugs, and cosmetics. This is attributed to its antioxidant, anti-inflammatory, antimutagenic, and anticarcinogenic properties coupled with its ability to regulate the body's metabolism (Panche et al., 2016).

Determination of Total Phenolic Content by UV-Vis Spectrophotometry

This research aimed to determine the total phenolic content contained in the ethanol extract of Sungkai leaves using the UV-Vis

Spectrophotometry method. This method is chosen because UV-Vis Spectrophotometry has the advantages of having a low detection limit and a high degree of accuracy and precision (Bahrum et al., 2023). The drawbacks of this method are that the compound to be analyzed must have a chromophore group (color carrier group), conjugated double bonds, and a wavelength that lies in the ultraviolet or visible region (Tetha & Sugiarso, 2016).

This research was conducted to determine the levels of total phenolic

compounds in the ethanol extract of Sungkai leaves with the reference compound used, namely gallic acid solution, the reason for choosing gallic acid is because gallic acid is a derivative of hydroxybenzoate acid, which is classified as simple phenolic acid and the phenol content of gallic acid is pure (Tahir et al., 2017). In addition, gallic acid has a hydroxy group and conjugated double bonds in each benzene ring (Hilma et al., 2021), and the reaction can be seen in Figure 3:




Figure 1. The reaction of gallic acid with the Folin-Ciocalteu reagent (Hilma et al., 2021)

The phenolic-hydroxy group reduces the heteropoly acid (phosphomolybdic phosphotungstate) contained in the Folin-Ciocalteu reagent into a blue molybdenum-tungsten complex which UV-Vis Spectrophotometry can detect phenolic compounds into phenolic compounds. Therefore, NaOH is used to make the atmosphere alkaline (Andriani & Murtisiwi, 2018).

The operating time on gallic acid was obtained at 57-61 minutes. In operating time, it is known that the absorption value is said to be stable if the absorbance value is the same or seen from the relatively small difference in absorbance value. With that in mind, the purpose of determining operating time is to determine the relationship between absorbance and time in the solution to be used (Riyanti & Wilianita, 2023).

The maximum absorption wavelength of gallic acid was obtained in the 61st minute after adding the Folin-ciocalteu reagent and

NaOH using a UV-Vis spectrophotometer, and a blue solution was obtained. Then, absorption readings were taken in the range of 550-800 nm, and this absorption was used as the maximum wavelength. And the maximum wavelength obtained is 750.2 nm.

In determining the gallic acid standard curve, absorption readings were carried out during the operating time for 61 minutes and measured at a wavelength of 750.2 nm. The results of gallic acid absorbance can be seen in Figure 4. The results of the measurements that have been made are the standard curve equation $Y = 0.0095x + 0.0507$ with an R-calculated value of 0.9965. From that day, according to the R table statistical test with the number of data (n) = 5 at the 95% confidence level, the value of the R table is 0.8783. From the results obtained, it shows that the R count is greater than the R table. This indicates a solid significant relationship between the standard gallic acid concentration and the absorbance and the linear regression equation

(Asmorowati, 2019). With that, the standard curve of the linear regression equation that has been obtained is used to calculate the phenol content contained in the ethanol extract of Sungkai leaves.

In determining the total phenolic content of the ethanol extract of Sungkai

leaves, measurements were made at 61 minutes and a wavelength of 750.2 nm. The absorption results read can be seen in Table 3:




Figure 4. Standard curve graph for gallic acid

Table 3. Results of Total Phenolic Content in Sungkai Leaf Ethanol Extract

Replication	Sample weight	Absorbance	Rate (mg GAE/g)	Average	SD	CV
1	0,2003	0,582	34,902	33,666	1,052	3,126%
2	0,2001	0,551	32,898			
3	0,2002	0,567	33,933			
4	0,2004	0,573	34,293			
5	0,2001	0,542	32,306			

The result of determining the total phenolic content was 33.666 ± 1.052 mg GAE/g, which means that every gram of ethanol extract of Sungkai leaves has the same value as 33.666 mg of gallic acid. From these calculations, it was found that the CV value was 3.126%, where this value fulfilled the homogeneity requirements because the value was <5% (Sari & Fitrianiingsih, 2020). In a study by (Adlis Santoni et al., 2023) on the hexane extract of Sungkai leaves originating from Padang City, West Sumatra, had a phenolic content of 3.582 mgGAE/g samples.

Several studies have shown that antioxidant activity is proportional to the content of phenolic compounds. Antioxidants

are related to the ability to perform free anti-radical action from natural cell damage. Therefore, the flavonoids and phenols contained in Sungkai leaves are thought to function as antitumor, anti-inflammatory, and antimicrobial (Lubis et al., 2018).

Differences in the growing areas of the sample plants can cause the difference in total phenolic and flavonoid levels. This study used samples from Samarinda City, East Kalimantan, while a survey conducted by (Adlis Santoni et al., 2023) used samples from Padang City, West Sumatra. In addition, differences can be influenced by environmental factors such as rainfall,

temperature, ultraviolet radiation, and soil composition (Borges et al., 2013).

ACKNOWLEDGMENT

The authors are grateful to Departement of Education and Culture Office of East Kalimantan for providing a "Kaltim Tuntas" Scholarship 2023 to complete the Master of Pharmacy at Universitas Ahmad Dahlan.

CONCLUSION

It can be concluded from this study that the results of the phytochemical screening showed that the ethanol extract of Sungkai leaves originating from Samarinda City, East Kalimantan, positively contained flavonoids, saponins, alkaloids, phenols, and steroids. The results of determining the total flavonoids and phenolic content of the ethanol extract of Sungkai leaves were 18.210 ± 0.271 mg QE/g sample and 33.666 ± 1.052 mg GAE/g sample. Suggestions for future researchers, namely conducting activity tests on antioxidants from the ethanol extract of Sungkai leaves from Palaran District, Samarinda City, East Kalimantan.

REFERENCES

- Adlis, S. (2020). Aktivitas Antibakteri dan Sitotoksik Ekstrak Daun Sungkai (*Peronema canescens* Jack). *Jurnal Kimia UNAND*, 9(4), 303–401.
- Adlis Santoni, Mai Efdi, & Fadhillah, N. (2023). Profil Fitokimia dan Penentuan Fenolik Total, Flavonoid Total, dan Uji Aktivitas Antioksidan Ekstrak Daun Sungkai (*Peronema canescens* Jack) dari Daerah Kota Padang. *Jurnal Kimia Unand*, 12(1 SE-Articles), 1–6. <http://jku.fmipa.unand.ac.id/index.php/jku/article/view/25>
- Andriani, D., & Murtisiwi, L. (2018). Penetapan Kadar Fenolik Total Ekstrak Etanol Bunga Telang (*Clitoria ternatea* L.) Dengan Spektrofotometri Uv Vis. *Cendekia Journal of Pharmacy*, 2, 32–38. <https://doi.org/10.31596/cjp.v2i1.15>
- Asmorowati, H. (2019). Penetapan kadar flavonoid total buah alpukat biasa (*Persea americana* Mill.) dan alpukat mentega (*Persea americana* Mill.) dengan metode spektrofotometri UV-Vis. *Jurnal Ilmiah Farmasi*, 15, 51–63. <https://doi.org/10.20885/jif.vol15.iss2.art1>
- Bahrum, S., Umar, A., & Jasril, J. (2023). Pengaruh terhadap lama perendaman buah rosella (*Hibiscus sabdariffa* L) melalui uji kadar vitamin C secara spektrofotometri UV-VIS. *Jurnal SAGO Gizi Dan Kesehatan*, 4, 249. <https://doi.org/10.30867/gikes.v4i2.1209>
- Borges, L., Alves, S., Leite Sampaio, B., Conceição, E., Bara, M., & Paula, J. (2013). Environmental factors affecting the concentration of phenolic compounds in *Myrcia tomentosa* leaves. *Revista Brasileira de Farmacognosia*, 5. <https://doi.org/10.1590/S0102-695X2013005000019>
- Carolina, M., Araya, W., Carolina, P., & Permatasari Iskandar, D. (2022). Efektifitas Pemberian Seduhan Daun Sungkai (*Peronema canescens* Jack) Terhadap Perubahan Tekanan Darah Pada Lansia Hipertensi Di Wilayah Upt Puskesmas Pahandut Palangka Raya. *Jurnal Kesehatan Tambusai*, 3(3 SE-Articles), 442–452. <https://doi.org/10.31004/jkt.v3i3.6448>
- Damayanti, P., Luhumingtyas, F., & Indrayati, L. (2023). Penetapan Kadar Fenolik dan Flavonoid Total Ekstrak Etanol Buah Parijoto (*Medinilla speciosa* Blume) dengan Metode Spektrofotometri UV-Vis. *Jurnal Farmasi (Journal of Pharmacy)*, 12. <https://doi.org/10.37013/jf.v12i1.222>
- Fadlilaturrahmah, F., Putra, A., Rizki, M., & Nor, T. (2021). Uji Aktivitas Antioksidan dan Antitirozinase Fraksi n-Butanol Daun Sungkai (*Peronema canescens* Jack.) Secara Kualitatif Menggunakan Kromatografi Lapis Tipis. *Jurnal Pharmascience*, 8, 90.

- <https://doi.org/10.20527/jps.v8i2.11160>
- Febri Fatwami, E., Royani, S., & Korespondensi, P. (2023). Skrining Fitokimia dan Uji Antioksidan Ekstrak Daun Cabai Rawit (*Capsicum frutescens* L.). *Journal Syifa Sciences and Clinical Research*, 5(2). <https://doi.org/10.37311/JSSCR.V5I2.20896>
- Hadi, S., Subekti, A., & Khairunnisa, A. (2023). Antioxidant Test and Flavonoids Determination of Tuber Pakis Kinca (*Nephrolepis cordifolia* (L) C. Presl). *IJCA (Indonesian Journal of Chemical Analysis)*, 6(1), 1–9. <https://doi.org/10.20885/ijca.vol6.iss1.art1>
- Hilma, H., Putri, N., & Lely, N. (2021). Penentuan Kandungan Total Fenol Dan Total Flavonoid Ekstrak Daun Kelengkeng (*Dimoncarpus longan* Lour). *Jurnal Ilmiah Farmako Bahari*, 12, 80. <https://doi.org/10.52434/jfb.v12i1.1037>
- Imran, Nurliana, L., Natsir, M., Nohong, Kadir, L., Rudi, L., Ruslan, & Azis, T. (2023). Identifikasi dan Uji Aktivitas Antioksidan Senyawa Metabolit Sekunder Ekstrak Tanaman *Cleome viscosa* L: Identification and Antioxidants Activity Tests of Secondary Metabolite Compounds of *Cleome viscosa* L. Plant Extracts. *KOVALEN: Jurnal Riset Kimia*, 9, 100–106. <https://doi.org/10.22487/kovalen.2023.v9.i1.16340>
- Jumiarni, W. O., & Komalasari, O. (2017). Inventory of Medicines Plant As Utilized By Muna Tribe in Kota Wuna Settlement. *Majalah Obat Tradisional*, 22(1), 45. <https://doi.org/10.22146/tradmedj.24314>
- Larasati, D., & Putri, F. M. S. (2023). Skrining Fitokimia dan Penentuan Kadar Flavonoid Ekstrak Etanol Limbah Kulit Pisang (*Musa acuminata* Colla). *Jurnal Mandala Pharmacon Indonesia*, 9(1 SE-Original Articles), 125–131. <https://doi.org/10.35311/jmpi.v9i1.330>
- Lindawati, N. Y., Hudzaifah Ma'ruf, S., Tinggi, S., Kesehatan, I., & Surakarta, N. (2020). Penetapan Kadar Total Flavonoid Ekstrak Etanol Kacang Merah (*Phaseolus vulgaris* L.) Secara Spektrofotometri Visibel. *Jurnal Ilmiah Manuntung*, 6(1), 83–91. <https://doi.org/10.51352/JIM.V6I1.312>
- Lubis, S., Masfria, & Dalimunthe, A. (2018). Determination Of Total Phenolic Content, Total Flavonoid Content, And Antimutagenic Activity Of Ethanol Extract Nanoparticles Of *Rhaphidophora Pinnata* (L.F) Schott Leaves. *Rasayan Journal of Chemistry*, 11, 505–510. <https://doi.org/10.31788/RJC.2018.1122068>
- Panche, A. N., Diwan, A. D., & Chandra, S. R. (2016). Flavonoids: an overview. *Journal of Nutritional Science*, 5, e47. <https://doi.org/10.1017/jns.2016.41>
- Rahman, A., Rengganis, G. P., Prayuni, S., Novriyanti, I., Sari, T. N., Pratiwi, P. D., & Pratama, S. (2022). Pengaruh Pemberian Infusa Daun Sungkai (*Peronema canescens*) Terhadap Jumlah Leukosit Pada Mencit. *JOURNAL OF HEALTHCARE TECHNOLOGY AND MEDICINE*, 7(2), 614–620. <https://doi.org/10.33143/JHTM.V7I2.1574>
- Ramadhani, N., Giri Samudra, A., Pertiwi, R., Dwi Utami, C., Muslimah, A., Syahidah, W., Siti Khodijah Program Studi, P. S., Matematika dan Ilmu Pengetahuan Alam, F., & Bengkulu JI Supratman Kandang Limun Bengkulu, U. W. (2022). Analisis Total Fenol Dan Flavonoid Ekstrak Etanol Kulit Batang Sungkai (*Peronema canescens* Jack). *PHARMACY: Jurnal Farmasi Indonesia (Pharmaceutical Journal of Indonesia)*,

- 19(1), 66–79.
<https://doi.org/10.30595/PHARMACY.V19I1.12554>
- Rifkia, V., & Prabowo, I. (2020). Pengaruh Variasi Suhu dan Waktu terhadap Rendemen dan Kadar Total Flavonoid pada Ekstraksi Daun Moringa Oleifera Lam. dengan Metode Ultrasonik. *Pharmacy: Jurnal Farmasi Indonesia*, 17(02), 387–395.
<https://doi.org/10.30595/pharmacy.v17i2.7752>
- Riyanti, H. B., & Wilianita, R. A. (2023). Penetapan Kadar Tanin Dalam Ekstrak Etanol Daun Angsana (*Pterocarpus indicus Willd*) Hasil Maserasi Dan Sokletasi Menggunakan Spektrofotometer Uv-Vis. 8(1), 241–252.
- Rohmah, S., Muadifah, A., & Martha, R. (2021). Validasi Metode Penetapan Kadar Pengawet Natrium Benzoat pada Sari Kedelai di Beberapa Kecamatan di Kabupaten Tulungagung Menggunakan Spektrofotometer Uv-Vis. *Jurnal Sains Dan Kesehatan*, 3, 120–127.
<https://doi.org/10.25026/jsk.v3i2.265>
- Sadik, F. (2021). Uji Efektivitas Ekstrak Etanol Daun Jarak Pagar (*Jatropha curcas*. L) Sebagai Antihipertensi Pada Tikus. *Kieraha Medical Journal*, 3, 74–81.
<https://doi.org/10.33387/kmj.v3i2.3949>
- Sari, D. E. M., & Fitrianiingsih, S. (2020). Analisis Kadar Nilai Sun Protection Factor (SPF) pada Kosmetik Krim Tabir Surya yang Beredar di Kota Pati Secara In Vitro. *Cendekia Journal of Pharmacy*, 4(1), 69–79.
- Shofi, M., Suwitasari, F., & Istiqomah, N. (2020). Aktivitas Antioksidan Ekstrak Etanol Kamboja Jepang (*Adenium obesum*) dan Kamboja Putih (*Plumeria acuminata*). *Al-Kauniah: Jurnal Biologi*, 13, 167–178.
<https://doi.org/10.15408/kauniah.v13i1.12631>
- Silvani, M., Riga, R., & Agustini, D. (2023). Aktivitas Antioksidan Jamur Endofitik BS-1 yang Diisolasi dari Bunga Sambiloto Menggunakan Beras Putih sebagai Media Pertumbuhan. *Jurnal Sains Dan Kesehatan*, 5, 149–156.
<https://doi.org/10.25026/jsk.v5i2.1734>
- Suharyanto, S., & Prima, D. A. N. (2020). Penetapan Kadar Flavonoid Total pada Juice Daun Ubi Jalar Ungu (*Ipomoea batatas* L.) yang Berpotensi Sebagai Hepatoprotektor dengan Metode Spektrofotometri UV-Vis. *Cendekia Journal of Pharmacy*, 4(2), 110–119.
- Susilowati, S., & Sari, I. (2021). Perbandingan Kadar Flavonoid Total Seduhan Daun Benalu Cengkeh (*Dendrophthoe Petandra* L.) pada Bahan Segar dan Kering. *Jurnal Farmasi (Journal of Pharmacy)*, 9, 33–40.
<https://doi.org/10.37013/jf.v9i2.108>
- Tahir, M., Muflihunna, A., & Syafrianti, S. (2017). Penentuan Kadar Fenolik Total Ekstrak Etanol Daun Nilam (*Pogostemon cablin* Benth.) Dengan Metode Spektrofotometri UV-Vis. *Jurnal Fitofarmaka Indonesia*, 4, 215–218.
<https://doi.org/10.33096/jffi.v4i1.231>
- Tetha, D. A., & Sugiarto, D. (2016). Pebandingan Metode Analisa Kadar Besi antara Serimetri dan Spektrofotometer UV-Vis dengan Pengompleks 1,10- Fenantrolin. *Akta Kimia Indonesia*, 1, 8.
<https://doi.org/10.12962/j25493736.v1i1.1419>
- Widiasari, S. (2018). Mekanisme Inhibisi Angiotensin Converting Enzym Oleh Flavonoid Pada Hipertensi Inhibition Angiotensin Converting Enzym Mechanism By Flavonoid In Hypertension. *Collaborative Medical Journal (CMJ)*, 1(2), 30–44.
<http://jurnal.univrab.ac.id/index.php/cmj/article/view/474>

Phytochemical Screening Analysis and Determination of Total Flavonoids and Total Phenolics Content of Ethanol Extract of Sungkai Leaf (*Penorema canescens* Jack) from Samarinda City

ORIGINALITY REPORT

15%

SIMILARITY INDEX

12%

INTERNET SOURCES

9%

PUBLICATIONS

2%

STUDENT PAPERS

PRIMARY SOURCES

1 publikasi.ildikti10.id 1%
Internet Source

2 Rachmawaty, Andi Mu'nisa, Hasri, Halifah Pagarra, Hartati. "Analysis of phenolic content and antioxidant activity of cocoa pod husk (*theobroma cacao* L.)", *Journal of Physics: Conference Series*, 2019 1%
Publication

3 Erica Mega Valentine Simangunsong, Yessi Febriani, Muharni Saputri, Dina Arisa, Ghianti Zarqa Afifah. "EFFECTIVENESS OF BUTTERFLY PEA ETHANOL EXTRACT ON DECREASING BLOOD GLUCOSE LEVELS OF MALE MICE", *Jambura Journal of Health Sciences and Research*, 2023 1%
Publication

4 www.frontiersin.org 1%
Internet Source

journal2.uad.ac.id

5	Internet Source	1 %
6	Aminah Aminah, Nurhayati Tomayahu, Zainal Abidin. "PENETAPAN KADAR FLAVONOID TOTAL EKSTRAK ETANOL KULIT BUAH ALPUKAT (<i>Persea americana</i> Mill.) DENGAN METODE SPEKTROFOTOMETRI UV-VIS", <i>Jurnal Fitofarmaka Indonesia</i> , 2017 Publication	1 %
7	discovery.researcher.life Internet Source	1 %
8	ijrpr.com Internet Source	1 %
9	www.ejournalunb.ac.id Internet Source	<1 %
10	statik.unesa.ac.id Internet Source	<1 %
11	Madyawati Latief, Indra Lasmana Tarigan, Putri Maya Sari, Fiolita Etsa Aurora. "Aktivitas Antihiperurisemia Ekstrak Etanol Daun Sungkai (<i>Peronema canescens</i> Jack) Pada Mencit Putih Jantan", <i>Pharmacon: Jurnal Farmasi Indonesia</i> , 2021 Publication	<1 %
12	journals.umkt.ac.id Internet Source	<1 %

13

Ika Ristia Rahman. "Antioxidant and Sunscreen Activity Purified Extract of Kenikir Leaves (ETDK) and Tampoi Fruit Peel Extract (EKBT) Nanocream", Jurnal Farmasi Galenika (Galenika Journal of Pharmacy) (e-Journal), 2021

Publication

<1 %

14

Siti Jubaidah, Reksi Sundu, Nur Sabriningsih. "PENETAPAN KADAR FENOLIK TOTAL FRAKSI POLAR DAN NONPOLAR DAUN RAMBAI LAUT (Sonneratia caseolaris L.) DENGAN METODE SPEKTROFOTOMETRI UV-Vis", Jurnal Riset Kefarmasian Indonesia, 2019

Publication

<1 %

15

ebin.pub
Internet Source

<1 %

16

nepjb.com
Internet Source

<1 %

17

jurnalnasional.ump.ac.id
Internet Source

<1 %

18

Khuswatun Khasanah, Desy Nawangsari, Ikhwan Yuda Kusuma. "Solid Dispersion of Acetosal Using Polyvinyl Pyrrolidone (PVP) K-30 in Tablets with Direct Compressing Method", Indo. J. Chem. Res., 2023

Publication

<1 %

19	Internet Source	<1 %
20	Submitted to Lambung Mangkurat University Student Paper	<1 %
21	Submitted to Udayana University Student Paper	<1 %
22	Submitted to Sriwijaya University Student Paper	<1 %
23	eprints.uad.ac.id Internet Source	<1 %
24	www.scielo.br Internet Source	<1 %
25	doaj.org Internet Source	<1 %
26	garuda.kemdikbud.go.id Internet Source	<1 %
27	www.neliti.com Internet Source	<1 %
28	www.semanticscholar.org Internet Source	<1 %
29	www.ukm.my Internet Source	<1 %
30	anyflip.com Internet Source	<1 %

31	ethesis.lib.ku.ac.th Internet Source	<1 %
32	farmasi.unw.ac.id Internet Source	<1 %
33	jurnal.ulb.ac.id Internet Source	<1 %
34	repository.stikesdrsoebandi.ac.id Internet Source	<1 %
35	tsukuba.repo.nii.ac.jp Internet Source	<1 %
36	www.scilit.net Internet Source	<1 %
37	R Naufalin, Erminawati, R Wicaksono, A T Febryani, N Latifasari. "Antioxidant activity of kecombrang preserving powder using foam mat drying method", IOP Conference Series: Earth and Environmental Science, 2021 Publication	<1 %
38	bsmiab.org Internet Source	<1 %
39	dspace.umkt.ac.id Internet Source	<1 %
40	ejournal.umm.ac.id Internet Source	<1 %

41	journal.universitaspahlawan.ac.id Internet Source	<1 %
42	jurnal.peneliti.net Internet Source	<1 %
43	online-journal.unja.ac.id Internet Source	<1 %
44	prosiding.farmasi.unmul.ac.id Internet Source	<1 %
45	www.dovepress.com Internet Source	<1 %
46	www.freepatentsonline.com Internet Source	<1 %
47	www.jpms-stifa.com Internet Source	<1 %
48	Fendy R. Mondong. "Skrining Fitokimia dan Uji Aktivitas Antioksidan Ekstrak Etanol Daun Patikan Emas (<i>Euphorbia prunifolia</i> Jacq.) dan Bawang Laut (<i>Proiphys amboinensis</i> (L.) Herb)", Jurnal MIPA, 2015 Publication	<1 %
49	journal.uad.ac.id Internet Source	<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On