

HISTORY ARTIKEL KE 2

Mapping Rheumatoid Arthritis Susceptibility through Integrative Bioinformatics and Genomic

Sistem yang digunakan editorial dapat diakses melalui Link: <http://www.journal.uad.ac.id/index.php/Media-Farmasi/author/submission/24912>. Artikel di submit pada tanggal 25 September 2022. Artikel di review oleh 4 orang reviewer. Revisi artikel dilakukan pada tanggal 19 Oktober 2022 kemudian pada tanggal 17 Januari 2023, artikel dinyatakan diterima oleh Editor. Proses editing dan layout dilakukan sekitar satu pekan, dan pada Maret 2023, artikel diterbitkan secara online. Untuk lebih jelasnya dapat dilihat pada gambar berikut. Dokumen perubahan dari artikel yang diterbitkan sejak proses submit, hasil review dari reviewer, artikel revisi, dan artikel versi terbit, terlampir.

Mapping Rheumatoid Arthritis Susceptibility through Integrative Bioinformatics and Genomic

Nining Medi Sushanti¹, Wirawan Adikusuma^{2,3}, Arief Rahman Afief¹, Anita Silas La'ah⁴, Firdayani³, Rockie Chong⁵, Zainul Amiruddin Zakaria⁷, Barkah Djaka Purwanto^{8,9}, Rahmat Dani Satria^{10,11}, Riat El Khair^{10,11}, Abdi Wira Septama⁶, Lalu Muhammad Irham^{1,6*}

¹Faculty of Pharmacy, University of Ahmad Dahlan, Yogyakarta, 55164, Indonesia,

²Departement of Pharmacy, University of Muhammadiyah Mataram, Mataram, 83127, Indonesia

³Research Center for Vaccine and Drugs, National Research and Innovation Agency (BRIN), South Tangerang, Indonesia.

⁴Taiwan International Graduate Program in Molecular Medicine, National Yang Ming Chiao Tung University and Academia Sinica, Taipei, Taiwan

⁵Department of Chemistry and Biochemistry, University of California, Los Angeles, USA

⁶Traditional Medicine, National Research and Innovation Agency (BRIN), South Tangerang, Indonesia

⁷Borneo Research on Algesia, Inflammation and Neurodegeneration (BRAIN) Group, Department of Biomedical Sciences, Faculty of Medicines and Health Sciences, University Malaysia Sabah, Jalan UMS, Kota Kinabalu 88400, Sabah, Malaysia

⁸Faculty of Medicine, University of Ahmad Dahlan, Yogyakarta, 55191, Indonesia

⁹PKU Muhammadiyah Bantul Hospital, Bantul, Yogyakarta, 55711, Indonesia

¹⁰Department of Clinical Pathology and Laboratory Medicine, Faculty of Medicine, Public Health and Nursing, Universitas Gadjah Mada, Yogyakarta 55281, Indonesia

¹¹Clinical Laboratory Installation, Dr. Sardjito Central General Hospital, Yogyakarta 55281, Indonesia

*Correspondence author: Lalu Muhammad Irham; email: lalu.irham@pharm.uad.ac.id

ARTICLE INFO

ABSTRACT

Article history

Received: 25-09-2022

Revised: 19-10-2022

Accepted: 17-01-2023

Keywords

Rheumatoid arthritis,
Autoimmune,
Genome,
Bioinformatics,
Genetic variation

Rheumatoid arthritis (RA) is an autoimmune disease that influences several organs and tissues, especially the synovial joints, and is associated with multiple genetic and environmental factors. Numerous databases provide information on the relationship between a specific gene and the disease pathogenesis. However, it is important to further prioritize biological risk genes for downstream development and validation. This study aims to map RA-association genetic variation using genome-wide association study (GWAS) databases and prioritize influential genes in RA pathogenesis based on functional annotations. These functional annotations include missense/nonsense mutations, cis-expression quantitative trait locus (cis-eQTL), overlap knockout mouse phenotype (KMP), protein-protein interaction (PPI), molecular pathway analysis (MPA), and primary immunodeficiency (PID). 119 genetic variants mapped had a potential high risk for RA based on functional scoring. The top eight risk genes of RA are TYK2 and IFNGR2, followed by TNFRSF1A, IL12RB1 and CD40, C5, NCF2, and IL6R. These candidate genes are potential biomarkers for rheumatoid arthritis that can aid drug discovery and disease diagnosis.

This is an open access article under the [CC-BY-SA](https://creativecommons.org/licenses/by-sa/4.0/) license.



1. Introduction

Rheumatoid arthritis (RA) is the most common autoimmune disease in the world. It is an uncontrollable chronic inflammatory disease with various multi-system modalities followed by the proliferation of the synovial tissue that causes pain, joint erosion, and functional disorder (Garner et al., 2014). This synovial tissue proliferation can cause chronic impairments (Song & Lin, 2017). One sign of RA is persistent inflammation in the synovial joints. RA prevalence is within the age range of 40 – 60 which is between 0.5% and 1%, with greater prevalence in women than in men (Song & Lin,

2017). The main risk factors of RA are gender, family history, advanced age, exposure to silicate, and smoking cigarette (Garner et al., 2014) The primary influential determinants for RA vulnerability are genetic and environmental factors (Deane et al., 2017).

In this study, Genome-Wide Association Study (GWAS) is used to identify genetic variants that are at risk for disease pathogenesis by applying functional annotation criteria. This can also be useful in medicine through the bioinformatics-based approach (Narayanan, 2016; Reay & Cairns, 2021). GWAS is a research approach used in finding the genetic variation (Single sNucleotide Polymorphism, SNPs) associated with certain diseases, this involves the entire genome of several individuals. GWAS has been widely developed as a screening method not only in early-stage patients but also for people who are at risk based on family history. This research approach is useful for finding SNPs that contribute to complex diseases such as cancer, infectious diseases (AIDS, leprosy, hepatitis), autoimmune diseases, neuropsychiatric, and various diseases (Fareed & Afzal, 2013).

Meta-Analysis and GWAS have identified more than 100 RA loci, and GWAS is known as a database with genetic architecture also for RA (Kwon et al., 2020). GWAS is recognized as a powerful approach to map genes responsible for various diseases. For example, earlier studies have analyzed Chinese population SNPs that affect RA by using GWAS (Ren et al., 2014). In addition, the whole genome case-control of the linkage disequilibrium (LD) mapping was conducted by utilizing SNPs and RA-associated polymorphisms in PADI4 and SLC22A4/A5 cluster was identified (Ren et al., 2014). These findings showed that PADI4 is a risk gene for RA while SLC22A4/A5 has several polymorphisms associated with many autoimmune diseases. Thus, large-scale LD mapping seems effective at identifying RA polymorphisms (Ren et al., 2014).

Information on the relationship between a given gene and the pathogenesis of a disease is largely available on the GWAS database. The priority given to the most influential genes (biological risk genes) is scarcely examined. Thus, this research was conducted based on six functional annotations resulting in the prioritization of the most influential genes in RA pathogenesis.

2. Materials and Methods

2.1. RA-associated genes

Genomic variants RA risk genes were collected from the GWAS database, and the most influential genes were prioritized based on six functional annotations (Figure 1). Single nucleotide polymorphisms (SNPs) related to RA were expanded using the criterion $r^2 > 0.8$ of HaploReg v4.1. The SNPs associated with RA were denoted as "RA-associated genes". The genomic data were then prioritized based on six functional annotations, where genes with scores ≥ 2 were classified as "biological RA risk genes".

2.2. Functional annotations of RA-associated genes

Candidate genes most likely to be RA targets were prioritized based on their biological function using six functional annotations. Missense or nonsense mutations are single amino acid changes that produce different protein functions and are selected as functional annotation criteria (Zhang et al., 2012). In accordance with HaploReg v4.1, the first functional annotation is missense or nonsense mutation because it contains functional annotations from the SNPs database (DB) (Leng et al., 2020). HaploReg v4.1 was used to link cis-expression quantitative trait loci (cis-eQTL) to genetic variants in blood target tissues (Ward & Kellis, 2016). We also used WebGestalt 2019 to understand the relationship between mutant genes and phenotypes (Wang et al., 2017). Phenotype information of mouse and other mammals was derived from the ontology of the Mammalian Phenotype (MP) (Smith & Eppig, 2009). To further understand the phenotypes of mouse, we used BioMart to convert genes with human Ensembl ID into mouse Ensembl ID (Drost & Paszkowski, 2017). Genes with False Discovery Rates (FDRs) in mouse phenotypes < 0.05 were considered to be significant results. RA-associated genes from biological process networks were identified using protein-protein interactions (PPIs) (Qiu et al., 2021). In addition, we used WebGestalt 2019 to conduct reinforcement analysis and to investigate whether the genes were collected on certain functional annotations (Liao et al., 2019). Kyoto Encyclopedia of Genes and Genomes (KEGG) was used to determine what molecular pathways were gained from the list of RA-associated genes and what genes were involved. Further analysis of the biochemical pathway was carried out by WebGestalt 2019 database. The last annotation criterion was primary immunodeficiency (PID). PID refers to inborn immunity diseases and is often reported

to be associated with cancer (Mortaz et al., 2016). PID attacks the immune system and causes inflammation in RA (Dimitriadis & Sorensen, 2016). The hypergeometric test was employed for reinforcement analysis of these data, with the significance criterion being p -value < 0.05 .

Candidate genes most likely to be RA targets were prioritized based on their biological function using six functional annotations. Missense or nonsense mutations are single amino acid changes that produce different protein functions and are selected as functional annotation criteria (Z. Zhang et al., 2012). In accordance with HaploReg v4.1, the first functional annotation is missense or nonsense mutation because it contains functional annotations from the SNPs database (DB) (Leng et al., 2020). HaploReg v4.1 was used to link *cis*-expression quantitative trait loci (*cis*-eQTL) to genetic variants in blood target tissues (Ward & Kellis, 2016). We also used WebGestalt 2019 to understand the relationship between mutant genes and phenotypes (Wang et al., 2017). Phenotype information of mouse and other mammals was derived from the ontology of the Mammalian Phenotype (MP) (Smith & Eppig, 2009). To further understand the phenotypes of mouse, we used BioMart to convert genes with human Ensembl ID into mouse Ensembl ID (Drost & Paszkowski, 2017). Genes with False Discovery Rates (FDRs) in mouse phenotypes < 0.05 were considered to be significant results. RA-associated genes from biological process networks were identified using protein-protein interactions (PPIs) (Qiu et al., 2021). In addition, we used WebGestalt 2019 to conduct reinforcement analysis and to investigate whether the genes were collected on certain functional annotations (Liao et al., 2019). Kyoto Encyclopedia of Genes and Genomes (KEGG) was used to determine what molecular pathways were gained from the list of RA-associated genes and what genes were involved. Further analysis of the biochemical pathway was carried out by WebGestalt 2019 database. The last annotation criterion was primary immunodeficiency (PID). PID refers to inborn immunity diseases and is often reported to be associated with cancer (Mortaz et al., 2016). PID attacks the immune system and causes inflammation in RA (Dimitriadis & Sorensen, 2016). The hypergeometric test was employed for reinforcement analysis of these data, with the significance criterion being p -value < 0.05 .




Fig 1. Schematic model showing how genome-based information can be integrated to map susceptibility genes for rheumatoid arthritis (RA). [*cis*-expression quantitative trait locus (*cis*-eQTL); protein-protein interaction (PPI); molecular pathway analysis (MPA); and primary immunodeficiency (PID).]

3. Results and Discussion

3.1. Prioritized SNPs potentially cause RA

2,732 RA-associated SNPs were dictated in the GWAS database (Table S1) and 2012 SNPs were obtained after filtering based on p -value $< 10^{-8}$. 396 SNPs were obtained after checking for duplicates, and further filtering based on the inclusion criterion $OR > 2$, led to a total of 360 RA-associated SNPs (Table S2). The SNPs collected from GWAS were expanded using HaploReg v4.1 while the RA-associated genes were prioritized based on six functional annotations, namely missense/nonsense

mutation, cis-eQTL, knockout mouse phenotype (KMP), protein-protein interaction (PPI), molecular pathway analysis (MPA), and primary immunodeficiency (PID).

3.2. RA risk SNPs Functional Annotations

This study was designed to prioritize the best candidate genes by using six functional annotations with a scoring system. In the functional annotation process, candidate genes with higher scores represent greater biological influences on RA pathogenesis (Figure 2). Following the six annotations, we discovered 39 SNPs of missense/nonsense mutation variants from the total 360 SNPs (Figure 3). We further examined the cis-eQTL effect using HaploReg v4.1, 101 SNPs were gained from the total 360 SNPs (Figure 3). Phenotype data was taken from the MP ontology which contains information about the phenotype of mice and other mammals. WebGestalt 2019 was applied to perform over-representation analysis (ORA) and we found 102 genes overlapping with RA risk genes (FDR <0.05). Gene Ontology (GO) annotations derived from WebGestalt 2019 were used to evaluate PPIs. Following the analysis on PPIs, 122 genes overlap with other RA risk genes (FDR < 0.05). KEGG was used to perform ORA on the molecular pathway. Thus, over 61 RA-associated genes on KEGG pathways were identified. We also used the PID data from IUIS to analyze and confirm the overlapping genes. Sixteen overlapping genes were statistically significant ($p < 0.05$) from PID analysis.

We also scored the risk genes from 0 to 6 and obtained 156 genes with a score of 0. 85 genes have a score of 1, 45 genes have a score of 2, 40 genes have a score of 3, 26 genes have a score of 4, 6 genes have a score of 5, and 2 genes have a score of 6 (Figure 4). In total, we obtained 119 genes with a score of ≥ 2 , that were categorized as ‘biological RA risk genes’ (Table S2). Using the aforementioned scoring criteria, the top eight RA risk genes are: tyrosine kinase 2 (TYK2), followed by interferon-gamma receptor 2 (IFNGR2). Members of the tumor necrosis factor receptor superfamily 1A (TNFRSF1A), interleukin receptor subunit 12 beta 1 (IL12RB1) and mass differentiation 40 (CD40), complement 5 (C5), neutrophil cytosolic factor 2 (NCF2), and interleukin receptor 6 (IL6R).

Gencode_id	Gencode_name	missense	cis-eQTL	KO mice	PPI	KEGG	PID	Total Score
ENSG00000105397	TYK2	1	1	1	1	1	1	6
ENSG00000159128	IFNGR2	1	1	1	1	1	1	6
ENSG00000067182	TNFRSF1A	0	1	1	1	1	1	5
ENSG00000096996	IL12RB1	0	1	1	1	1	1	5
ENSG00000101017	CD40	0	1	1	1	1	1	5
ENSG00000106804	C5	0	1	1	1	1	1	5
ENSG00000116701	NCF2	1	0	1	1	1	1	5
ENSG00000160712	IL6R	1	1	1	1	1	0	5
ENSG00000003400	CASP10	0	1	0	1	1	1	4
ENSG00000020633	RUNX3	1	0	1	1	1	0	4
ENSG00000056558	TRAF1	0	1	1	1	1	0	4
ENSG00000064012	CASP8	0	0	1	1	1	1	4
ENSG00000076662	ICAM3	1	1	0	1	1	0	4
ENSG00000081237	PTPRC	0	0	1	1	1	1	4
ENSG00000081985	IL12RB2	0	1	1	1	1	0	4
ENSG00000100906	NFKBIA	0	0	1	1	1	1	4
ENSG00000103313	MEFV	1	0	1	1	0	1	4
ENSG00000103811	CTSH	1	1	0	1	1	0	4
ENSG00000110448	CD5	1	0	1	1	1	0	4
ENSG00000112486	CCR6	0	1	1	1	1	0	4
ENSG00000115267	IFIH1	1	0	1	1	1	0	4
ENSG00000115604	IL18R1	0	1	1	1	1	0	4
ENSG00000117560	FASLG	0	0	1	1	1	1	4
ENSG00000118503	TNFAIP3	1	0	1	1	1	0	4

Fig. 2. Functional annotations for rheumatoid arthritis (RA) with scores ≥ 2 based on defined scoring criteria

Tyrosine kinase 2 (TYK2) is a member of the Janus kinase (JAK) family of transforming growth factors and cytokines. The TYK2 loss analysis function reveals its importance toward immunity against infections, inflammation (automatic), and immunity (automatic). Many genome-wide association studies in humans propose relationships between TYK2 genetic variants and many autoimmune diseases, inflammatory diseases, and tumors. Hence, TYK2 emerges as a target of interest for therapeutic interventions (Leitner et al., 2017). The human interferon-gamma receptor is a multimer of IFN- γ R1 (coded by IFNGR1) and IFN- γ R2 (two chains). Interferon-gamma receptor 2

(IFN- γ R2) is a gene containing a protein and in humans, this gene is coded by the IFNGR2 gene. The IFNGR2 gene codes for beta chains that are not bound to the ligands of the IFN- γ R.

Tumor necrosis factor receptor 1A (TNFRSF1A) is a CD120a which is a member of the superfamily tumor necrosis factor alpha-binding membrane receptor. The TNFRSF1A gene instructs the making of tumor necrosis factor receptor 1 (TNFR1) protein (Chen et al., 2015). This protein is found stretching on the cell membrane, both extracellularly and intracellularly (Chen et al., 2015). Extracellularly, the TNFR1 protein enchains tumor necrosis factor/TNF. The TNF and the TNFR1 protein interaction catalyzes the latter to chain to two other TNFR1 proteins, constituting trimer, a three-protein complex (Chen et al., 2015). This formation of the trimer is important for the TNFR1 protein to function. The chains of the protein of TNF and TNFR1 prompt the latter to deliver a signal into the cell (Chen et al., 2015). The TNFR1 protein signal can stimulate inflammation or cell self-destruction (apoptosis). The signal into the cell starts a routeway activating the nuclear factor kappa B protein triggering inflammation and directing to the cytokine production, the immune system protein (Chen et al., 2015).




Fig. 3. Number of genes for each functional annotation that is prioritized for rheumatoid arthritis. [cis-expression quantitative trait locus (cis-eQTL); Knockout mice (KOmice); protein-protein interaction (PPI); Kyoto Encyclopedia of Genes and Genomes (KEGG); and primary immunodeficiency (PID).]




Fig. 4. Gene frequency distribution across defined functional scores for rheumatoid arthritis, showing 156 genes with a score of 0.85, genes with a score of 1, and 119 (45 + 40 + 26 + 6 + 2) genes with a total score of ≥ 2 are classified as RA risk genes.

Interleukin-12 receptor beta-1 subunit (IL12RB1) functions as an interleukin receptor that binds to interleukin-12 with low affinity which is involved in IL12 transduction (Floss et al., 2016). IL12RB2 makes a functional high-affinity receptor for IL12 (Chen et al., 2015). IL23R forms the receptor of interleukin-23 functioning in IL23 signal transduction via JAK/STAT signaling cascade activation (Floss et al., 2016). CD40 is a member of the tumor necrosis factor (TNF) superfamily found in antigen-presenting cells, including dendritic cells (DC), B cells, monocytes, and macrophages, which play an important role in activating the immune system (Karnell et al., 2019). The expression expands

to various non-hematopoietic cells, including nerve cells, epithelial cells, endothelial cells, and fibroblasts (Karnell et al., 2019). Typical genetic variants at the TRAF1-C5 locus on the chromosome correlate with an increased risk of anti-ccp-positive RA (Huang et al., 2019).

Genetic variants rs7021206 and rs3761847 at the TRAF1-C5 locus are associated with RA in the Han Chinese population, indicating that TRAF1-C5 may play an important role in the development of RA thereby reducing the pathogenic role of TRAF1-C5 in ethnic variations (van Steenbergen et al., 2015). NCF2 is carried to the membrane of the cell to be combined with other components to build a NOX system that is active through microbial stimulation. In other studies, the polymorphism NCF2 rs10911363 was linked to SLE risk in Sweden and US populations. The variant of NCF2 rs789181 was discovered to have a relationship with mild RA in men (T.-P. Zhang et al., 2020). IL6 is an important cytokine in mediating inflammation and RA systemic overview, including synovitis, fatigue, anemia, anorexia, and osteoporosis (Narazaki et al., 2017).

In this study, we used the GWAS database to obtain the genomic variants for RA. The GWAS database is easily accessible and effective because it is diverse and structured. The GWAS method effectively identifies genetic loci associated with the pathogenesis of a disease, including prioritizing candidate loci, exploring disease mechanisms, predicting disease risk, identifying new drug targets, and gathering information about the desired trait or population (Caliskan et al., 2021). However, the GWAS database has drawbacks in that detailed phenotypic descriptions of the study's ethnic background are not provided by many GWAS papers, and even nowadays, there is no standard way to make an ethnicity report (Hart & Kranzler, 2015). The prospect for further research is to develop a wider use of functional annotation criteria to get more results to determine genetic variations that affect RA pathogenesis and identify drug candidates that target biological genes for RA risk. The impact of this research is to find out the genetic variation that influences the pathogenesis of RA and gain an understanding by utilizing genetic variation that can be used for drug development by using a bioinformatics approach.

4. Conclusion

Our research showed that genomic information is highly important for mapping RA genomic variants based on defined functional annotations from this study. Two candidate genes, TYK2 and IFNGR2, were obtained based on defined scoring criteria, and we propose these genes as potential RA biomarkers to be prioritized for downstream clinical development and validation.

Author Contributions: Nining Medi Sushanti, Lalu Muhammad Irham, Wirawan Adikusuma, and Lalu Muhammad Irham conceived and designed the study. Nining Medi Sushanti and Lalu Muhammad Irham performed the computational analysis. Nining Medi Sushanti wrote the manuscript. Nining Medi Sushanti, Lalu Muhammad Irham., Wirawan Adikusuma, Anita Silas La'ah, Arief Rahman Afief, Zainul Amiruddin Zakaria, Firdayani, Rahmat Dani Satria, Riat El Khair, Abdi Wira Septama and Rockie Chong. revised the manuscript. Lalu Muhammad Irham supervised and coordinated this study. All authors have read and approved the manuscript and have made significant contributions to this study.

Funding

No funding was provided for this research.

Competing Interests

The authors have not declared a conflict of interest for this article.

Acknowledgment

All authors would like to appreciate Universitas Ahmad Dahlan for allowing them to carry out this study.

References

Caliskan, M., Brown, C. D., & Maranville, J. C. (2021). A catalog of GWAS fine-mapping efforts in autoimmune disease. *The American Journal of Human Genetics*, 108(4), 549–563. <https://doi.org/10.1016/j.ajhg.2021.03.009>

- Chen, W., Xu, H., Wang, X., Gu, J., Xiong, H., & Shi, Y. (2015). The tumor necrosis factor receptor superfamily member 1B polymorphisms predict response to anti-TNF therapy in patients with autoimmune disease: A meta-analysis. *International Immunopharmacology*, 28(1), 146–153. <https://doi.org/10.1016/j.intimp.2015.05.049>
- Deane, K. D., Demoruelle, M. K., Kelmenson, L. B., Kuhn, K. A., Norris, J. M., & Holers, V. M. (2017). Genetic and environmental risk factors for rheumatoid arthritis. *Best Practice & Research Clinical Rheumatology*, 31(1), 3–18. <https://doi.org/10.1016/j.berh.2017.08.003>
- Dimitriades, V. R., & Sorensen, R. (2016). Rheumatologic manifestations of primary immunodeficiency diseases. *Clinical Rheumatology*, 35(4), 843–850. <https://doi.org/10.1007/s10067-016-3229-6>
- Drost, H. G., & Paszkowski, J. (2017). Biomart: Genomic data retrieval with R. *Bioinformatics*, 33(8), 1216–1217. <https://doi.org/10.1093/bioinformatics/btw821>
- Fareed, M., & Afzal, M. (2013). Single nucleotide polymorphism in genome-wide association of human population: A tool for broad spectrum service. *Egyptian Journal of Medical Human Genetics*, 14(2), 123–134. <https://doi.org/10.1016/j.ejmhg.2012.08.001>
- Floss, D. M., Klöcker, T., Schröder, J., Lamertz, L., Mrotzek, S., Strobl, B., Hermanns, H., & Scheller, J. (2016). Defining the functional binding sites of interleukin 12 receptor β 1 and interleukin 23 receptor to Janus kinases. *Molecular Biology of the Cell*, 27(14), 2301–2316. <https://doi.org/10.1091/mbc.E14-12-1645>
- Garner, R., Ding, T., & Deighton, C. (2014). Management of rheumatoid arthritis. *Medicine*, 42(5), 237–242. <https://doi.org/10.1016/j.mpmed.2014.02.004>
- Hart, A. B., & Kranzler, H. R. (2015). Alcohol dependence genetics: lessons learned from genome-wide association studies (GWAS) and post-GWAS analyses. *Alcoholism: Clinical and Experimental Research*, 39(8), 1312–1327. <https://doi.org/10.1111/acer.12792>
- Huang, S.-C., Hua, D.-J., Sun, Q.-Q., Zhang, L.-N., Cen, H., & Zhou, L. (2019). Associations of TRAF1/C5 rs10818488 and rs3761847 polymorphisms with genetic susceptibility to rheumatoid arthritis: a case-control study and updated meta-analysis. *Central European Journal of Immunology*, 44(2), 159–173. <https://doi.org/10.5114/ceji.2019.87067>
- Karnell, J. L., Rieder, S. A., Ettinger, R., & Kolbeck, R. (2019). Targeting the CD40-CD40L pathway in autoimmune diseases: Humoral immunity and beyond. *Advanced Drug Delivery Reviews*, 141, 92–103. <https://doi.org/10.1016/j.addr.2018.12.005>
- Kwon, Y.-C., Lim, J., Bang, S.-Y., Ha, E., Hwang, M. Y., Yoon, K., Choe, J.-Y., Yoo, D.-H., Lee, S.-S., Lee, J., Chung, W. T., Kim, T.-H., Sung, Y.-K., Shim, S.-C., Choi, C.-B., Jun, J.-B., Kang, Y. M., Shin, J.-M., Lee, Y.-K., ... Bae, S.-C. (2020). Genome-wide association study in a Korean population identifies six novel susceptibility loci for rheumatoid arthritis. *Annals of the Rheumatic Diseases*, 79(11), 1438–1445. <https://doi.org/10.1136/annrheumdis-2020-217663>
- Leitner, N. R., Witalisz-Siepracka, A., Strobl, B., & Müller, M. (2017). Tyrosine kinase 2 – Surveillant of tumours and bona fide oncogene. *Cytokine*, 89, 209–218. <https://doi.org/10.1016/j.cyto.2015.10.015>
- Leng, R. X., Di, D. S., Ni, J., Wu, X. X., Zhang, L. L., Wang, X. F., Liu, R. S., Huang, Q., Fan, Y. G., Pan, H. F., Wang, B., & Ye, D. Q. (2020). Identification of new susceptibility loci associated with rheumatoid arthritis. *Annals of the Rheumatic Diseases*, 79(12), 1565–1571. <https://doi.org/10.1136/annrheumdis-2020-217351>
- Liao, Y., Wang, J., Jaehnig, E., Shi, Z., & Zhang, B. (2019). WebGestalt 2019: gene set analysis toolkit with revamped UIs and APIs. *Nucleic Acids Research*, 47. <https://doi.org/10.1093/nar/gkz401>

- Mortaz, E., Tabarsi, P., Mansouri, D., Khosravi, A., Garssen, J., Velayati, A., & Adcock, I. M. (2016). Cancers related to immunodeficiencies: Update and perspectives. *Frontiers in Immunology*, 7(SEP), 1–13. <https://doi.org/10.3389/fimmu.2016.00365>
- Narayanan, R. (2016). Repurposing drugs in the genomics era: bioinformatics approaches. *MOJ Proteomics & Bioinformatics*, 3(4). <https://doi.org/10.15406/mojpb.2016.03.00092>
- Narazaki, M., Tanaka, T., & Kishimoto, T. (2017). The role and therapeutic targeting of IL-6 in rheumatoid arthritis. *Expert Review of Clinical Immunology*, 13(6), 535–551. <https://doi.org/10.1080/1744666X.2017.1295850>
- Qiu, S., Munir, A., Malik, S. I., Khan, S., & Hassan, A. (2021). Identification of differentially expressed genes and pathways crosstalk analysis in Rheumatoid and Osteoarthritis using next-generation sequencing and protein-protein networks. *Saudi Journal of Biological Sciences*, 28(8), 4656–4663. <https://doi.org/10.1016/j.sjbs.2021.04.076>
- Reay, W. R., & Cairns, M. J. (2021). Advancing the use of genome-wide association studies for drug repurposing. *Nature Reviews Genetics*, 22(10), 658–671. <https://doi.org/10.1038/s41576-021-00387-z>
- Ren, T., Han, Z., Yang, C., Hang, Y., Fang, D., Wang, K., Zhu, X., Ji, X., & Zhou, F. (2014). Association of SLC22A4 gene polymorphism with rheumatoid arthritis in the chinese population. *Journal of Biochemical and Molecular Toxicology*, 28(5), 206–210. <https://doi.org/10.1002/jbt.21554>
- Smith, C. L., & Eppig, J. T. (2009). The mammalian phenotype ontology: Enabling robust annotation and comparative analysis. *Wiley Interdisciplinary Reviews: Systems Biology and Medicine*, 1(3), 390–399. <https://doi.org/10.1002/wsbm.44>
- Song, X., & Lin, Q. (2017). Genomics, transcriptomics and proteomics to elucidate the pathogenesis of rheumatoid arthritis. *Rheumatology International*, 37(8), 1257–1265. <https://doi.org/10.1007/s00296-017-3732-3>
- van Steenberg, H. W., Rodríguez-Rodríguez, L., Berglin, E., Zhernakova, A., Knevel, R., Ivorra-Cortés, J., Huizinga, T., Fernández-Gutiérrez, B., Gregersen, P. K., Rantapää-Dahlqvist, S., & van der Helm-van Mil, A. (2015). A genetic study on C5-TRAF1 and progression of joint damage in rheumatoid arthritis. *Arthritis Research & Therapy*, 17(1), 1. <https://doi.org/10.1186/s13075-014-0514-0>
- Wang, J., Vasaikar, S., Shi, Z., Greer, M., & Zhang, B. (2017). WebGestalt 2017: A more comprehensive, powerful, flexible and interactive gene set enrichment analysis toolkit. *Nucleic Acids Research*, 45(W1), W130–W137. <https://doi.org/10.1093/nar/gkx356>
- Ward, L. D., & Kellis, M. (2016). HaploReg v4: Systematic mining of putative causal variants, cell types, regulators and target genes for human complex traits and disease. *Nucleic Acids Research*, 44(D1), D877–D881. <https://doi.org/10.1093/nar/gkv1340>
- Zhang, T.-P., Li, R., Huang, Q., Pan, H.-F., Ye, D.-Q., & Li, X.-M. (2020). Association of NCF2, NCF4, and CYBA Gene Polymorphisms with Rheumatoid Arthritis in a Chinese Population. *Journal of Immunology Research*, 2020, 1–11. <https://doi.org/10.1155/2020/8528976>
- Zhang, Z., Miteva, M. A., Wang, L., & Alexov, E. (2012). Analyzing effects of naturally occurring missense mutations. *Computational and Mathematical Methods in Medicine*, 2012. <https://doi.org/10.1155/2012/805827>