Rekapitulasi Proses Penerbitan Artikel Syarat Khusus Pengajuan Lektor Kepala

"Influence of croscarmellose in fast disintegrating tablet of *Syzygium polyanthum* extract" International Journal of Public Health Science (IJPHS), Vol 13, No. 1, 2024 p-ISSN 2252-8806, e-ISSN 2620-4126 pp. 438-446 apt. Widyasari Putranti, M.Sc

No	Keterangan	Tanggal
1	Submit Artikel	1 November 2022
2	Email mendapatkan respon dari pengelola jurnal	1 November 2022
3	Email informasi melacak kemajuan naskah dalam proses	1 November 2022
	editorial	
4	Catatan Penelaahan Awal dan PermohonanPerbaikan Artikel	24 Maret 2023
5	Pengiriman revisi naskah sesuai catatanpenelaahan	9 Mei 2023
6	Pemberitahuan Penerbitan	16 Mei 2023
7	Permohonan Penyesuaian Konten Artikel	10 Oktober 2023
8	Pengiriman Revisi artikel, bukti transfer dan form transfer	28 Juli 2023
	copyright	
9	Penerbitan Artikel	21 Desember 2023

Lampiran :

Submit Artikel

Home > User > Author > Submissions > #22666 > Summary

#22666 Summary

SUMMARY REVIEW EDITING

Submission

Nuthors Widyasari Putranti, Desty Restia Rahmawati, Nining Sugihartini, Teuku Nanda Saifullah				
Title	Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract			
Original file	22666-44108-1-SM.DOCX 2022-11-01			
Supp. files	None			
Submitter	apt widyasari putranti 🕮			
Date submitted	November 1, 2022 - 06:36 AM			
Section	Other_Topics_in_Public_Health_Science			
Editor	Veronique Gucht 🕮 (Review) Henry Imhonde 🖻 (Review) Jennifer Spencer 🕮 (Review)			
Abstract Views	172			
Status				
Status	Published Vol 13, No 1: March 2024			
Initiated	2023-12-21			
Last modified	2023-12-22			
Submission Met	adata			
A 11				

Authors

Name

Widyasari Putranti 🖾

Email mendapatkan respon dari pengelola jurnal



Bukti Revisi dalam bentuk tabel

Tabel Perbaikan

"Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract"

No	Original text	Reviewer's comment	Revised text
1	Plant germplasm resources as raw materials for	Separate citations with	Plant germplasm resources are raw
	medicines, one of which is bay leaf (Syzygium polyanthum)[1].	commas, For example in	materials for medicines, one of
	This plant belongs to the Myrtaceae family[2]. Empirically,	your paper, citation [7][8]	which is a bay leaf (Syzygium
	herbal medicinal ingredients are widely used by Indonesian	can be written: [7], [8].	polyanthum). Bay leaves are known
	people as alternative therapies in overcoming various diseases,	- Simplify multiple	to contain secondary metabolites,
	such as: hyperlipidemia or high cholesterol levels.[3][4][5][6]	citations, for example in	such as saponins, terpenoids,
	Bay leaves are known to contain secondary metabolites,	your paper [3][4][5][6]., it	flavonoids (quercetin), polyphenols,
	such as saponins, terpenoids, flavonoids (quercetin),	should be write: [3]-[6].	alkaloids, steroids, and essential oils
	polyphenols, alkaloids, steroids, and essential oils	-Write citation from small	(sesquiterpenes)[1]-[2]
	(sisquiterpenes)[7][8]. Based on the Decree of the Minister of	number. For example in	This properties of FDT can release
	Health of the Republic of Indonesia (2009), bay leaves contain a	your paper: [10][6] it	the active drug substance
	total flavonoid not less than 0.40% which is calculated as	should be write: [6], [10].	immediately, speed up the onset,
	quercetin[9]. Quercetin in bay leaves has potential as an	-If you put citation at the	increase the oral bioavailability of
	antihyperlipidemic which has been shown to be able to	end of the sentence make	the drug in the body so that
	significantly reduce levels of triglycerides, total cholesterol, and	sure it comes before	therapeutic effectiveness can be
	LDL cholesterol in plasma and tissues in hyperlipidemic rats	".". For example in your	achieved [6]-[7].
	with a parallel increase in HDL, as well as inhibiting HMG -CoA	paper: " cholesterol	
	activity, reductase and LDL oxidation.[10][6]	levels.[3][4][5][6]" it	
		should be write:	
		cholesterol levels [3]-[6].	
		CHECK ALL citing	
		references and do the	
		revision. Use	
		Mendeley/reference	
		manager to help you	
		manage the citations.	
2	Requirements and weight uniformity test are determined based	In subsection "Weight	Requirements and weight uniformity
	on the Perka BPOM RI concerning the quality requirements of	uniformity test" the Perka	tests are determined based on the
	traditional medicines	BPOM RI can be write in	regulation of the National Food and

3	Quercetin in bay leaves has potential as an antihyperlipidemic which has been shown to be able to significantly reduce levels of triglycerides, total cholesterol, and LDL cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in HDL, as well as inhibiting HMG -CoA activity, reductase and LDL oxidation.[10][6]	italic style and describe it, since it is using local language. Please note that not all the readers understand your language. Use English so that your idea is understood by all the readers. Or you can inform the "BPOM RI" in nomenclature or explain more in the introduction. In second paragraph of introduction, there are abbrevitations: LDL, HDL, HMG -CoA, for the first appearance in a paper, please expand the abbreviation(s)/acronym(s). Also, write explanation first then abbrevitation. For example in your paper: LOD (Loss on Drying), MC (Moisture Content), write as: loss on drying (LOD), moisture content (MC). CHECK ALL abbrevitation	Drug Agency of the Republic of Indonesia concerning the quality requirements of traditional medicines. The test was carried out by weighing 20 tablets one by one, from 20 tablets no more than 2 tablets, each of which deviated from the average weight by more than 5% and not one tablet whose weight deviated from the average weight. the average is greater than 10% [12]. Quercetin in bay leaves the potential as an antihyperlipidemic which can significantly reduce levels of triglycerides, total cholesterol, and low-density lipoprotein (LDL) cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in high-density lipoprotein (HDL), as well as inhibiting 3- hidroksi-3-metilglutaril coenzyme A (HMG-CoA), reductase, and LDL oxidation [4].
4	Bay leaf simplicia was tested for macroscopic (organoleptic) specifications on fresh and dried simplicia, as well as microscopic tests for simplicia parts and identification fragments of simplicia powder (figures 1 and 2). The simplicia specification test was carried out to ensure the correctness of the identity and quality of the bay leaf simplicia based on the Indonesian Herbal Pharmacopoeia compendia and Indonesian Materia Medika. The results of the specification test for the simplicia of bay leaves showed that the simplicia used met the requirements of organoleptic and microscopic terms. The determination test of bay leaf simplicia was in accordance with its species or identity.[9] $figure 1. fragments of bay leaf powder (a) upper epidermis (b) lower epidermis figure 2. identification fragments of bay leaf, (a) upper epidermis, (b) floem, (c) calcium oxalate crystal, (c) sclerenchyma$	ub-Figures 1(a)-(b) and 2(a)-(d) are not mentioned in the text. If a figure contains subfigures, explain what the whole figure illustrates before explaining what the subfigures do. This is important to make the readers get what the relation between those subfigures is, and why they are put in the same figure's number instead of making them two different figures in separate numbering. Complete the description by explaining the relation of the images, why these images should be grouped together in the same image as subfigures of 2: (c) sclerenchyma maybe it is (d) sclerenchyma.	Fresh simplicia of bay leaves were sorted wet, dried using an oven (50- 60°C) with a moisture content of not more than 10%, then sorted dry. Simplicia powder for extract preparation is a fine simplicial (No. 60 mesh) in order to optimize the extraction process [18]. The results of the simplicial extraction of bay leaves weighing 3.7 kg obtained a thick extract of 347 grams. The total yield of the thick extract obtained was 9.38%. The morphology of the viscous extract is shown in Figure 2. Screening and determination of the chemical compound flavonoid quercetin contained in the thick extract of bay leaves was carried out at the Integrated Research Laboratory of the Faculty of Pharmacy, UAD. According to the Ministry of Health (2017), bay leaves contain a total flavonoid of not less than 0.40% calculated as quercetin. Based on the results of the test the flavonoid content of 0.8079±0.0045%, so that it meets the specifications for the total flavonoid content of bay leaves[17]. Quercetin is an active substance that has medicinal effects such as antiherlipidemia, antibacterial, antidiabetic, antioxidant [18], [19, □

			441 (a) (b) (c) (d) (e) Figure 1. Identification fragments of bay leaf; (a) lower epidermis, (b) upper epidermis, (c) floem, (d) calcium oxalate crystal, (e) sclerenchyma
5	Superdisintegrant croscarmellose sodium intra granular can increase the speed of disintegration, increase the extra-granular concentration significantly by reducing the wetting time and disintegration time, reducing the brittleness rate, and increasing the hardness of FDT. The composition of the optimum formula FDT bay leaf extract consists of CCS intra granular (10 mg) and extra granular (25 mg).	What implication of Conclusion	In the FDT research, to improve the poor flow properties of the extract, the wet granulation method was carried out. The IPC results for the three formulas showed granules that met the requirements. This research provides information that the use of extra granular CCS at higher levels will increase the rate of disintegration. The combination of CCS intra granular 10 mg (2%) and extra granular 25 mg (5%) can produce FDT with the most effective. By combining intra and extra granular super disintegrants, a good level of tablet hardness can be obtained so that the tablet is not brittle and has a faster disintegration time. In addition, FDT bay leaves extract can be used as an alternative treatment for hyperlipidemia in patients who experience dysphagia. This research still needs development to determine the stability of FDT bay leaves extract where the active substance is herbal medicine
6	Formulation of Intra and Extra-Granularly Croscarmellose in	Title <10	Influence of croscarmellose in fast
	Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium		disintegrating tablet of Syzygium
i	poryanunani	1	poryannium exitaet

Email informasi melacak kemajuan naskah dalam proses editorial

	÷		13 dari 23	<	>
10.075		[IJPHS] Submission Acknowledgement (Eksternal) Kotak Masuk ×		ð	ß
		Lina Handayani ijphs@iaescore.com l <u>ewat</u> smtpcorp.com 1 No kepada saya ✔	ov 2022, 13.36 🛛 📩	¢	:
46	H	lati-hati dengan pesan ini.			
	ir	ina handayani mirip dengan sebuah nama di organisasi Anda, tetapi alamat emalinya bukan bagian dari domain Anda atau Gmali tidak dapat memverinkasi banwa pesan tersebut berasai dari ijpris@laescore ii kecuali jika Anda menghubungi pengirimnya dengan cara lain untuk memastikan alamat email ini sah.	.com. Jangan balas	email	
×		Laporkan phishing Kelihatannya aman		(j	
++22		Terjemahkan ke Indonesia × The following message is being delivered on behalf of International Journal of Public Health Science (JPHS). apt widyasari putranti: Thank you for submitting the manuscript, "Formulation of Intra and Extra-Granularly Crosscamellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)" to International Journal of Public Health Science (JPHS). Science (JPHS). wild be able to track its progress through the editorial process by logging in to the journal web site.			
		Manuscript URL: <u>https://fiphs_iaescore.com/index.php/UPHS</u> /author/submission/22666 Username: widyasari			
× -		If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.			

Catatan Penelaahan Awal dan Permohonan Perbaikan Artikel

Q ijphs	× 辈		• Aktif ~	0 \$		UNIVERSITAS AHMAD D	AHLAN	•
					12 da	ıri 23 🔇	: >	
						ć	3 C	
Lina Handayani ijphs@iaescore.com <u>lewat</u> kirimdata.id kepada saya ▼				Jum, 24 Mar 202	3, 06.57	☆ ←	n i	
Hati-hati dengan pesan ini. Lina Handayani mirip dengan sebuah nama di organisasi Anda, tetapi alamat emailnya bu Ini kecuali jika Anda menghubungi pengirimnya dengan cara lain untuk memastikan alami Laporkan phishing Kelihatannya aman	ikan bagian dari domain Anda ata at email ini sah.	ı Gmail tidak dapat memverifikasi	bahwa pesan tersebut berasal dari ijph:	@iaescore.com	n. Jangan	balas em	ail (j)	
Terjemahkan ke Indonesia X The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).								
Dear Prof/Dr/Mr/Mrs: apt widyasari putranti, We have reached a decision regarding your submission entitled "Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)" to International Journal of Public Health Science (IJPHS), a peer-reviewed and an OPEN ACCESS journal that makes significant contributions to major areas of public health science.								
Our decision is to revisions The goal of your revised paper is to describe novel technical results. A high quality paper MUST has: (1) a clear statement of the problem the paper is addressing> explain in "Introduction" section (2) the proposed solution(s)/method(s)/approach(es)/framework(s)/ (3) results achieved. It describes clearly what has been done before on the problem, and what is new.								

Pengirman revisi naskah sesuai catatan penelaahan

	Syzygium polyananam excluse
Section	Other_Topics_in_Public_Health_Science
	Veronique Gucht 🖾 (Review)
Editor	Henry Imhonde 🖾 (Review)
	Jennifer Spencer 🖾 (Review)

Peer Review

Round 1

Version

Review Version	22666-44109-1-RV.DOCX	2022-11-01
Initiated	2022-11-01	
Last modified	2023-03-23	
Uploaded file	None	

Editor Decision

Decision	Accept Submission 2023-05-16				
Notify Editor		Editor/Author E	mail Record 쯱	2023-05-16	
Editor Version	None				
Author	22666-	46084-1-ED.DOCX	2023-05-09	DELETE	
Version	22666-	46084-2-ED.DOCX	2023-05-15	DELETE	
	<u>22666-</u>	46084-3-ED.DOCX	2023-08-14	DELETE	
	<u>22666-</u>	46084-4-ED.DOCX	2023-10-14	DELETE	
	<u>22666-</u>	46084-5-ED.DOCX	2023-10-19	DELETE	
Upload Author	Choo	ose File No file o	hosen	Upload	

Pemberitahuan Penerbitan artikel



Permohonan Penyesuaian Konten Artikel



Pengiriman Revisi artikel, bukti transfer dan form transfer copyright

	۱	widyasari putranti «widyasari@pharm.uad.ac.id» kepada IJPHS ▼	@ 14 Okt 2023, 1	16.58 🍸	ک ک
		Widyasari Putranti. M. Sc. Apt Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770			
		Satu lampiran • Dipindai dengan Gmail ()			
	IJ	IJPHS IAES ⊲ijphsiaes.core@gmail.com> kepada saya ▼	@ 17 Okt 2023, 1	14.48 🕇	γ (
		Terjemahkan ke Indonesia × Thanks for revision. Please revise the following files again.			
	Paym	ent article 22666 (Eksternal) Kotak Masuk ×	:	× 8	Ľ
۲	widyasa kepada ijp Berikut te	iri putranti <widyasari@pharm.uad.ac.id> hs ▼ rlampir payment utk artikel 22666. Utk camera ready paper dan similirity menyusul. Terima kasih</widyasari@pharm.uad.ac.id>	🗢 Jum, 28 Jul 2023, 15.22 🔥	r <	:
	Satu lar Neminar Ir Matode Tra Tujuer Tran Total Trans Total Trans MitDyASJ	mpiran - Dipindai dengan Gmail () watar Patasano atar Patasano			A
IJ	IJPHS Ed kepada sa Baik bu te On Fri, Ju Berikut	ditor ≺ijphs⊜iaescore.com> ya ❤ arima kasih, kami tunggu segera untuk camera ready dan hasil similartynya il 28, 2023 at 3·22 PM widyasari putranti < <u>widyasari@pharm uad.ac.id</u> > wrote: terlampir payment utk artikel 22666. Utk camera ready paper dan similirity menyusul. Terima kasih	Jum, 11 Agu 2023, 10.37 🛛 🗙	r 41	I
	UNIVERS Kampus Kampus Kampus Kampus	SITAS AHMAD DAHLAN 1: Jin. Kapas No. 9 Yogyakarta 2: Ji. Pramuka 42, Sidkan, Umbulharjo, Yogyakarta 55161 3: Ji. Prof. Dr. Sceponno, S. H., Janturan, Warungboto, Umbulharjo, Yogyakarta 55164 4: Ji.Ringroad Selatan, Yogyakarta 5: Ji. Ki Ageng Pemanahan 19, Yogyakarta			

Penerbitan Artikel



DAFTAR ISI CORRESPONDING AUTHOR ARTIKEL JURNAL

"Influence of crascarmellose in fast disintegrating tablet of Syzygium polyanthum extract"

IJPHS, Vol 13 No 1, 2024

No	Keterangan	Tanggal
1	Submitted to the journal "IJPHS"	1 november 2022
2	First revision : accepted with major revision	24 maret 2023
3	Second revision : minor revision (reference)	10 mei 2023
4	Paper accepted	16 mei 2023
5	Author payment APC	28 Juli 2023
6	Author Sent camera ready paper and similarity cek	14 agustus 2023
7	3th revision	10 oktober 2023
8	4 th revision	17 oktober 2023
9	5 th final revision	28 oktober 20223
10	Certificate of acceptance	2 november 2023
11	Published	21 desember 2023

UNIVERSITAS AHMAD DAHLAN

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Submission Acknowledgement

1 pesan

Lina Handayani <ijphs@iaescore.com> Kepada: apt widyasari putranti <widyasari@pharm.uad.ac.id>

The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).

apt widyasari putranti:

Thank you for submitting the manuscript, "Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)" to International Journal of Public Health Science (IJPHS). With the online journal management system that we are using, you will be able to track its progress through the editorial process by logging in to the journal web site:

Manuscript URL: https://ijphs.iaescore.com/index.php/IJPHS/author/submission/22666 Username: widyasari

If you have any questions, please contact me. Thank you for considering this journal as a venue for your work.

Lina Handayani International Journal of Public Health Science (IJPHS)

International Journal of Public Health Science (IJPHS) http://ijphs.iaescore.com 1 November 2022 pukul 13.36

UNIVERSITAS AHMAD DAHLAN

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Editor Decision

1 pesan

Lina Handayani <ijphs@iaescore.com> Balas Ke: "Dr. Lina Handayani" <ijphs@iaescore.com> Kepada: apt widyasari putranti <widyasari@pharm.uad.ac.id>

The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).

Dear Prof/Dr/Mr/Mrs: apt widyasari putranti,

We have reached a decision regarding your submission entitled "Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)" to International Journal of Public Health Science (IJPHS), a peer-reviewed and an OPEN ACCESS journal that makes significant contributions to major areas of public health science.

Our decision is to revisions

The goal of your revised paper is to describe novel technical results.

A high quality paper MUST has:

(1) a clear statement of the problem the paper is addressing --> explain in "Introduction" section

(2) the proposed solution(s)/method(s)/approach(es)/framework(s)/

(3) results achieved. It describes clearly what has been done before on the problem, and what is new.

In preparing your revised paper, you should pay attention to:

1. Please ensure that: all references have been cited in your text; Each citation should be written in the order of appearance in the text; The references must be presented in numbering and CITATION ORDER is SEQUENTIAL [1], [2], [3], [4],

Please download & study our published papers for your references:

- http://ijphs.iaescore.com

- http://ijere.iaescore.com

- http://journal.uad.ac.id/index.php/edulearn

- http://iaescore.com/journals (other journals)

(Please use "Search" menu under "JOURNAL CONTENT" menu in right side of the site)

2 An Introduction should contain the following three (3) parts: - Background: Authors have to make clear what the context is. Ideally, authors should give an idea of the state-of-the art of the field the report is about.

- The Problem: If there was no problem, there would be no reason for writing a manuscript, and definitely no reason for reading it. So, please tell readers why they should proceed reading. Experience shows that for this part a few lines are often sufficient.

- The Proposed Solution: Now and only now! - authors may outline the contribution of the manuscript. Here authors have to make sure readers point out what are the novel aspects of authors work. Authors should place the paper in proper context by citing relevant papers. At least, 5 references (recently journal articles) are used in this section.

3. Results and discussion section: The presentation of results should be simple and straightforward in style. This section report the most important findings, including results of statistical analyses as appropriate. You should present the comparison between performance of your approach and other researches. Results given in figures should not be repeated in tables. It is very important to prove that your manuscript has a significant value and not trivial.

Please submit your revised paper within 6 weeks.

I look forward for hearing from you

Thank you

Best Regards, Dr. Lina Handayani

The following template should be used for responses to reviewers:

I would like to thank the reviewers for their insightful feedback. All comments from Reviewer 1 are highlighted in yellow, those from Reviewer 2 are highlighted in red, and those from Reviewer 3 are highlighted in green.

Reviewer 1

Comment 1: There are some references that are not required. Response: We thoroughly updated our references; 5 references were eliminated, and two were replaced by more recent publications.

Comment 2: The presentation of Figures 2 and 3 should be improved. Response: The necessary adjustments have been made.

Comment 3: Equation (2) seems to be incorrect.

Response: Equation (2) is correct. This can be proven as follows:... In order to clarify equation 9 in the manuscript, the following remarks have been added... etc.

All changes for reviewer 1 are highlighted in yellow in the main text.

Reviewer 2

Comment 1: Response:

Comment 2: Response:

Comment 3: Response:

All changes for reviewer 2 are highlighted in red in the main text.

Etc.

Such a document clarifies everything and will aid the reviewers in evaluating the work fast. When providing your amended primary document files, you must also upload your corrections statement. Before your manuscript, the declaration of revisions should appear.

Reviewer J:

Does the paper contain an original contribution to the field?: Yes

Is the paper technically sound?:

Yes

```
Does the title of the paper accurately reflect the major focus contribution of this paper?:
```

Yes

Please suggest change of the title as appropriate within 10 words:

Is the abstract a clear description of the paper?

.

:

Yes

Please suggest change of the abstract

Is the paper well written (clear, concise, and well organized)?: No

Are the equations, figures and tables in this journal style, clear, relevant, and are the captions adequate?:

No

Please score the paper on a scale of 0 - 10 as per the directions below:

9-10 Excellent - Outstanding 7-8 Good 5-6 Average 3-4 Poor 0-2 Very Poor : 5

Comments to the Authors (how to improve this paper)::

-The template of paper needs a double check to follow all preparation rules and requirements rigidly. See:

https://ijphs.iaescore.com/index.php/IJPHS/about/submissions#authorGuidelines

-Separate citations with commas, For example in your paper, citation [7][8] can be written: [7], [8].

- Simplify multiple citations, for example in your paper [3][4][5][6]., it should be write: [3]-[6].

-Write citation from small number. For example in your paper: [10][6] it should be write: [6], [10].

-If you put citation at the end of the sentence make sure it comes before ".". For example in your paper: "... cholesterol levels.[3][4][5][6]" it should be write: ...cholesterol levels [3]-[6].

CHECK ALL citing references and do the revision. Use Mendeley/reference manager to help you manage the citations.

-In subsection "Weight uniformity test" the Perka BPOM RI can be write in italic style and describe it, since it is using local language. Please note that not all the readers understand your language. Use English so that your idea is understood by all the readers. Or you can inform the "BPOM RI" in nomenclature or explain more in the introduction.

-In second paragraph of introduction, there are abbrevitations: LDL, HDL, HMG -CoA, for the first appearance in a paper, please expand the abbreviation(s)/acronym(s). Also, write explanation first then abbrevitation. For example in your paper: LOD (Loss on Drying), MC (

Moisture Content), write as: loss on drying (LOD), moisture content (MC). CHECK ALL abbrevitation on the whole paper.

-Sub-Figures 1(a)-(b) and 2(a)-(d) are not mentioned in the text. If a figure contains subfigures, explain what the whole figure illustrates before explaining what the subfigures do. This is important to make the readers get what the relation between those subfigures is, and why they are put in the same figure's number instead of making them two different figures in separate numbering.

Complete the description by explaining the relation of the images, why these images should be grouped together in the same image as subfigures, and not split up as several different images.

-The subfigures of 2: (c) sclerenchyma maybe it is (d) sclerenchyma.

-As written in the journal template, table is written with a font size of 8pt, no bold in the row/colum.

-In the method section, re-write "Bay Leaf Ethanol Extract, FDT Formulation of Bay Leaf Extract, Granule Physical Properties Test Parameters, Flowability test, Average diameter test, Fragility test, Absorption test" as sub-section. For example: 2.1.Bay Leaf Ethanol Extract

2.1 Bay Leaf Ethanol Extract

2.2 FDT Formulation of Bay Leaf Extract

2.3Granule Physical Properties Test Parameters.

... etc

-Some references are in local language, translate it. As this is an international journal, readers are from all over the world, not all the readers understand your language. Use English so that your idea is understood by all the readers.

-Elaborate the biographies into min. 3 sentences. It should include: the links of the researchers social media accounts of the authors as complete as possible: Scopus (if available), ORCID (required), Google.

Reviewer K:

Does the paper contain an original contribution to the field?: Yes

Is the paper technically sound?: Yes

Does the title of the paper accurately reflect the major focus contribution of this paper?: Yes

Please suggest change of the title as appropriate within 10 words:

Is the abstract a clear description of the paper?

Yes

Please suggest change of the abstract

Is the paper well written (clear, concise, and well organized)?: Yes

Are the equations, figures and tables in this journal style, clear, relevant, and are the captions adequate?:

Yes

Please score the paper on a scale of 0 - 10 as per the directions below:

9-10 Excellent - Outstanding 7-8 Good 5-6 Average 3-4 Poor 0-2 Very Poor : 6

Comments to the Authors (how to improve this paper)::

The paper has been written quite well. However, there are some things that need to be improved to improve the quality of the paper: The references should be writen in international language (English)

Reviewer L:

Does the paper contain an original contribution to the field?: Yes

Is the paper technically sound?: Yes

Does the title of the paper accurately reflect the major focus contribution of this paper?:

Yes

Please suggest change of the title as appropriate within 10 words: Not Needed. Title is clear

Is the abstract a clear description of the paper?

Yes

:

•

Please suggest change of the abstract

Not needed

Is the paper well written (clear, concise, and well organized)?: Yes

Are the equations, figures and tables in this journal style, clear, relevant, and are the captions adequate?:

Yes

Please score the paper on a scale of 0 - 10 as per the directions below:

9-10 Excellent - Outstanding 7-8 Good 5-6 Average 3-4 Poor 0-2 Very Poor : 8

Comments to the Authors (how to improve this paper):: Paper is well structured.

Ref. from Journal article must be completed with vol., issue, pages, DOI

- Update references in recent 10 years

- Cite references in IEEE Style, not APA Style

- Write biographies of authors after ref. section

- Complete the ORCID ID for each author in Biographies section.

-Make sure that each paragraph at least contained three sentences.

- Each reference must be completed with DOI and can be traced online.

- Similarity should be no more than 20 percent.

- Proof read the English to expert.

- State the research funding and its contract number, if any in the

acknowledgment section

International Journal of Public Health Science (IJPHS) http://ijphs.iaescore.com

UNIVERSITAS AHMAD DAHLAN

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Editor Decision

1 pesan

Lina Handayani <ijphs@iaescore.com> Balas Ke: "Dr. Lina Handayani" <ijphs@iaescore.com> Kepada: apt widyasari putranti <widyasari@pharm.uad.ac.id>

The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).

Dear Prof/Dr/Mr/Mrs: apt widyasari putranti,

some references are old and missed the DOI.

-Make sure that each paragraph at least contained three sentences. - Add implication of your research in Conclusion section.

International Journal of Public Health Science (IJPHS) http://ijphs.iaescore.com 10 Mei 2023 pukul 15.17

UNIVERSITAS

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Editor Decision

1 pesan

Lina Handayani <ijphs@iaescore.com> Balas Ke: "Dr. Lina Handayani" <ijphs@iaescore.com> Kepada: apt widyasari putranti <widyasari@pharm.uad.ac.id>

The following message is being delivered on behalf of International Journal of Public Health Science (IJPHS).

-- Paper ID# 22666 -- Authors must strictly follow the guidelines for authors at http://iaescore.com/gfa/ijphs.docx

Dear Prof/Dr/Mr/Mrs: apt widyasari putranti,

It is my great pleasure to inform you that your paper entitled "Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)" is ACCEPTED and will be published on the International Journal of Public Health Science (IJPHS). This journal is accredited SINTA 1 by Ministry of Research and Technology/National Research and Innovation Agency, Republic of Indonesia (RISTEK-BRIN) and has ACCEPTED for inclusion (indexing) in Scopus (https://suggestor.step.scopus.com/progressTracker/?trackingID=D331D503BA1584BF) since 2020 issues (https://www.scopus.com/results/results.uri?src=s&st1=&st2=&sot=b&sdt=b&origin=searchbasic&rr=&sl=57&s= SRCTITLE%20(International%20Journal%20of%20Public%20Health%20Science). Congratulations!

Please prepare your final camera-ready paper (in MS Word or LATEX file format) adheres to every detail of the guide of authors (MS Word: http://iaescore.com/gfa/ijphs.docx, or http://iaescore.com/gfa/ijphs.rar for LATEX file format), and check it for spelling/grammatical mistakes. Then you should upload your final paper though our online system (as "author version" under our decision, NOT as new submission).

You should submit your camera-ready paper along with your payment receipt and similarity report (that less than 20%) within 6 weeks.

I look forward to hearing from you.

Thank you

Best Regards, Dr. Lina Handayani

You should submit all your documents:

to email: jphs@iaescore.com within 6 weeks.

All correspondence should be addressed to the emails (support by phone is not provided).

16 Mei 2023 pukul 12.30

^{1.} Camera-ready paper

^{2.} Similarity report, along with

^{3.} Your payment evidence

IMPORTANT!!!

1). PLEASE ADHERE STRICTLY THE GUIDE OF AUTHORS !! http://iaescore.com/gfa/ijphs.docx and pay attention to the checklist for preparing your FINAL paper for publication: http://ijphs.iaescore.com/index.php/IJPHS/about/editorialPolicies#custom-5

2). It is mandatory to present final paper in the sections structure "IMRADC style":

1. INTRODUCTION

2. The Proposed Method/Framework/Procedure specifically designed (optional)

3. METHOD

4. RESULTS AND DISCUSSION

5. CONCLUSION

See http://iaescore.com/gfa/ijphs.docx

3). It is mandatory!! Add biographies of authors as our template (include links to the authors' profiles, do not delete any icons in the template). See http://iaescore.com/gfa/ijphs.docx

--> Provide links for all authors to the 4 icons (Scholar, Scopus, Publons and ORCID)

4). Use different PATTERNS for presenting different results in your graphics (instead of different colors). It is mandatory!! Re-check all your figures. See http://iaescore.com/gfa/ijphs.docx

5). Please ensure that all references have been cited in your text. Use a tool such as EndNote, Mendeley, or Zotero for reference management and formatting, and choose IEEE style. Each citation should be written in the order of appearance in the text in square brackets. For example, the first citation [1], the second citation [2], and the third and fourth citations [3], [4]. When citing multiple sources at once, the preferred method is to list each number separately, in its own brackets, using a comma or dash between numbers, as such: [1], [3], [5]. It is not necessary to mention an author's name, pages used, or date of publication in the in-text citation. Instead, refer to the source with a number in a square bracket, e.g. [9], that will then correspond to the full citation in your reference list. Examples of in-text citations:

This theory was first put forward in 1970 [9]. Zadeh [10] has argued that ... Several recent studies [7], [9], [11]-[15] have suggested that.... ... end of the line for my research [16].

6). Please present all references as complete as possible and use IEEE style (include information of DOIs, volume, number, pages, etc). If it is available, DOI information is mandatory!! See http://iaescore.com/gfa/ijphs.docx

In order to cover part of the publication cost, each accepted paper is charged: USD 265 (~IDR 3850K). This charge is for the first 8 pages, and if any published manuscript over 8 pages will incur extra charges USD 50 (~IDR 800K) per page. Your paper is 9 pages.

The payment should be made by bank transfer (T/T):

Bank Account name (please be exact)/Beneficiary: LINA HANDAYANI Bank Name: CIMB NIAGA Bank Branch Office: Kusumanegara Yogyakarta City: Yogyakarta Country : Indonesia Bank Account #: 760164155700 -----

SWIFT Code: BNIAIDJAXXX (PT. BANK CIMB NIAGA, TBK. in JAKARTA)

or as an alternative of the bank transfer, through PayPal to email: info@iaesjournal.com

International Journal of Public Health Science (IJPHS) http://ijphs.iaescore.com

UNIVERSITAS

widyasari putranti <widyasari@pharm.uad.ac.id>

Payment article 22666

4 pesan

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: ijphs@iaescore.com 28 Juli 2023 pukul 15.22

Berikut terlampir payment utk artikel 22666. Utk camera ready paper dan similirity menyusul. Terima kasih



IJPHS Editor <ijphs@iaescore.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id> 11 Agustus 2023 pukul 10.37

Baik bu terima kasih, kami tunggu segera untuk camera ready dan hasil similartynya

On Fri, Jul 28, 2023 at 3:22 PM widyasari putranti <widyasari@pharm.uad.ac.id> wrote: Berikut terlampir payment utk artikel 22666. Utk camera ready paper dan similirity menyusul. Terima kasih

UNIVERSITAS AHMAD DAHLAN Kampus 1: Jln. Kapas No. 9 Yogyakarta Kampus 2: Jl. Pramuka 42, Sidikan, Umbulharjo, Yogyakarta 55161 Kampus 3: Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta 55164 Kampus 4: Jl.Ringroad Selatan, Yogyakarta Kampus 5: Jl. Ki Ageng Pemanahan 19, Yogyakarta

Kontak

Email: info@uad.ac.id Telp. : (0274) 563515, 511830, 379418, 371120 Fax. : (0274) 564604

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS Editor <ijphs@iaescore.com>

berikut kami lampirkan camera ready paper dan cek similirity <u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

[Kutipan teks disembunyikan]

14 Agustus 2023 pukul 14.46

2 lampiran

Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Ex.pdf 3189K

22666-46084-2-ED_destyrr_v1.docx 4929K

IJPHS Editor <ijphs@iaescore.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id> 6 September 2023 pukul 09.21

Received, thank you [Kutipan teks disembunyikan]

Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)

Submission date: 11-Aug-2023 03:14PM (UTC+0700) Submission ID: 2144347899 File name: 22666-46084-2-ED_destyrr_v1_1.docx (4.81M) Word count: 5245 Character count: 28789 **International Journal of Public Health Science (IJPHS)** Vol. 99, No. 1, Month 2099, pp. 1~1x ISSN: 2252-8806, DOI: 10.11591/ijphs.v99i1.paperID

Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)

Widyasari Putrant 13 Desty Restia Rahmawati¹, Nining Sugihartini¹, Teuku Nanda Saifullah² ¹Faculty of Pharmacy, Ahmad Dahlan University, Yogyakarta, Indonesia ²Faculty of Pharmacy, Gadjah mada University, Yogyakarta, Indonesia

Article Info

Article history:

Received month dd, yyyy Revised month dd, yyyy Accepted month dd, yyyy

Keywords:

Bay Leaf Extract Fast Disintegrating Tablet Croscarmellose sodium Extra granular Intra granular

ABSTRACT

Bay leaves contain the flavonoid quercetin which can be used as an antihype12 idemic drug. The development of antihyperlipidemic drug formula in the form of Fast Disintegrating Tablet (FDT) is needed for patients who experience dysphagia. FDT preparations require 12 optimal super disintegrant concentration to produce a good drug formula. This study aims to develop the FDT formula for 70% ethanol extract of bay leaves using the super disintegrant croscarmellose sodium (CCS) intra and extra-granular. FDT Formulation using the wet granulation method with variations in CCS concentrations, namely F1: 2%, F2: 3.5%, and F3: 5% (extra granular), while 2% (intra granular). The formulation process, in-process control (IPC) granules, weight uniform 28 tests, and various physical properties tests of tablets were carried out. Data were statistically analyzed using One Way ANOVA. test (a=95%). The results of statistical tests of IPC granules, uniformity of weight, and tablet size of all FDT formulas were not significantly different (p> 0.05). CCS concentration difference extra granular significantly affected the wetting time, disintegration time, hardness, and the value of % friability of FDT (p< 0.05). The combination of intra and extra granular CCS (2%:5%) gave the most optimum physical properties of bay leaf extract FDT.

This is an open-access article under the CC BY-SA license.



Corresponding Author:

Widyasari Putranti Faculty of Pharmac 2 Ahmad Dahlan University, Yogyakarta, Indonesia Email: Widyasari@pharm.uad.ac.id

1. INTRODUCTION

Plant germplasm resources as raw materials for medicines 21 e of which is a bay leaf (*Syzygium polyanthum*). Bay leaves are known to contain secondary metabolites, such as saponins, terpenoids, 19 vonoids (quercetin), polyphenols, alkaloids, steroids, and essential oils (sesquiterpenes)[1]-[2]. Based on the Decree of the Minister of Health of the Republic of Indonesia (2009), bay leaves contain a total flavonoid of not less than 0.40% which is calculated as quercetin[3]. Querc 24 in bay leaves the potential as an antihyperlipidemic which has been shown to be able to significantly reduce levels of triglycerides, total cholesterol, and anLow-Densityty Lipoprotein (LDL) cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in High-Density Lipoprotein (HDL), as well as inhibiting 3-hidroksi-3-metilglutaril coenzyme A (HMG -CoA) activity, reductase, and LDL oxidation.[4]-[5]

Conventional drugs for hyperlipidemia are generally available in tablet form. However, the tablet dosage form has several disadvantages. Some of them require a long absorption time in the body, slow drug action, and elderly patients may experience difficulty swallowing or dysphagia[6]. Seeing the problems that

Journal homepage: http://ijphs.iaescore.com

¢

ISSN: 2252-8806

2

0

arise from conventional tablets, the herbal Fast Disintegrating Tablet (FDT) of bay leaf extract is the right alternative so that the drug can be used comfortably and has a therapeutic effect more quickly. FDT is a solid dosage form containing pharmaceutically active substances that can be disintegrated quickly, generally within seconds when placed on the tongue, so that FDT can release the active drug substance immediately, speed up the onset, increase the oral bioavailability of the drug in the body so that therapeutic effect more quickly. FDT preparations combine 2 advantages of preparations, namely liquid which can increase the solubility and bioavailability of drugs, as well as the advantages of solid dosage forms that have high physicochemical stability, relatively constant homogeneity of the active ingredients of the preparation, and easier manufacturing.

FDT is formulated with croscarmellose sodium (CCS) super disintegrant which is classified as a strong swelling product with a good wa 11 absorption mechanism[8],[9]. CCS is effectively used in FDT preparations because it rapidly expands to 4-8 times its initial volume when in contact with water, thus accelerating the tablet-crushing process. CCS is effective intra-granular which allows for better capillary and swelling processes, causing the tablet to disintegrate quickly and reduce disintegration time[9]. The addition of intra and extra-granular crushing materials can increase the effectiveness of tablet disintegration through two stages, namely extra granular which breaks the tablet into granules, and intra granular breaks down the granules into smaller particles, with the mechanism of the crushing material that experiences rapid swelling and water absorption (wicking) through the gaps between the granules and the particles that make up the granules[10]. This study aims to develop herbal fast-disintegrating tablets (herbal FDT) with the active ingredient of bay leaf ethanol extract. The concentration of CCS super disintegrant intra and extra-granular needs to be determined to obtain the FDT of bay leaf extract with optimum physical properties.

2. METHOD

The ingredients used were simplicia and bay leaf extract (Beringharjo Market, Yogyakarta Indonesia) in the form of fresh bay leaf simplicia, ethanol, croscarmellose sodium (PT. Phapros Tbk), lactose, magnesium stearate, aspartame, menthol and corn starch paste. The tools used are Ohaus Analytical Balance, Absorption test equipment granules, Volumenometer (Tapped Density Tester), Flow Tester granule, Oven, Rotary evaporator calipers, Sieving Machine, Vacuum Pump, Dry Granule Sieve, Wet Granule Sieve, Pan, Halogen Moisture Analyzer, Single Punch Tablet Printing Machine, Hardness Tester, Friability Tester, and Disintegration Tester.

2.1 Bay Leaf Ethanol Extract

Bay leaves were extracted by maceration method using 70% ethanol solvent with a sample: solvent ratio (1:10) (w/v)[3]. Maserati and pulp were separated by the vacuum pump and filter paper. The solvent was evaporated using a rotary evaporator. The viscous extract obtained was calculated for its yield.

2.2 FDT Formulation of Bay Leaf Extract

FDT was made by the wet granulation method. At the granulation stage, the process of mixing, extracting, super disintegrant CCS (intra granular), lactose, and corn starch. The mixture was sieved through a wet (No. 18 mesh), oven-dried (50-60°C), then sifted dry (No. 20 mesh). Extra granular granules and excipients consisting of magnesium stearate, aspartame, menthol, and CCS (extra granular) mixed until homogeneous (30 minutes). Mixture Extra granular granules and excipients that have been homogenized are compressed using a single punch tablet press.

2.3 Granule Physical Properties Test Parameters Moisture content (MC) test

The moisture content of granules can be determined based on the value of LOD (Loss on Drying) and MC. LOD is a test for measuring the difference in heavy total granules when before and after drying or a moisture content statement based on wet weight. Meanwhile, MC is a statement of moisture content based on dry weight. Measurements using the Halogen Moisture Analyzer tool. Good granule moisture content if the LOD and MC values are 10%.[11]

Flowability test

The flow properties of the granules can be tested by calculating the angle of repose, the flow time test, and the determination index (Hausner ratio). The granule angle of repose is said to be very good (25 -30); good (30-35°) or $25^{\circ} > < 35$. A good granule flow time is not more than 10 seconds for 100 grams of granules. Excellent granule setting index (\leq 10); good (11–15), and the Hausner ratio index that meets the requirements for the physical properties of the granule mass, which is very good (1.00–1.11); good (1.12–1.18).[12] **Average diameter test**

Int J Public Health Sci

3

ISSN: 2252-8806

Sieve No. 14, 16, 20, 30, 50 mesh and pan, installed in stages on the sieving machine (50 amplitude, 15 minutes). The granules left on each sieve were weighed and the percentage was calculated.[11] **Fragility test**

The fragility of the granules is based on the number of fines that occur after the friability test by sieving[13]. The granules were put into a graded sieve with the top sieve No. 30 mesh and bottom pan, using a sieving machine (50 amplitude, 30 minutes).

Absorption test

The water absorption of the granules affects the tablet's disintegration time (disintegration). It was carried out with a series of absorption test equipment by placing 300 mg of granules on a water-saturated sheath paper. The absorbed water was measured using a digital scale for 15 minutes or until it was constant.[8]

2.4 Physical Properties Test Parameters of FDT Weight uniformity test

Requirements and weight uniformity tests are determined based on the regulation of the National Food and Drug Agency of the Republic of Indonesia concerning the quality requirements of traditional medicines. The test was carried out by weighing 20 tablets one by one, from 20 tablets no more than 2 tablets, each of which deviated from the average weight by more than the price specified in column A and not one tablet whose weight deviated from the average weight. the average is greater than the price specified in column B.[14] Size uniformity test (diameter and thickness)

Each tablet was measured in diameter in a horizontal position with a caliper, and the thickness of the the was measured in a vertical position with a caliper (millimeters).

Hardness test

The tablet hardness test was carried out by taking a sample of at least 6 tablets from each formula. The tablet hardness test is carried out using hardness tester tablets. One by one the tablets are tested by placing the tablets in the center perpendicular to the hardness tester. A good FDT hardness is 3-5 kg/cm².[12],[15] **Fragility test**

For tablets with a single weight of exactly or less than 650 mg, the total weight of the tested samples was close to 6.5 grams. Prior to testing, the tablets were dusted and weighed. All tablets were put into the friability tester (speed 25 rpm, 4 minutes). The tablets were cleaned of adhering fines and re-weighed. A good friability [3] he should not be more than 1%.[12]

Wetting time test

The wetting time test was done by placing a sheet of filter paper that has been folded twice into a petri dish with a 10 meter of 5 cm. The petri dish was filled with 5.0 mL of distilled water containing the strawberry red dye. A tablet is then 14 heed on the filter paper, and simultaneously with the start of the test instrument, the stopwatch is turned on. Wetting time was calculated as the time required for a red color to appear on the entire surface of the tablet.[16]

Disintegration time 27st (disintegration)

A total 6 tablets of each formula were placed in each tube and a disk was placed on it, the test was carried out with water medium at a temperature of 37 ± 2 °C or 35-39 C using a 1000 mL glass beaker. The disintegration time requirement for FDT preparations is at least < 1 minute disintegration time.[12],[17]

2.5 Data analysis

The results of the physical examined on f granules and tablets of FDT bay leaf extract were analyzed using the One Way ANOVA test method with a 95% confidence level.

3. RESULTS AND DISCUSSION

Bay leaf simplicia was tested for macroscopic (organoleptic) specifications on fresh and dried simplicia, as well as micro opic tests for simplicia parts and identification fragments of simplicia powder. The simplicia specification test was carried out to ensure the correctness of the identity and quality of the bay leaf simplicia based on the Indonesian Herbal Pharmacopoeia compendia and Indonesian Materia Medika. Based on the microscopic test results on figure 1 there are identifier fragments namely (a) lower epidermis with stomata, (b) upper epidermis, (c)elements with dots, (d) prism-shaped calcium oxalate crystals, and (e) sclerenchyma. The results of the specification test for the simplicia of bay leaves showed that the simplicia at the Biology Laboratory of Ahmad Dahlan University (UAD) was carried out to ensure that the selected bay leaf simplicia was in accordance with its species or identity.[3]

Paper's should be the fewest possible that accurately describe ... (First Author)

¢

4

ISSN: 2252-8806



0

Figure 1. identification fragments of bay leaf; (a) lower epidermis, (b) upper epidermis, (c) floem, (d) calcium oxalate crystal, (e) sclerenchyma

3.1 Extraction Bay Leaf

Fresh simplicia of bay leaves were sorted wet, dried using an oven (50-60°C) with a moisture content of not more than 10%, then sorted dry. Simplicia powder for extract preparation is a fine Simplicial (No. 60 mesh) in order to optimize the extraction process[18]. The results of the simplicial extraction of bay leaves weighing 3.7 kg obtained a thick extract of 347 grams. The total yield of the thick extract obtained was 9.38%. The morphology of the viscous extract is shown in Figure 2. Screening at 20 etermination of the chemical compound flavonoid quercetin contained in the thick extract of bay leaves was carried out at the Integrated Research Laboratory of the Faculty of Pharmacy, UAD. According to the N29 stry of Health (2017), bay leaves contain a total flavonoid of not less than 0.40% calculated as quercetin. Based on the results of the test the flavonoid content of 0.8079 \pm 0.0045%, so that it meets the specifications for the total flavonoid content of bay leaves.[18]



Figure 2. Bay leaf Extract

3.2 FDT Formulation

FDT formulation method with wet granulation is got 7 sed for water-resistant active ingredients and heating temperature[19]. The active ingredient in bay leaf is quercetin (3,3',4',5,7-pentahydroxyflavone) is a flavonoid derivative compound that is resistant to heating with a melting point of 316.5 °C [20], while the wet granulation process ranges from at a temperature of 40-60°C. The wet granulation method can improve the flowability and compactibility of the material and facilitate the FDT formulation process from the active ingredient in the form of a thick extract[21]. Wet granulation has the advantage of good content distribution and uniformity, besides that the hydrophobic surface of the tablet becomes more hydrophilic which affects the water absorption process into the tablet, thereby increasing the speed of tablet disintegration[22]. The work of woven fibers between granules can increase the speed of tablet disintegration[23], it is synergistic with the presence of CCS super disintegrant extra granular which facilitates swelling and absorption of water into granules and their small particles.

The concentration of CCS super disintegrant intra-granular of the three formulas was the same, ie 2% (10 mg/tablet), and extra granular was formulated with varying levels, F1 2% (10), F2 3.5% (17.5), and F3 5%

Int J Public Health Sci, Vol. 99, No. 1, Month 2099: 1-1x

Int J Public Health Sci

5

ISSN: 2252-8806

(25 mg/tablet). The FDT formula for bay leaf extract is shown in Table 1. Based on the Handbook of pharmaceutical excipients the use of CCS in the wet granulation process should be added in the wet or granulation (intra-granular) stage, as well as in the dry (extra-granular) process[9],[24]. Because this method can increase the wicking ability (water absorption) and swelling of the CCS super disintegrant.[9]

Ingradiant	Composition (mg)		
Ingredient	F1	F2	F3
134 leaf extract	70	70	70
Croscarmellose sodium (intra granular)	10	10	10
Lactose	380	374	366
Croscarmellose sodium (extra granular)	10	17.5	25
Magnesium stearate	5	5	5
Aspartame	20	20	20
Menthol	1	1	1
Corn starch paste 5% w/v	qs	qs	qs
Total Weight (mg)	500	500	500

3.2.1 Granule Physical Properties Test Parameters

MC and LOD values 19 granules from all formulas (table 2), the results obtained meet the requirements of 10% (p>0.05). The flow properties of the granules show the results of the angle of repose of the granules ranging from 30-35 (good), granule flow time of 10-14 grams/second (good), and the determination test/Hausner ratio (very good). The flow properties of granules are very influential on the tablet compression process, granules with good flow properties will cause the granules to enter the die space to be relatively constant so that the tablet weights with small weight variations can be obtained and can increase the uniformity of the resulting tablet dosage.[8],[25]

Table 2. Results of Examination of Physical Properties of FDT Granules of Bay Leaf Extract

m . D	Formula			
Test Parameters	F1	F2	F3	
LOD ± SD (%)*	7.03±0.76	6.29±1.28	6.57±1.18	
MC ± SD (%)*	2.98±0.17	2.35±0.33	2.69±0.31	
Angle of repose ± SD (°)*	30.66±1.75	31.18±2.92	30.08±2.30	
Flow time ± SD (grams/second)*	12.98±1.28	13.12±0.89	11.85±0.48	
Assignment ± SD (%)*	3.17±1.04	2.83±1.04	3.83±0.76	
Hausner ratio Index ± SD*	1.03 ± 0.01	1.03±0.01	1.04 ± 0.01	
Average diameter \pm SD (μ m)*	694.64±28.87	657.58±39.65	698.14±26.09	
Fragility ± SD (%)*	0.8028 ± 0.20	0.8199±0.05	0.5863±0.04	
Absorption ± SD (mg/min)*	0.186±0.05	0.188±0.05	0.127±0.01	

Information:

(*) sig value. (p>0.05) means that the three formulas are not significantly different

The average diameter of all granules ranged from 650-700 m (p> 0.05) and the value of % friability of the granules ranged from 0.5-0.9% (p> 0.05). The granule absorption test for all formulas had absorption between 0.120-0.190 mg/minute (p> 0.05). The granule absorption shows the speed of the granules in absorbing water to crush the granules into small particles so as to speed up the tablet disintegration time. All In Process Control (IPC) granule parameters have met the requirements so that the tableting process can be continued with a single punch tablet press with a punch pressure of 13 kg.

3.2.2 Physical Properties Test Parameters of FDT

The uniformity of tablet weight is related to the uniformity of active substance levels and therapeutic effects. The uniformity of tablet weight (table 3) can be seen from the coefficient of variation F1, F2, are F3 all <5%, (0.0942, 0.1246, and 0.2103) %, respectively. The weight uniformity test based on the National Food and Drug Agency of the Republic of Indonesia requirements, shows the results that meet the requirements, where the tablet meets the specified weight deviation limit value range.

Hardness ranges FDT be good tablet from 3-5 kg/cm 2[15], while the conventional tablet, is 4-8 kg/cm². The FDT hardness (table 3) shows results that meet the requirements. Hardness affects the brittleness and disintegration time of tablets, as well as the dissolution/release of the active drug substance[9]. The results of statistical analysis of tablet hardness with sig. (p<0.05) indicates that the hardness of the three FDT formulas is significantly different. The hardness of FDT tablets is strongly influenced by the CCS super disintegrant excipient. FDT (F3) produced the hardest tablet with the highest extra-granular CCS concentration (5%),

¢

ISSN: 2252-8806

causing the particles in the tablet to be tightly bound. Croscarmellose sodium (Ac-Di-Sol) is a super disintegrant with a fibrous cross-linked polymer structure that functions as a disintegrant with a strong swelling mechanism, fiber is able to bind strongly to the particles so as to increase the hardness and reduce the brittleness of the tablet[26]. This causes the tablet to have good compatibility, so there is a tendency to increase hardness as the amount of CCS in the tablet increases[27]. Another factor that affects the hardness of FDT is the upper punch pressure during the tableting process, the greater the pressure used, the harder the FDT will be. The results of the diameter and thickness measurements of the tablets showed that the size of the entire tablet was uniform (homogeneous).

0

The frability value of the tablet is based on the requirements, which should not be more than 33%[12]. The fragility value of FDT is shown in Table 3 with a friability value of <1% so that FDT meets the requirements. The results of the tablet friability statistical test have a sig value. (p <0.05) means that the fragility value between formulas is significantly different. CCS super disintegrant is a factor influencing fragility. Formula 3 with the highest number of CCS has the smallest friability value, where there is a term of the tablet friability value as the amount of CCS in the tablet increases. The friability value is inversely proportional to the tablet's hardness, where the harder a tablet is, the smaller the friability value will be.

BDT wetting time is shown in Table 3, FDT (F3) with the highest number of CCS has the smallest wetting time. Wetting time is a parageter to determine the speed of FDT in absorbing water, which affects the speed of tab to lisintegration. The faster the wetting time, the faster the tablet disintegration. The wetting time of tablets is influenced by the structure of the tablet matrix and the hydrophilicity of the excipients [28]. The result of statistical analysis of FDT wetting time has a sig value. (p < 0.05) means that the formulas differ significantly. FDT (F3) with the highest number of CCS had the fastest time for wettable tablets, super disintegrant CCS being the most influential factor. The cross-linked chemical structure of CCS creates excipients that have hydrophilic properties and very easy-to-absorb solvents resulting in outstanding swelling properties in the disintegration process.

Table 3. Results of Examination of Physical Properties of Bay Leaf Extract FDT Tablets

Terr		Formula		Note.
Test Parameters	F1 F2		F3	
Weight uniformity ± SD (mg)	$50.4.42 \pm 0.48$	$50\ 4.78\ \pm\ 0.63$	504.43 ± 1.06	Appropriate *
Diameter ± SD (mm)	12.00 ± 0.00	12.00 ± 0.00	12.00 ± 0.00	Appropriate *
Thickness ± SD (mm)	3.67±0.0 0	3.67±0.0 0	3.68 ±0.0 1	Appropriate *
Hardness ± SD (kg)	4.46±0.25 °	4.50±0.27 °	5.24±0.32 a.b	In accordance
Fragility ± SD (%)	0.77±0.05 °	0.72±0.04 °	0.52±0.04 a.b	In accordance
Wetting time ± SD (seconds)	50.33 ± 3.06 °	43.33 ± 4.04 °	20.00 ± 2.65 ^{a,b}	In accordance
Disintegration time ± SD (seconds)	45.33±5.86 °	38.33±3.79 °	19.67±3.06 a.b	In accordance

Information:

(*) sig value. (p>0.05) means that the three formulas are not significantly different

(*) means a formula that 16 gnificantly different from the formula 1

(b) means a formula that is significantly different from the formula 2

(c) means a formula that is significantly different from formula 3;

F1: CCS intra granular 2% : extra granular 2%

F2: CCS intra granular 2% : extra granular 3.5%

F3: CCS intra granular 2% : extra granular 5%

Disintegration (disintegration) time FDT has at least a disintegration time of less than 1 minute[17]. Another source says that in-vitro about 30 seconds or less[29], whereas according to Ph. euros. is <3 min in the oral cavity before swallowing[30],[31]. The faster disintegration time will increase the speed of drug release from the tablet which affects the effectiveness of therapy. The results of the disintegration time of FDT in table 3 show that it meets the requirements of the disintegration time of FDT (<1 minute). FDT disintegration time test are shown in table 3 with FDT (F3) having the fastest disintegration time, due to the higher levels of CCS used in F3 than F1 and F2. The results of the statistical test of tablet disintegration time have a sig value. (p <0.05) means that each formula is significantly different. According to Rowe (2009), CCS has a dual mechanism, namely water wicking and rapid swelling[9]. The highly porous form of the structure speeds up the disintegration time, because water quickly enters the tablet and increases the tablet wetting rate, through the gaps between the granules and the granule pores[32]. This is related to the factor of the amount of CCS super disintegration used in each formula intra and extra-granular. The use of intra and extra-granular affects the decrease in tablet disintegration time through the swelling mechanism, and along with the increase

Int J Public Health Sci, Vol. 99, No. 1, Month 2099: 1-1x

6

Int J Public Health Sci

7

ISSN: 2252-8806

in the extra-granular CCS concentration, the disintegration time will be faster. The FDT disintegration pattern is also facilitated by intra-granular CCS which causes the tablets to be finely crushed not in the form of granules but into small particles so that the speed of disintegration is optimal. The optimum formula for FDT of bay leaf extract is shown in Table 4 with the results of all evaluations of the physical properties of the granules and FDT (F3) that meet the requirements and show the optimum results of physical parameters.

Table 4 . Various Parameters of C	Optimized Tablet Formula ((F3)
-----------------------------------	----------------------------	------

No.	Test Parameters 22	Optimized Formula Value
1.	Weight uniformity ± SD (mg)	504.43 ± 1.06
2.	Diameter ± SD (mm)	12.00 ± 0.00
3.	Thickness ± SD (mm)	3.68 ±0.01
4.	Violence ± SD (kg)	5.24±0.32
5.	Fragility ± SD (%)	0.52±0.04
6.	Wetting time ± SD (seconds)	20.00 ± 2.65
7.	Disintegration time ± SD (seconds)	19.67±3.06

1

The results of this study are in line with the research of Puri et al.[25] and Ainurofiq & Azizah[33], that the combination of intra-granular and extra-granular can produce tablets with the most effective disintegration time by providing better tablet physical properties, compared to intra-granular or extra-granular only. Extra-crushing material Granules have a high tendency to absorb water from the surrounding liquid through the tablet-breaking mechanism[34]. Tablets with disintegrant intra-granular have a higher hardness than tablets with extra granular crushing agents [33]. By combining intra and extra granular super disintegrants, a good level of tablet hardness can be obtained so that the tablet is not brittle and has a faster disintegration time with the mechanism of tablet disintegration into granules (extra granular factor) and granules into small particles (intra granular factor).

4. CONCLUSION

The combination of CCS intra granular 10mg (2%) and extra granular 25mg(5%) can produce FDT with the most effective. In addition, super disintegrant CCS extra granular can affect the physical properties of FDT.

ACKNOWLEDGEMENTS (10 PT)

This research was carried out with the assistance of the Higher Education Research Grant No PKPT-013/SKPP/III/2018.

REFERENCES

- L. Hartanti, S. M. K. Yonas, J. J. Mustamu, S. Wijaya, H. K. Setiawan, and L. Soegianto, 'Influence of extraction methods of bay leaves (Syzygium polyanthum) on antioxidant and HMG-CoA Reductase inhibitory activity', *Heliyon*, vol. 5, no. 4, p. e01485, Apr. 2019, doi: 10.1016/j.heliyon.2019.e01485.
- [2] E. Kato, R. Nakagomi, M. D. P. T. Gunawan-puteri, and J. Kawabata, 'Identification of hydroxychavicol and its dimers, the lipase inhibitors contained in the Indonesian spice, Eugenia polyantha', *Food Chem.*, vol. 136, no. 3–4, pp. 1239–1242, 2013, doi: 10.1016/j.foodchem.2012.09.013.
- [3] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia, 1st ed. Jakarta Indonesia, 2009.
- [4] K. Zeka, K. Ruparelia, R. Arroo, R. Budriesi, and M. Micucci, 'Flavonoids and Their Metabolites: Prevention in Cardiovascular Diseases and Diabetes', *Diseases*, vol. 5, no. 3, p. 19, 2017, doi: 10.3390/diseases5030019.
- [5] W. Putranti, L. Widiyastuti, M. S. Bachri, and Defianti, 'Test Activity Combinations of Celery Herb (*Apium graveolens* L.) and Bay Leaf (*Syzygium polyanthum* W.) Ethanol Extract Toward DecreasedLipid Profile Levels in Hypertensive Mice', *Adv. Mater. Res.*, vol. 1162, pp. 166–172, 2021, doi: 10.4028/www.scientific.net/amr.1162.166.
- [6] I. Safitri, S. Sulistiyaningsih, and A. Y. Chaerunisaa, 'Review : Superdisintegran dalam Sediaan Oral', *Farmasetika.com* (Online), vol. 4, no. 3, p. 56, 2019, doi: 10.24198/farmasetika.v4i3.22945.
- [7] A. Masih, A. Kumar, S. Singh, and A. K. Tiwari, 'FAST DISSOLVING TABLETS: A REVIEW', Int. J. Curr. Pharm. Res., vol. 9, no. 2, p. 8, Mar. 2017, doi: 10.22159/ijcpr.2017v9i2.17382.
- [8] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, 'Formulasi Fast

Paper's should be the fewest possible that accurately describe ... (First Author)

¢

ISSN:	2252-8806

8

Disintegrating Tablet Ekstrak Etanol Daun Salam dengan Kombinasi Crospovidone dan Croscarmellose Sodium sebagai Superdisintegrants', *J. Sains Farm. Klin.*, vol. 8, no. 3, p. 285, 2021, doi: 10.25077/jsfk.8.3.285-295.2021.

[9] R. Rowe, P. Sheskey, and M. Quinn, *Handbook of Pharmaceutical Excipients*, vol. E.28. 2009.

- [10] M. O. Okekunle, O. D. Akin-Ajani, B. L. Ekpukpon, O. A. Odeku, B. I. Olu-Owolabi, and K. O. Adebowale, 'Intra and Extra-granular Disintegrant Properties of Modified Underutilised Red Lima Bean Starch in Paracetamol Tablet Formulation', *Brazilian J. Pharm. Sci.*, vol. 58, 2022, doi: 10.1590/s2175-97902022e19967.
- [11] L. Lachman and Kanig, *Teori dan Praktek Farmasi Industri*, 2nd ed. Jakarta Indonesia: UI Press, 1994.
- [12] M. 20852 The United States Pharmacopeial Convention 12601 Twinbrook Parkway, Rockville, Ed., *The United States Pharmacopeia, USP 41/The National Formulary, NF 36*. Baltimore, MD. Rockville.: United Book Press, Inc., 2018.
- [13] G. K. Bolhuis and Z. T. Chowhan, Materials for Direct Compaction, in Pharmaceutical Powder Compaction Technology, 420th ed. New York and Basel: Marcel Dekker, 1996.
- [14] Food and Drug Supervisory Agency of the Republic of Indonesia, No Title 1200, 2014, BPOM. Obat Tradisional. Mutu. Persyaratan., no. 1200. 2014.
- [15] R. Panigrahi and S. Behera, 'A Review On Fast Dissolving Tablets', vol. 1, no. 11, pp. 1–15, 2010, doi: 10.5958/0974-360X.
- [16] C. P. Jain and P. S. Naruka, 'Formulation and Evaluation of Fast Dissolving Tablets of Doxazosin Mesylate', Asian J. Pharm. Res., vol. 6, no. 3, p. 131, 2016, doi: 10.5958/2231-5691.2016.00020.4.
- [17] Allen, L.V., N. G. Popovisch, and H. C. Ansel, Ansel's Dossage Forms and Drug Delivery Systems, 9th Ed., no. 1. 2011.
- [18] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia (FHI), II. Jakarta: k, 2017.
- [19] L. Lachman and D. Ph, The Theory and Practice of Industrial Pharmacy: Tablets, Gilbert S. Banker, Neil R. Anderson, 3rd ed. Garden City, New York; Livingston, New Jersey; Ridgefield, Connecticut, 1986.
- [20] H. Maurya and T. Kumar, 'Formulation, standardization, and evaluation of polyherbal dispersible tablet', Int. J. Appl. Pharm., vol. 11, no. 1, pp. 158–167, 2019, doi: 10.22159/ijap.2019v11i1.30113.
- [21] P. Sirame, R. Khairnar, and R. Tasgaonkar, 'Granulation', Int. J. Res. Appl. Sci. Eng. Technol., vol. 11, no. 1, pp. 347–355, Jan. 2023, doi: 10.22214/ijraset.2023.48564.
- [22] C. M. Shinko, 'Granulation Characterization: Methods and Significance, Dalam: Parikh, D.M.(ed.), Handbook of Pharmaceutical Granulation Technology, Marcel Dekker, New York', p. 422, 1997.
- [23] J. Swarbrick, J. C. Boylan., and 2002, *Encyclopaedia of Pharmaceutical Technology*, 2nd ed., vol. 3, no. 4. New York: Organic Process Research * Development 2003, 613, 2003.
- [24] Dinesh Puri et al., 'Formulation of Herbal Fast Disintegrating Tablets and its', Curr. Clin. Pharmacol. Bentham Sci., vol. 13, no. 2, pp. 128–135, 2018, doi: 10.2174/1574884713666180309153942.
- [25] D. Puri et al., 'Formulation of Herbal Fast Disintegrating Tablets and its ex-vivo Study for Antihistaminic Activity in Guinea Pig Ileum', Curr. Clin. Pharmacol., vol. 13, no. 2, pp. 128–135, 2018, doi: 10.2174/1574884713666180309153942.
- [26] J. Balasubrmaniam, Balasubrmaniam, J. 2009, Influence of Superdisintegrants on The Rate of Drug Dissolution from Oral Solid Dosage Forms, Pharmaceutical Technology, April., April. Pharmaceutical Technology, 2009.
- [27] A. N. Bestari, T. N. S. Sulaiman, and A. Rohman, 'Formulasi Orally Disinetgration Tablet (ODT)', vol. 12, no. 2, pp. 453–465, 2016, doi: 10.22146/farmaseutik.v12i2.26454.
- [28] C. G. Jange, C. R. Wassgren, and K. Ambrose, 'The Significance of Tablet Internal Structure on Disintegration and Dissolution of Immediate-Release Formulas: A Review', *Powders*, vol. 2, no. 1, pp. 99–123, Feb. 2023, doi: 10.3390/powders2010008.
- [29] F. CDER, 'Guidance for Industry Orally Disintegrating Tablets Guidance for Industry Orally Disintegrating Tablets', no. December, 2008.
- [30] 2010 The Council of Europe: The European Pharmacopoeia, 7th Edition, Strasbourg, France, 'Dalam: Brniak, W Jachowicz, R, Pelka, P, The practical approach to the evaluation of methods used to determine the disintegration time of orally disintegrating table', *Saudi Pharm. J.*, vol. 23, no. 4, pp. 437–443, 2015, doi: 10.1016/j.jsps.2015.01.015.
- [31] S. Kakar, R. Singh, and S. Kumar, 'Orodispersible tablets: an overview', MOJ Proteomics

Int J Public Health Sci, Vol. 99, No. 1, Month 2099: 1-1x

Int J F	Public Health Sci	ISSN: 2252-8806	0
	Apoinforma., vol. 7, no. 3,	op. 180–182, 2018, doi: 10.15406/mojpb.2018.07	7.00230.
[32]	P. Dey and S. Maiti, 'Orod	ispersible tablets: A new trend in drug delivery',	J. Nat. Sci. Biol. Med.,
	vol. 1, no. 1, p. 2, 2010, do	i: 10.4103/0976-9668.71663.	
[33]	A. Ainurofiq and N. Aziza	n, 'Perbandingan Penggunaan Bahan Penghancur	r Secara Intragranular,
	Ekstragranular, Dan Komb	inasinya', J. Pharm. Sci. Clin. Res., vol. 01, no. 0	01, pp. 1–9, 2016, doi:
	10.20961/jpscr.v1i1.682.		
[34]	S. U. Pradhan et al., 'Chara	cteristics of multi-component formulation granu	les formed using
	distributive mixing elemen 1826–1837, Nov. 2018, do	ts in twin screw granulation', <i>Drug Dev. Ind. Pha</i> : 10.1080/03639045.2018.1503293.	arm., vol. 44, no. 11, pp
35			

BIOGRAPHIES OF AUTHORS

The recommended number of authors is at least 2. One of them as a corresponding author. *Please attach clear photo (3x4 cm) and vita. Example of biographies of authors:*



Paper's should be the fewest possible that accurately describe ... (First Author)

	0
N: 2252-8806	75. 275.
11. 2252 0000	
	TN Saifullah is lecturer in department of technology pharmaceutical Gadjah Mada university, Yogyakarta Indonesia.

Int J Public Health Sci, Vol. 99, No. 1, Month 2099: 1-1x

Formulation of Intra and Extra-Granularly Croscarmellose in Fast Disintegrating Tablet of Bay Leaf Extract (Syzygium polyanthum)

ORIGIN	ALITY REPORT				
SIMIL	4% ARITY INDEX	10% INTERNET SOURCES	8% PUBLICATIONS	6% STUDENT PA	PERS
PRIMA	RY SOURCES				
1	Submitt Kement Student Pape	ed to Badan PP erian Kesehatar	SDM Kesehata า	an	3%
2	eprints. Internet Sour	uad.ac.id			1%
3	Submitt Student Pape	ed to Universita	as Kristen Duta	a Wacana	1%
4	Submitt Student Pape	ed to Ain Sham	s University		1%
5	www.ph Internet Sour	iytojournal.com			1%
6	Siti Fario Sahlan, Rohmat Antioxic Acute To Capsule Journal Publication	da, Diah Kartika Dian Ratih Laks in, Herbert Situ lant, In-Vivo Ant oxicity Study of from Tetragon of Biological Sci	Pratami, Muh mitawati, Etin morang. "In-Vi ti-Inflammator Indonesian Pr ula sapiens", S ences, 2021	amad tro y, and opolis audi	1 %

7	Muhammad Fariez Kurniawan, Fadhilah Alvari Yusuf. "The Possible Antidiabetic Effect of Ficus carica L. Tablet on Iloxan-Induced Diabetes Model in Rats", Open Access Macedonian Journal of Medical Sciences, 2021 Publication	1%
8	doaj.org Internet Source	<1%
9	repository.unib.ac.id	<1%
10	f1000research.com Internet Source	<1%
11	repository-tnmgrmu.ac.in	<1%
12	semnaskes.unipasby.ac.id	<1%
13	ijcc.chemoprev.org Internet Source	<1%
14	mdpi-res.com Internet Source	<1 %
15	www.aaspyoungscicof.com	<1%
16	Submitted to Rochester Institute of Technology Student Paper	<1%

17	nutritionandmetabolism.biomedcentral.com	<1 %
18	the-medical-dictionary.com	<1 %
19	Submitted to University of Bradford Student Paper	<1 %
20	A Akrom, Mustofa, L H Nurani. "Ethanolic Extract of Black Cumin Seed Reduced Radical Reactive from Dimethylbenzanthracene Compounds", IOP Conference Series: Earth and Environmental Science, 2021 Publication	<1%
21	Eliya Mursyida, Raissa Almira, Santi Widiasari, Olvaria Misfa. "Antibacterial Activity of Bay Leaf (Syzygium polyanthum) Ethanol Extract on Escherichia coli Growth", Photon: Jurnal Sain dan Kesehatan, 2021 Publication	<1%
22	Iria Seoane-Viaño, Xiaoyan Xu, Jun Jie Ong, Ahmed Teyeb et al. "A case study on decentralized manufacturing of 3D printed medicines", International Journal of Pharmaceutics: X, 2023 Publication	<1 %

23 M A Dhina, D A Deswati, S R Mubaroq. "Increasing dissolution rate of telmisartan tablet by dispersion method", IOP Conference

<1 %
Series: Materials Science and Engineering, 2018 Publication

Syarief Thaufik Hidayat, Inu Mulyantoro, Suhartono Damas, Raymond R Tjandrawinata. "The Effect and Safety Assessment of Metformin and DLBS3233 (A Bioactive Fraction of Lagerstroemia speciosa and Cinnamomum burmannii) on Improving Metabolic Parameters in Women with Polycystic Ovary Syndrome", International Journal of Women's Health, 2023 Publication

Titiek Hidayati, Akrom Akrom, Indrayanti Indrayanti, Sagiran Sagiran. "Black cumin seed oil increase leucocyte and CD4Thelper number in sprague-dawley rats induced with dimethylbenzanthracene", International Journal of Public Health Science (IJPHS), 2019 Publication

26

innovareacademics.in Internet Source

- Jaime Conceição, Oluwatomide Adeoye, Helena Cabral-Marques, Angel Concheiro, Carmen Alvarez-Lorenzo, José Manuel Sousa Lobo. "Carbamazepine bilayer tablets combining hydrophilic and hydrophobic cyclodextrins as a quick/slow biphasic release
- <1%

<1 %

<1%

system", Journal of Drug Delivery Science and Technology, 2020 Publication

28	academic.oup.com	<1%
29	repository.stikim.ac.id	<1%
30	storage.googleapis.com	<1%
31	www.researchgate.net	<1%
32	WWW.rroij.com Internet Source	<1%
33	www.sciencegate.app	<1%
34	J. ZH. Gao. "Investigation of human pharmacoscintigraphic behavior of two tablets and a capsule formulation of a high dose, poorly water soluble/highly permeable drug (efavirenz)", Journal of Pharmaceutical Sciences, 11/2007 Publication	<1 %
35	Submitted to Universitas Brawijaya	<1%

Exclude quotesOnExclude matchesOffExclude bibliographyOn

UNIVERSITAS AHMAD DAHLAN

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] REVISION ID 22666

3 pesan

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: Widyasari@pharm.uad.ac.id

Dear Author,

We inform you that your article needs improvement, with this email we attach the file that needs to be revised and we have provided comments on the file. Please revise the attached file and put a yellow mark on the part you are revising and ask us to edit the Bibtex file. We give until Oct 12, 2023. Thank you very much for your attention.

Kind Regards, Maria Ulfah Editorial Staff on behalf of Editor-in-Chief Layout and Editing Team International Journal of Public Health Science Institute of Advanced Engineering and Science (IAES) ISSN: 2252-8806 https://ijphs.iaescore.com/index.php/IJPHS/index email: ijphsiaes.core@gmail.com, ijphs@iaescore.com

22666.docx 4935K

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

<u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

[Kutipan teks disembunyikan]

त्रिको	22666.docx
E	4365K

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

Thanks for revision. Please revise the following files again.

[Kutipan teks disembunyikan]

[Kutipan teks disembunyikan]

UNIVERSITAS AHMAD DAHLAN

Kampus 1: Jln. Kapas No. 9 Yogyakarta Kampus 2: Jl. Pramuka 42, Sidikan, Umbulharjo, Yogyakarta 55161 Kampus 3: Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta 55164 14 Oktober 2023 pukul 16.58

10 Oktober 2023 pukul 16.28

17 Oktober 2023 pukul 14.47

5/31/24, 10:44 AM

Kampus 4: Jl.Ringroad Selatan, Yogyakarta

Kampus 5: Jl. Ki Ageng Pemanahan 19, Yogyakarta

Kontak

Email: info@uad.ac.id

Telp. : (0274) 563515, 511830, 379418, 371120

Fax. : (0274) 564604





Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract

Widyasari Putranti¹, Desty Restia Rahmawati¹, Nining Sugihartini¹, Teuku Nanda Saifullah²

¹Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia ²Department of Pharmaceutical Technology, Faculty of Pharmacy, Gadjah Mada University, Yogyakarta, Indonesia

Article Info ABSTRACT

Article history:

Received month dd, yyyy Revised month dd, yyyy Accepted month dd, yyyy

Keywords:

Croscarmellose Superdisintegrant Syzygium polyanthum extract Fast disintegrating tablet Intra granular Extra granular Bay leaves (Syzygium polyanthum) contain the flavonoid quercetin which can be used as an antihyperlipidemic drug. The development of antihyperlipidemic drug formula in the form of fast disintegrating tablet (FDT) is needed for patients who experience dysphagia. FDT preparations require an optimal super disintegrant concentration to produce a good drug formula. This study aims to develop the FDT formula of bay leaves extract using the super disintegrant croscarmellose sodium (CCS) intra and extragranular. FDT formulation using the wet granulation method with variations of CCS concentrations; F1: 2%, F2: 3.5%, and F3: 5% for extra granular, and 2% for intra granular. The formulation process, in-process control (IPC) granules, weight uniformity tests, and various physical properties tests of tablets were carried out. Data were statistically analyzed using one way ANOVA test (α =95%). The results of statistical tests of IPC granules, uniformity of weight, and tablet size of all FDT formulas were not significantly different (p>0.05). The CCS concentration for extra granular significantly affected the wetting time, disintegration time, hardness, and the value of friability percentage of FDT (p<0.05). The combination of intra and extra-granular CCS (2%:5%) gave the most optimum physical properties of bay leaf extract FDT.

This is an open access article under the <u>CC BY-SA</u> license.



Corresponding Author:

Widyasari Putranti Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan Yogyakarta, Indonesia Email: widyasari@pharm.uad.ac.id

1. INTRODUCTION

Plant germplasm resources are raw materials for medicines, one of which is a bay leaf (*Syzygium polyanthum*). Bay leaves are known to contain secondary metabolites, such as saponins, terpenoids, flavonoids (quercetin), polyphenols, alkaloids, steroids, and essential oils (sesquiterpenes)[1]-[2]. Based on the Decree of the Minister of Health of the Republic of Indonesia (2009), bay leaves contain a total flavonoid of not less than 0.40% which is calculated as quercetin [3]. Quercetin in bay leaves the potential as an antihyperlipidemic which can significantly reduce levels of triglycerides, total cholesterol, and low-density lipoprotein (LDL) cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in high-density lipoprotein (HDL), as well as inhibiting 3-hidroksi-3-metilglutaril coenzyme A (HMG-CoA), reductase, and LDL oxidation [4].

Conventional drugs for hyperlipidemia are generally available in tablet form. However, the tablet dosage form has several disadvantages. Some of them require a long absorption time in the body, and slow drug action, and elderly patients may experience difficulty swallowing or dysphagia [5]. Seeing the problems that arise from conventional tablets, the herbal fast disintegrating tablet (FDT) of bay leaf extract is the right alternative so that the drug can be used comfortably and has a therapeutic effect more quickly. FDT is a solid

1

dosage form containing pharmaceutically active substances that can be disintegrated quickly, generally within seconds when placed on the tongue. This properties of FDT can release the active drug substance immediately, speed up the onset, increase the oral bioavailability of the drug in the body so that therapeutic effectiveness can be achieved [6]-[7]. FDT preparations combine two advantages of preparations, namely liquid which can increase the solubility and bioavailability of drugs, as well as the advantages of solid dosage forms that have high physicochemical stability, relatively constant homogeneity of the active ingredients of the preparation, and easier manufacturing.

FDT is formulated with croscarmellose sodium (CCS) super disintegrant which is classified as a strong swelling product with a good water absorption mechanism [7]-[8]. CCS is effectively used in FDT preparations because it rapidly expands to 4-8 times its initial volume when in contact with water, thus accelerating the tablet-crushing process. CCS is effective intra-granular which allows for better capillary and swelling processes, causing the tablet to disintegrate quickly and reduce disintegration time [8]. The addition of intra and extra-granular crushing materials can increase the effectiveness of tablet disintegration through two stages, i) extra granular which breaks the tablet into granules, and ii) intra granular breaks down the granules into smaller particles. The mechanism is crushing material that experiences rapid swelling and water absorption (wicking) through the gaps between the granules and the particles that make up the granules [9]. This study aims to develop herbal FDT with the active ingredient of bay leaf ethanol extract. The concentration of CCS super disintegrant intra and extra-granular needs to be determined to obtain the FDT of bay leaf extract with optimum physical properties.

2. METHOD

The ingredients used were bay leaf extract in the form of fresh bay leaf collected from tradisional market, Beringharjo market, Yogyakarta, Indonesia, ethanol 70%, croscarmellose sodium (PT. Phapros Tbk), lactose, magnesium stearate, aspartame, menthol and corn starch paste (pharmaceutical grade). The tools for extraction used are oven, rotary evaporator calipers, waterbath, vacuum pump. The tools for granule physical properties test used are Ohaus analytical balance, absorption test equipment granules, volumenometer (tapped density tester), flow tester granule, sieving machine, dry granule sieve, wet granule sieve, pan, halogen moisture analyzer. the tools for tablet test used are single punch tablet printing machine, hardness tester, friability tester, and disintegration tester.

2.1. Bay leaf ethanol extract

Bay leaves were dried in an oven at 50-60°C until a moisture content of less than 10% was obtained. Simplicia were extracted by maceration method using 70% ethanol solvent (1:10) (w/v) [3]. Maserati and pulp were separated by the vacuum pump and filter paper. The solvent was evaporated using a rotary evaporator. The viscous extract obtained was calculated for its yield.

2.2. FDT formulation of bay leaf extract

FDT was made by the wet granulation method. At the granulation stage, bay leaves extract was mixed with intra granular CCS, lactose and corn starch. The mixture was sieved through a wet (no. 18 mesh), ovendried (50-60°C), then sifted dry (no. 20 mesh). The granules and excipients consisting of magnesium stearate, aspartame, menthol, and CCS (extra granular) mixed until homogeneous (30 minutes). Mixture of granules and excipients that have been homogenized are compressed using a single punch tablet press.

2.3. Granule physical properties test parameters

The moisture content of granules can be determined based on the value of loss on drying (LOD) and MC. LOD is a test for measuring the difference in heavy total granules when before and after drying or a moisture content statement based on wet weight. Meanwhile, MC is a statement of moisture content based on dry weight. Measurements using the halogen moisture analyzer tool. Good granule moisture content if the LOD and MC values are 10% [10].

The flow properties of the granules can be tested by calculating the angle of repose, the flow time test, and the determination index (Hausner ratio). The granule angle of repose α (degree) is said to be very good (25°-30°); good (30-35°) or 25°><35°. A good granule flow time is not more than 10 seconds for 100 grams of granules [11]. Excellent granule setting index (\leq 10); good (11–15), and the Hausner ratio index that meets the requirements for the physical properties of the granule mass, which is very good (1.00–1.11); good (1.12–1.18).

Sieve no. 14, 16, 20, 30, 50 mesh and pan, installed in stages on the sieving machine (50 amplitude, 15 minutes). The granules left on each sieve were weighed and the percentage was calculated [10]. The fragility of the granules is based on the number of fines that occur after the friability test by sieving. The granules were

D 3

put into a graded sieve with the top sieve no. 30 mesh and bottom pan, using a sieving machine (50 amplitude, 30 minutes). The water absorption of the granules affects the tablet's disintegration time (disintegration). It was carried out with a series of absorption test equipment by placing 300 mg of granules on a water-saturated sheath paper. The absorbed water was measured using a digital scale for 15 minutes or until it was constant [7].

2.4. Physical properties test parameters of FDT

Requirements and weight uniformity tests are determined based on the regulation of the National Food and Drug Agency of the Republic of Indonesia concerning the quality requirements of traditional medicines. The test was carried out by weighing 20 tablets one by one, from 20 tablets no more than 2 tablets, each of which deviated from the average weight by more than 5% and not one tablet whose weight deviated from the average weight. the average is greater than 10% [12].

Each tablet was measured in diameter in a horizontal position with a caliper, and the thickness of the tablet was measured in a vertical position with a caliper (millimeters). The tablet hardness test was carried out by taking a sample of at least 6 tablets from each formula. The tablet hardness test is carried out using hardness tester tablets. One by one the tablets are tested by placing the tablets in the center perpendicular to the hardness tester. A good FDT hardness is 3-5 kg/cm² [11], [13].

For tablets with a single weight of exactly or less than 650 mg, the total weight of the tested samples was close to 6.5 grams. Prior to testing, the tablets were dusted and weighed. All tablets were put into the friability tester (speed 25 rpm, 4 minutes). The tablets were cleaned of adhering fines and re-weighed. A good friability value should not be more than 1% [11].

The wetting time test was done by placing a sheet of filter paper that has been folded twice into a petri dish with a diameter of 5 cm. The petri dish was filled with 5.0 mL of distilled water containing the strawberry red dye. A tablet is then placed on the filter paper, and simultaneously with the start of the test instrument, the stopwatch is turned on. Wetting time was calculated as the time required for a red color to appear on the entire surface of the tablet [14]. A total of 6 tablets of each formula were placed in each tube and a disk was placed on it, the test was carried out with water medium at a temperature of 37 ± 2 °C or 35-39 °C using a 1000 ml glass beaker. The disintegration time requirement for FDT preparations is at least <1 minute disintegration time [11], [15].

2.5. Data analysis

The results of the physical examination of granules and tablets of FDT bay leaf extract were analyzed using the one-way ANOVA test method with a 95% confidence level.

3. RESULTS AND DISCUSSION

Bay leaf simplicia was tested for macroscopic (organoleptic) specifications on fresh and dried simplicia, as well as microscopic tests for simplicia parts and identification fragments of simplicia powder. Fragments prepared by dissolving in chloralhydrate and water [16]. The simplicia specification test was carried out to ensure the correctness of the identity and quality of the bay leaf simplicia based on the Indonesian Herbal Pharmacopoeia compendia and Indonesian Materia Medika. Based on the microscopic test results on Figure 1 there are identifier fragments namely i) lower epidermis with stomata, ii) upper epidermis, iii) elements with dots, iv) prism-shaped calcium oxalate crystals, and v) sclerenchyma. The results of the specification test for the simplicia of bay leaves showed that the simplicia used met the requirements of organoleptic and microscopic terms. The determination test of bay leaf simplicia at the Biology Laboratory of Universitas Ahmad Dahlan (UAD) was carried out to ensure that the selected bay leaf simplicia was in accordance with its species or identity [3], [17].

3.1. Extraction bay leaf

Fresh simplicia of bay leaves were sorted wet, dried using an oven $(50-60^{\circ}C)$ with a moisture content of not more than 10%, then sorted dry. Simplicia powder for extract preparation is a fine simplicial (No. 60 mesh) in order to optimize the extraction process [18]. The results of the simplicial extraction of bay leaves weighing 3.7 kg obtained a thick extract of 347 grams. The total yield of the thick extract obtained was 9.38%. The morphology of the viscous extract is shown in Figure 2. Screening and determination of the chemical compound flavonoid quercetin contained in the thick extract of bay leaves was carried out at the Integrated Research Laboratory of the Faculty of Pharmacy, UAD. According to the Ministry of Health (2017), bay leaves contain a total flavonoid of not less than 0.40% calculated as quercetin. Based on the results of the test the flavonoid content of 0.8079±0.0045%, so that it meets the specifications for the total flavonoid content of bay leaves[17]. Quercetin is an active substance that has medicinal effects such as antiherlipidemia, antibacterial, antidiabetic, antioxidant [18], [19].



Figure 1. Identification fragments of bay leaf; (a) lower epidermis, (b) upper epidermis, (c) floem, (d) calcium oxalate crystal, (e) sclerenchyma



Figure 2. Bay leaf Extract

3.2. FDT formulation

FDT formulation method with wet granulation is good used for water-resistant active ingredients and heating temperature [10]. The active ingredient in bay leaf is quercetin (3,3',4',5,7-pentahydroxyflavone) is a flavonoid derivative compound that is resistant to heating with a melting point of 316.5 °C [20], while the wet granulation process ranges from at a temperature of 40-60 °C. The wet granulation method can improve the flowability and compatibility of the material and facilitate the FDT formulation process from the active ingredient in the form of a thick extract [21]. Wet granulation has the advantage of good content distribution and uniformity, besides that the hydrophobic surface of the tablet becomes more hydrophilic which affects the water absorption process into the tablet, thereby increasing the speed of tablet disintegration [22]. The work of woven fibers between granules can increase the speed of tablet disintegration[23], it is synergistic with the presence of CCS super disintegrant extra granular which facilitates swelling and absorption of water into granules and their small particles.

The concentration of CCS super disintegrant intra-granular of the three formulas was the same, 2% (10 mg/tablet), and extra granular was formulated with varying levels, F1 2% (10), F2 3.5% (17.5), and F3 5% (25 mg/tablet). The FDT formula for bay leaf extract is shown in Table 1. Based on the Handbook of pharmaceutical excipients the use of CCS in the wet granulation process should be added in the wet or granulation (intra-granular) stage, as well as in the dry (extra-granular) process [21]. Because this method can increase the wicking ability (water absorption) and swelling of the CCS super disintegrant.[8]

Table 1. I official of I DI bay leaf extract				
Ingredient		position	(mg)	
	F1	F2	F3	
Bay leaf extract	70	70	70	
Croscarmellose sodium (intra granular)	10	10	10	
Lactose	380	374	366	
Croscarmellose sodium (extra granular)	10	17.5	25	
Magnesium stearate	5	5	5	
Aspartame	20	20	20	
Menthol	1	1	1	
Corn starch paste 5% w/v	qs	qs	qs	
Total weight (mg)	500	500	500	

Table 1. Formula of FDT bay leaf extract

3.2.1. Granule physical properties test parameters

One of the most crucial elements of wet granulation is the water content, MC and LOD has been was measured and the values of granules from all formulas as shown in Table 2, the results obtained meet the requirements of 10% (p>0.05). Water content will affect granule characteristics, including granule size and compressibility. The flow properties of the granules show the results of the angle of repose of the granules ranging from 30-35 (good), granule flow time of 10-14 grams/second (good), and the determination test/Hausner ratio (very good). The flow properties of granules are very influential on the tablet compression process, granules with good flow properties will cause the granules to enter the die space to be relatively constant so that the tablet weights with small weight variations can be obtained and can increase the uniformity of the resulting tablet dosage [8], [24], [25].

The average diameter of all granules ranged from 650-700 m (p>0.05) and the value of friability percentage of the granules ranged from 0.5-0.9% (p>0.05)[13]. The granule absorption test for all formulas had absorption between 0.120-0.190 mg/minute (p>0.05). The granule absorption shows the speed of the granules in absorbing water to crush the granules into small particles so as to speed up the tablet disintegration time. All in process control (IPC) granule parameters have met the requirements so that the tableting process can be continued with a single punch tablet press with a punch pressure of 13 kg.

Table 2. Results of examination of physical properties of FDT granules of bay leaf extract

Test parameters	Formula				
	F1	F2	F3		
LOD±SD (%)*	7.03±0.76	6.29±1.28	6.57±1.18		
MC±SD (%)*	2.98 ± 0.17	2.35±0.33	2.69±0.31		
Angle of repose±SD (°)*	30.66±1.75	31.18 ± 2.92	30.08±2.30		
Flow time±SD (grams/second)*	12.98 ± 1.28	13.12±0.89	11.85±0.48		
Assignment±SD (%)*	3.17±1.04	2.83±1.04	3.83±0.76		
Hausner ratio Index±SD*	1.03 ± 0.01	1.03 ± 0.01	1.04 ± 0.01		
Average diameter±SD (µm)*	694.64 ± 28.87	657.58±39.65	698.14±26.09		
Fragility±SD (%)*	0.8028 ± 0.20	0.8199 ± 0.05	0.5863 ± 0.04		
Absorption±SD (mg/min)*	0.186 ± 0.05	0.188 ± 0.05	0.127±0.01		

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

3.2.2. Physical properties test parameters of FDT

The uniformity of tablet weight is related to the uniformity of active substance levels and therapeutic effects. The uniformity of tablet weight (Table 3) can be seen from the coefficient of variation F1, F2, and F3 all <5%, (0.0942, 0.1246, and 0.2103) %, respectively. The weight uniformity test based on the National Food and Drug Agency of the Republic of Indonesia requirements, shows the results that meet the requirements, where the tablet meets the specified weight deviation limit value range.

Hardness ranges FDT be good tablet from 3-5 kg/cm² [13], while the conventional tablet, is 4-8 kg/cm². The FDT hardness as shown in Table 3 shows results that meet the requirements. Hardness affects the brittleness and disintegration time of tablets, as well as the dissolution/release of the active drug substance [8]. The results of statistical analysis of tablet hardness with sig. (p<0.05) indicates that the hardness of the three FDT formulas is significantly different. The hardness of FDT tablets is strongly influenced by the CCS super disintegrant excipient. FDT (F3) produced the hardest tablet with the highest extra-granular CCS concentration (5%), causing the particles in the tablet to be tightly bound. Croscarmellose sodium (Ac-Di-Sol) is a super disintegrant with a fibrous cross-linked polymer structure that functions as a disintegrant with a strong swelling mechanism, fiber is able to bind strongly to the particles so as to increase the hardness and reduce the brittleness of the tablet [26]. This causes the tablet to have good compatibility, so there is a tendency

to increase hardness as the amount of CCS in the tablet increases [27]. Another factor that affects the hardness of FDT is the upper punch pressure during the tableting process, the greater the pressure used, the harder the FDT will be. The results of the diameter and thickness measurements of the tablets showed that the size of the entire tablet was uniform (homogeneous).

The friability value of the tablet is based on the requirements, which should not be more than 1% [11]. The fragility value of FDT is shown in Table 3 with a friability value of <1% so that FDT meets the requirements. The results of the tablet friability statistical test have a sig. value (p < 0.05) means that the fragility value between formulas is significantly different. CCS super disintegrant is a factor influencing fragility. Formula 3 with the highest number of CCS has the smallest friability value, where there is a tendency to decrease the friability value as the number of CCS in the tablet increases. The friability value is inversely proportional to the tablet's hardness, where the harder a tablet is, the smaller the friability value will be.

FDT wetting time is shown in Table 3, FDT (F3) with the highest number of CCS has the smallest wetting time. Wetting time is a parameter to determine the speed of FDT in absorbing water, which affects the speed of tablet disintegration. The faster the wetting time, the faster the tablet disintegration. The wetting time of tablets is influenced by the structure of the tablet matrix and the hydrophilicity of the excipients [28]. The result of statistical analysis of FDT wetting time has a sig. value (p<0.05) means that the formulas differ significantly. FDT (F3) with the highest number of CCS had the fastest time for wettable tablets, super disintegrant CCS being the most influential factor. The cross-linked chemical structure of CCS creates excipients that have hydrophilic properties and very easy-to-absorb solvents resulting in outstanding swelling properties in the disintegration process.

Table 3. Results of examination of physical properties of bay leaf extract FDT tablets

Test Demonstern		Note.		
Test Parameters	F1	F2	F3	
Weight uniformity±SD (mg)	50 4.42±0.48	50 4.78±0.63	50 4.43±1.06	Appropriate*
Diameter±SD (mm)	12.00±0.00	12.00±0.00	12.00±0.00	Appropriate*
Thickness±SD (mm)	3.67±0.00	3.67±0.00	3.68±0.01	Appropriate*
Hardness±SD (kg)	4.46±0.25°	4.50±0.27°	5.24±0.32 ^{a,b}	In accordance
Fragility±SD (%)	0.77±0.05°	0.72±0.04°	$0.52 \pm 0.04^{a,b}$	In accordance
Wetting time±SD (seconds)	50.33±3.06°	43.33±4.04°	20.00±2.65 ^{a,b}	In accordance
Disintegration time±SD (seconds)	45.33±5.86°	38.33±3.79°	19.67±3.06 ^{a,b}	In accordance

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

(a) means a formula that is significantly different from the formula 1

(b) means a formula that is significantly different from the formula 2

(c) means a formula that is significantly different from formula 3

F1: CCS intra granular 2%: extra granular 2%

F2: CCS intra granular 2%: extra granular 3.5%

F3: CCS intra granular 2%: extra granular 5%

Disintegration (disintegration) time FDT has at least a disintegration time of less than 1 minute [15]. Another source says that in-vitro about 30 seconds or less [29], whereas according to Ph. euros. is <3 min in the oral cavity before swallowing [29], [30]. The faster disintegration time will increase the speed of drug release from the tablet which affects the effectiveness of therapy. The results of the disintegration time of FDT in table 3 show that it meets the requirements of the disintegration time of FDT (<1 minute). FDT disintegration time test is shown in Table 3 with FDT (F3) having the fastest disintegration time, due to the higher levels of CCS used in F3 than F1 and F2. The results of the statistical test of tablet disintegration time have a sig value (p < 0.05) means that each formula is significantly different. CCS has a dual mechanism, namely water wicking and rapid swelling [8]. The highly porous form of the structure speeds up the disintegration time, because water quickly enters the tablet and increases the tablet wetting rate, through the gaps between the granules and the granule pores [31]. This is related to the factor of the number of CCS super disintegrant excipient used in each formula intra and extra-granular. The use of intra and extra-granular affects the decrease in tablet disintegration time through the swelling mechanism, and along with the increase in the extra-granular CCS concentration, the disintegration time will be faster. The FDT disintegration pattern is also facilitated by intra-granular CCS which causes the tablets to be finely crushed not in the form of granules but into small particles so that the speed of disintegration is optimal. The optimum formula for FDT of bay leaf extract is shown in Table 4 with the results of all evaluations of the physical properties of the granules and FDT (F3) that meet the requirements and show the optimum results of physical parameters.

The results of this study are in line with the research of Puri *et al.* [24], that the combination of intragranular and extra-granular can produce tablets with the most effective disintegration time by providing better tablet physical properties, compared to intra-granular or extra-granular only. Extra-crushing material granules

D 7

have a high tendency to absorb water from the surrounding liquid through the tablet-breaking mechanism [32]. Tablets with disintegrant intra-granular have a higher hardness than tablets with extra granular crushing agents. By combining intra and extra granular super disintegrants, a good level of tablet hardness can be obtained so that the tablet is not brittle and has a faster disintegration time with the mechanism of tablet disintegration into granules (extra granular factor) and granules into small particles (intra granular factor).

1 u 0 0 - 1, $u 1 0 u 0 p u 1 u 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0$	Ta	ble	4.	Va	arious	parameters	of	or	otimized	tab	olet	formula	a (F3)
--	----	-----	----	----	--------	------------	----	----	----------	-----	------	---------	-----	----	---

No.	Test parameters	Optimized formula value
1.	Weight uniformity±SD (mg)	504.43±1.06
2.	Diameter±SD (mm)	12.00±0.00
3.	Thickness±SD (mm)	3.68±0.01
4.	Violence±SD (kg)	5.24±0.32
5.	Fragility±SD (%)	0.52±0.04
6.	Wetting time±SD (seconds)	20.00 ±2.65
7.	Disintegration time±SD	19.67±3.06
	(seconds)	

4. CONCLUSION

In the FDT research, to improve the poor flow properties of the extract, the wet granulation method was carried out. The IPC results for the three formulas showed granules that met the requirements. This research provides information that the use of extra granular CCS at higher levels will increase the rate of disintegration. The combination of CCS intra granular 10 mg (2%) and extra granular 25 mg (5%) can produce FDT with the most effective. By combining intra and extra granular super disintegration time. In addition, FDT bay leaves extract can be used as an alternative treatment for hyperlipidemia in patients who experience dysphagia. This research still needs development to determine the stability of FDT bay leaves extract where the active substance is herbal medicine.

ACKNOWLEDGEMENTS

This research was carried out with the assistance of the Higher Education Research Grant No PKPT-013/SKPP/III/2018.

REFERENCES

- L. Hartanti, S. M. K. Yonas, J. J. Mustamu, S. Wijaya, H. K. Setiawan, and L. Soegianto, "Influence of extraction methods of bay leaves (Syzygium polyanthum) on antioxidant and HMG-CoA Reductase inhibitory activity," *Heliyon*, vol. 5, no. 4, p. e01485, Apr. 2019, doi: 10.1016/j.heliyon.2019.e01485.
- [2] E. Kato, R. Nakagomi, M. D. P. T. Gunawan-puteri, and J. Kawabata, "Identification of hydroxychavicol and its dimers, the lipase inhibitors contained in the Indonesian spice, Eugenia polyantha," *Food Chem.*, vol. 136, no. 3–4, pp. 1239–1242, 2013, doi: 10.1016/j.foodchem.2012.09.013.
- [3] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia, 1st ed. Jakarta Indonesia, 2009.
- [4] K. Zeka, K. Ruparelia, R. Arroo, R. Budriesi, and M. Micucci, "Flavonoids and Their Metabolites: Prevention in
- Cardiovascular Diseases and Diabetes," Diseases, vol. 5, no. 3, p. 19, 2017, doi: 10.3390/diseases5030019.
- [5] I. Safitri, S. Sulistiyaningsih, and A. Y. Chaerunisaa, "Review: Superdisintegran dalam Sediaan Oral," *Farmasetika.com* (Online), vol. 4, no. 3, p. 56, 2019, doi: 10.24198/farmasetika.v4i3.22945.
- [6] A. Masih, A. Kumar, S. Singh, and A. K. Tiwari, "Fast Dissolving Tablets: A Review," *Int. J. Curr. Pharm. Res.*, vol. 9, no. 2, p. 8, Mar. 2017, doi: 10.22159/ijcpr.2017v9i2.17382.
 [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam
- [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam leaves ethanol extract with combination of crospovidone and croscarmellose sodium as superdisintegrants," *J. Sains Farm. Klin.*, vol. 8, no. 3, p. 285, 2021, doi: 10.25077/jsfk.8.3.285-295.2021.
- [8] M. E. Rowe, R. C., Sheskey, P. J. dan Quinn, *Handbook of Pharmaceutical Excipients*, Sixth. London: Pharmaceutical press., 2009.
- [9] M. O. Okekunle, O. D. Akin-Ajani, B. L. Ekpukpon, O. A. Odeku, B. I. Olu-Owolabi, and K. O. Adebowale, "Intra and Extragranular Disintegrant Properties of Modified Underutilised Red Lima Bean Starch in Paracetamol Tablet Formulation," *Brazilian J. Pharm. Sci.*, vol. 58, 2022, doi: 10.1590/s2175-97902022e19967.
- [10] L. Lachman and D. Ph, The Theory and Practice of Industrial Pharmacy: Tablets, Gilbert S. Banker, Neil R. Anderson, 3rd ed. Garden City, New York; Livingston, New Jersey; Ridgefield, Connecticut, 1986.
- [11] M. 20852 The United States Pharmacopeial Convention 12601 Twinbrook Parkway, Rockville, Ed., *The United States*
- *Pharmacopeia, USP 41/The National Formulary, NF 36.* Baltimore, MD. Rockville.: United Book Press, Inc., 2018.
 [12] Food and Drug Supervisory Agency of the Republic of Indonesia, *No Title 1200, 2014, BPOM. Obat Tradisional. Mutu.*
- Persyaratan., no. 1200. 2014.
 [13] R. Rahane and P. R. Rachh, "A Review On Fast Dissolving Tablet," J. Drug Deliv. Ther., vol. 8, no. 5, pp. 50–55, Sep. 2018, doi: 10.22270/jddt.v8i5.1888.

8	
0	

- [14] C. P. Jain and P. S. Naruka, "Formulation and Evaluation of Fast Dissolving Tablets of Doxazosin Mesylate," Asian J. Pharm. Res., vol. 6, no. 3, p. 131, 2016, doi: 10.5958/2231-5691.2016.00020.4.
- [15] Allen, L.V., N. G. Popovisch, and H. C. Ansel, *Ansel's Dossage Forms and Drug Delivery Systems, 9th Ed.*, no. 1. 2011.
- [16] T. S. Wahyuni, N. Khoiriyah, L. Tumewu, W. Ekasari, A. Fuad, and A. Widyawaruyanti, "Microscopic and physicochemical evaluation of Ruta angustifolia leaves," *J. Public Health Africa*, vol. 14, no. S1, pp. 123–129, 2023, doi: 10.4081/jphia.2023.2520.
- [17] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia (FHI), II. Jakarta: k, 2017.
- [18] E. Sutrisna, Y. Nuswantoro, and R. F. Said, "Hypolipidemic of ethanolic extract of Salam bark (Syzygium polyanthum (Wight) Walp.) from Indonesia (Preclinical study)," *Drug Invent. Today*, vol. 10, no. 1, pp. 55–58, 2018.
- [19] M. A. Syabana, N. D. Yuliana, I. Batubara, and D. Fardiaz, "α-glucosidase inhibitors from Syzygium polyanthum (Wight) Walp leaves as revealed by metabolomics and in silico approaches," *J. Ethnopharmacol.*, vol. 282, p. 114618, Jan. 2022, doi: 10.1016/j.jep.2021.114618.
- [20] H. Maurya and T. Kumar, "Formulation, standardization, and evaluation of polyherbal dispersible tablet," *Int. J. Appl. Pharm.*, vol. 11, no. 1, pp. 158–167, 2019, doi: 10.22159/ijap.2019v11i1.30113.
- [21] P. Sirame, R. Khairnar, and R. Tasgaonkar, "Granulation," *Int. J. Res. Appl. Sci. Eng. Technol.*, vol. 11, no. 1, pp. 347–355, Jan. 2023, doi: 10.22214/ijraset.2023.48564.
- [22] C. M. Shinko, "Granulation Characterization: Methods and Significance, Dalam: Parikh, D.M.(ed.), Handbook of Pharmaceutical Granulation Technology, Marcel Dekker, New York," p. 422, 1997.
- [23] J. Swarbrick, J. C. Boylan., and 2002, Encyclopaedia of Pharmaceutical Technology, 2nd ed., vol. 3, no. 4. New York: Organic Process Research * Development 2003, 613, 2003.
- [24] D. Puri *et al.*, "Formulation of Herbal Fast Disintegrating Tablets and its ex-vivo Study for Anti-histaminic Activity in Guinea Pig Ileum," *Curr. Clin. Pharmacol.*, vol. 13, no. 2, pp. 128–135, 2018, doi: 10.2174/1574884713666180309153942.
- [25] E. Schlick-Hasper, J. Bethke, N. Vogler, and T. Goedecke, "Flow properties of powdery or granular filling substances of dangerous goods packagings—Comparison of the measurement of the angle of repose and the determination of the Hausner ratio," *Packag. Technol. Sci.*, vol. 35, no. 10, pp. 765–782, Oct. 2022, doi: 10.1002/pts.2678.
- [26] J. Balasubrmaniam, Balasubrmaniam, J. 2009, Influence of Superdisintegrants on The Rate of Drug Dissolution from Oral Solid Dosage Forms, Pharmaceutical Technology, April., April. Pharmaceutical Technology, 2009.
- [27] A. N. Bestari, T. N. S. Sulaiman, and A. Rohman, "Formulasi Orally Disinetgration Tablet (ODT)," vol. 12, no. 2, pp. 453– 465, 2016, doi: 10.22146/farmaseutik.v12i2.26454.
- [28] C. G. Jange, C. R. Wassgren, and K. Ambrose, "The Significance of Tablet Internal Structure on Disintegration and Dissolution of Immediate-Release Formulas: A Review," *Powders*, vol. 2, no. 1, pp. 99–123, Feb. 2023, doi: 10.3390/powders2010008.
- [29] 2010 The Council of Europe: The European Pharmacopoeia, 7th Edition, Strasbourg, France, "Dalam: Brniak, W Jachowicz, R, Pelka, P, The practical approach to the evaluation of methods used to determine the disintegration time of orally disintegrating table," *Saudi Pharm. J.*, vol. 23, no. 4, pp. 437–443, 2015, doi: 10.1016/j.jsps.2015.01.015.
- [30] S. Kakar, R. Singh, and S. Kumar, "Orodispersible tablets: an overview," *MOJ Proteomics Bioinforma.*, vol. 7, no. 3, pp. 180–182, 2018, doi: 10.15406/mojpb.2018.07.00230.
- [31] P. Dey and S. Maiti, "Orodispersible tablets: A new trend in drug delivery," J. Nat. Sci. Biol. Med., vol. 1, no. 1, p. 2, 2010, doi: 10.4103/0976-9668.71663.
- [32] S. U. Pradhan *et al.*, "Characteristics of multi-component formulation granules formed using distributive mixing elements in twin screw granulation," *Drug Dev. Ind. Pharm.*, vol. 44, no. 11, pp. 1826–1837, Nov. 2018, doi: 10.1080/03639045.2018.1503293.

BIOGRAPHIES OF AUTHORS



Widyasari Putranti (D) (S) (S) (S) (C) (C



Desty Restia Rahmawati D S S C was bachelor of faculty of pharmacy, Universitas Ahmad Dahlan, Yogyakarta Indonesia. She is a doctoral Student in Pharmaceutical Sciences Universitas Gadjah Mada. She achieved Best oral Presentation in the 4th International Conference on Pharmacy and Pharmaceutical Science (ICPPS) in Global Front, Surugadai Campus, Meiji University, Tokyo, Japan. She can be contacted by email: desty.restia@gmail.com

https://id.linkedin.com/in/desty-restia-rahmawati-282373193

D 9



Nining Sugihartini 💿 🔀 🖻 🌣 is lecturer in department of technology pharmaceutical Universitas Ahmad Dahlan, Yogyakarta Indonesia. Her research interest in drug and cosmetic development. Currently working on optimization of green tea extract topical dosage formulas as anti-inflammatory. nining.sugihartini@pharm.uad.ac.id

Teuku Nanda Saifullah D S S C is Lecturer in Department of Technology Pharmaceutical Universitas Gadjah Mada, Yogyakarta Indonesia. Medical And Health Sciences/Pharmaceutical Technology, Pharmaceutics Department. His research interest are drug delivery system, technology and formulation of pharmaceutical products. He can be reached by email: <u>in_saifullah@ugm.ac.id</u>

UNIVERSITAS

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Certificate of Acceptance 22666

1 pesan

ijphsiaes.core@gmail.com <ijphsiaes.core@gmail.com> Kepada: widyasari@pharm.uad.ac.id 2 November 2023 pukul 14.05

Dear Author,

Please find attached the certificate of acceptance of your article in our journal.

Certificate of Acceptance ID 22666.pdf





ndexed

CiteScore 2022

6.0

scopus

International Journal of Public Health Science (IJPHS)

CERTIFICATE OF ACCEPTANCE

The manuscript (IJPHS- 22666) entitled:

OPEN ACCESS

SJR Scimago Journal & Country Rank

o.159 0.284 0.284

Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract

Authored by:

Widyasari Putranti, Desty Restia Rahmawati, Nining Sugihartini, Teuku Nanda Saifullah

The manuscript has been accepted in IJPHS (ISSN 2252-8806)

https://ijphs.iaescore.com

November 2, 2023

Prof. Dr. Jay G. Silverman Editor-in-Chief

UNIVERSITAS

widyasari putranti <widyasari@pharm.uad.ac.id>

Fwd: REMINDER [IJPHS] REVISION ID 22666

12 pesan

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id> 25 Oktober 2023 pukul 15.52

------ Forwarded message ------Dari: IJPHS IAES <ijphsiaes.core@gmail.com> Date: Sel, 17 Okt 2023 pukul 14.47 Subject: Re: [IJPHS] REVISION ID 22666 To: widyasari putranti <widyasari@pharm.uad.ac.id>

Thanks for revision. Please revise the following files again.

Pada Sab, 14 Okt 2023 pukul 16.58 widyasari putranti <widyasari@pharm.uad.ac.id> menulis:

<u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

Pada Sel, 10 Okt 2023 pukul 16.28 IJPHS IAES <ijphsiaes.core@gmail.com> menulis: Dear Author,

We inform you that your article needs improvement, with this email we attach the file that needs to be revised and we have provided comments on the file. Please revise the attached file and put a yellow mark on the part you are revising and ask us to edit the Bibtex file. We give until Oct 12, 2023. Thank you very much for your attention.

--

Kind Regards, Maria Ulfah Editorial Staff on behalf of Editor-in-Chief Layout and Editing Team International Journal of Public Health Science Institute of Advanced Engineering and Science (IAES) ISSN: 2252-8806 https://ijphs.iaescore.com/index.php/IJPHS/index email: ijphsiaes.core@gmail.com, ijphs@iaescore.com

UNIVERSITAS AHMAD DAHLAN

Kampus 1: Jln. Kapas No. 9 Yogyakarta

Kampus 2: Jl. Pramuka 42, Sidikan, Umbulharjo, Yogyakarta 55161

Kampus 3: Jl. Prof. Dr. Soepomo, S.H., Janturan, Warungboto, Umbulharjo, Yogyakarta 55164

Kampus 4: Jl.Ringroad Selatan, Yogyakarta

Kampus 5: Jl. Ki Ageng Pemanahan 19, Yogyakarta

Kontak

Email: info@uad.ac.id Telp. : (0274) 563515, 511830, 379418, 371120 Fax. : (0274) 564604

Kind Regards, Maria Ulfah Editorial Staff on behalf of Editor-in-Chief Layout and Editing Team International Journal of Public Health Science Institute of Advanced Engineering and Science (IAES) ISSN: 2252-8806 https://ijphs.iaescore.com/index.php/IJPHS/index email: ijphsiaes.core@gmail.com, ijphs@iaescore.com

Kind Regards, Maria Ulfah Editorial Staff on behalf of Editor-in-Chief Layout and Editing Team International Journal of Public Health Science Institute of Advanced Engineering and Science (IAES) ISSN: 2252-8806 https://ijphs.iaescore.com/index.php/IJPHS/index email: ijphsiaes.core@gmail.com, ijphs@iaescore.com

22666.docx W 4362K

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

Article was revised and upload on ojs [Kutipan teks disembunyikan]

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

Thanks for the information.

Here is the file we have checked and there are still some things you need to revise. please find the trealmpir file. we give you time to revise until October 28, 2023.

[Kutipan teks disembunyikan]

W	22666.docx
ΞE	4364K

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

I've revised <u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

[Kutipan teks disembunyikan]

25 Oktober 2023 pukul 17.13

26 Oktober 2023 pukul 07.55

28 Oktober 2023 pukul 11.52

22666 rev 28okt.docx 4370K

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

30 Oktober 2023 pukul 08.23

Thanks for revision..

However, your colleague Desty Restia Rahmawati has not included an Orcid ID. If you don't have one, please create one first.

[Kutipan teks disembunyikan]

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

i've revised (add orchid id desty restia R on article) <u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

[Kutipan teks disembunyikan]

22666 rev 28okt.docx 4370K

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

Thank you. We will inform you again if there is an update. [Kutipan teks disembunyikan] 30 Oktober 2023 pukul 09.36

30 Oktober 2023 pukul 15.05

IJPHS IAES <ijphsiaes.core@gmail.com> 15 Desember 2023 pukul 16.57 Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>, desty.restia@gmail.com, tn_saifullah@ugm.ac.id

Dear Author,

We inform you that your article needs improvement, with this email we attach the file that needs to be revised and we have provided comments on the file. Please revise the attached file and put a yellow mark on the part you are revising and ask us to edit the Bibtex file. We give until Dec 17, 2023. Thank you very much for your attention.

NOTE: Please revise using the file that we attached earlier. Do not create new files.

[Kutipan teks disembunyikan]

53. 22666 edit sept fin Ez.docx

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

I've revised <u>Widyasari Putranti, M.Sc., Apt</u> Departement of Technology Pharmaceutical Pharmacy Faculty of Ahmad Dahlan University Yogyakarta Indonesia Hp 081.328.899.770

[Kutipan teks disembunyikan]

18 Desember 2023 pukul 10.38

53. 22666 edit sept fin Ez.docx 4349K

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

add ID Orcid Desty Restia Rahmawati. We look forward to hearing from you soon. Thanks

[Kutipan teks disembunyikan]

widyasari putranti <widyasari@pharm.uad.ac.id> Kepada: IJPHS IAES <ijphsiaes.core@gmail.com>

Desty = 0009-0007-9290-6369 [Kutipan teks disembunyikan]

IJPHS IAES <ijphsiaes.core@gmail.com> Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>

Thanks. [Kutipan teks disembunyikan] 18 Desember 2023 pukul 11.50

18 Desember 2023 pukul 12.49

18 Desember 2023 pukul 14.13

Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract

Widyasari Putranti¹, Desty Restia Rahmawati¹, Nining Sugihartini¹, Teuku Nanda Saifullah²

¹Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia ²Department of Pharmaceutical Technology, Faculty of Pharmacy, Gadjah Mada University, Yogyakarta, Indonesia

Article Info ABSTRACT

Article history:

Received month dd, yyyy Revised month dd, yyyy Accepted month dd, yyyy

Keywords:

Croscarmellose Superdisintegrant Syzygium polyanthum extract Fast disintegrating tablet Intra granular Extra granular Bay leaves (Syzygium polyanthum) contain the flavonoid quercetin which can be used as an antihyperlipidemic drug. The development of antihyperlipidemic drug formula in the form of fast disintegrating tablet (FDT) is needed for patients who experience dysphagia. FDT preparations require an optimal super disintegrant concentration to produce a good drug formula. This study aims to develop the FDT formula of bay leaves extract using the super disintegrant croscarmellose sodium (CCS) intra and extragranular. FDT formulation using the wet granulation method with variations of CCS concentrations; F1: 2%, F2: 3.5%, and F3: 5% for extra granular, and 2% for intra granular. The formulation process, in-process control (IPC) granules, weight uniformity tests, and various physical properties tests of tablets were carried out. Data were statistically analyzed using one way ANOVA test (α =95%). The results of statistical tests of IPC granules, uniformity of weight, and tablet size of all FDT formulas were not significantly different (p>0.05). The CCS concentration for extra granular significantly affected the wetting time, disintegration time, hardness, and the value of friability percentage of FDT (p<0.05). The combination of intra and extra-granular CCS (2%:5%) gave the most optimum physical properties of bay leaf extract FDT.

This is an open access article under the <u>CC BY-SA</u> license.



Corresponding Author:

Widyasari Putranti Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan Yogyakarta, Indonesia Email: widyasari@pharm.uad.ac.id

1. INTRODUCTION

Plant germplasm resources are raw materials for medicines, one of which is a bay leaf (*Syzygium polyanthum*). Bay leaves are known to contain secondary metabolites, such as saponins, terpenoids, flavonoids (quercetin), polyphenols, alkaloids, steroids, and essential oils (sesquiterpenes)[1]-[2]. Based on the Decree of the Minister of Health of the Republic of Indonesia (2009), bay leaves contain a total flavonoid of not less than 0.40% which is calculated as quercetin [3]. Quercetin in bay leaves the potential as an antihyperlipidemic which can significantly reduce levels of triglycerides, total cholesterol, and low-density lipoprotein (LDL) cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in high-density lipoprotein (HDL), as well as inhibiting 3-hidroksi-3-metilglutaril coenzyme A (HMG-CoA), reductase, and LDL oxidation [4].

Conventional drugs for hyperlipidemia are generally available in tablet form. However, the tablet dosage form has several disadvantages. Some of them require a long absorption time in the body, and slow drug action, and elderly patients may experience difficulty swallowing or dysphagia [5]. Seeing the problems that arise from conventional tablets, the herbal fast disintegrating tablet (FDT) of bay leaf extract is the right alternative so that the drug can be used comfortably and has a therapeutic effect more quickly. FDT is a solid

1

dosage form containing pharmaceutically active substances that can be disintegrated quickly, generally within seconds when placed on the tongue. This properties of FDT can release the active drug substance immediately, speed up the onset, increase the oral bioavailability of the drug in the body so that therapeutic effectiveness can be achieved [6]-[7]. FDT preparations combine two advantages of preparations, namely liquid which can increase the solubility and bioavailability of drugs, as well as the advantages of solid dosage forms that have high physicochemical stability, relatively constant homogeneity of the active ingredients of the preparation, and easier manufacturing.

FDT is formulated with croscarmellose sodium (CCS) super disintegrant which is classified as a strong swelling product with a good water absorption mechanism [7]-[8]. CCS is effectively used in FDT preparations because it rapidly expands to 4-8 times its initial volume when in contact with water, thus accelerating the tablet-crushing process. CCS is effective intra-granular which allows for better capillary and swelling processes, causing the tablet to disintegrate quickly and reduce disintegration time [8]. The addition of intra and extra-granular crushing materials can increase the effectiveness of tablet disintegration through two stages, i) extra granular which breaks the tablet into granules, and ii) intra granular breaks down the granules into smaller particles. The mechanism is crushing material that experiences rapid swelling and water absorption (wicking) through the gaps between the granules and the particles that make up the granules [9]. This study aims to develop herbal FDT with the active ingredient of bay leaf ethanol extract. The concentration of CCS super disintegrant intra and extra-granular needs to be determined to obtain the FDT of bay leaf extract with optimum physical properties.

2. METHOD

The ingredients used were bay leaf extract in the form of fresh bay leaf collected from tradisional market, Beringharjo market, Yogyakarta, Indonesia, ethanol 70%, croscarmellose sodium (PT. Phapros Tbk), lactose, magnesium stearate, aspartame, menthol and corn starch paste (pharmaceutical grade). The tools for extraction used are oven, rotary evaporator calipers, waterbath, vacuum pump. The tools for granule physical properties test used are Ohaus analytical balance, absorption test equipment granules, volumenometer (tapped density tester), flow tester granule, sieving machine, dry granule sieve, wet granule sieve, pan, halogen moisture analyzer. the tools for tablet test used are single punch tablet printing machine, hardness tester, friability tester, and disintegration tester.

2.1. Bay leaf ethanol extract

Bay leaves were dried in an oven at 50-60°C until a moisture content of less than 10% was obtained. Simplicia were extracted by maceration method using 70% ethanol solvent (1:10) (w/v) [3]. Maserati and pulp were separated by the vacuum pump and filter paper. The solvent was evaporated using a rotary evaporator. The viscous extract obtained was calculated for its yield.

2.2. FDT formulation of bay leaf extract

FDT was made by the wet granulation method. At the granulation stage, bay leaves extract was mixed with intra granular CCS, lactose and corn starch. The mixture was sieved through a wet (no. 18 mesh), ovendried (50-60°C), then sifted dry (no. 20 mesh). The granules and excipients consisting of magnesium stearate, aspartame, menthol, and CCS (extra granular) mixed until homogeneous (30 minutes). Mixture of granules and excipients that have been homogenized are compressed using a single punch tablet press.

2.3. Granule physical properties test parameters

The moisture content of granules can be determined based on the value of loss on drying (LOD) and MC. LOD is a test for measuring the difference in heavy total granules when before and after drying or a moisture content statement based on wet weight. Meanwhile, MC is a statement of moisture content based on dry weight. Measurements using the halogen moisture analyzer tool. Good granule moisture content if the LOD and MC values are 10% [10].

The flow properties of the granules can be tested by calculating the angle of repose, the flow time test, and the determination index (Hausner ratio). The granule angle of repose α (degree) is said to be very good (25°-30°); good (30-35°) or 25°><35°. A good granule flow time is not more than 10 seconds for 100 grams of granules [11]. Excellent granule setting index (\leq 10); good (11–15), and the Hausner ratio index that meets the requirements for the physical properties of the granule mass, which is very good (1.00–1.11); good (1.12–1.18).

Sieve no. 14, 16, 20, 30, 50 mesh and pan, installed in stages on the sieving machine (50 amplitude, 15 minutes). The granules left on each sieve were weighed and the percentage was calculated [10]. The fragility of the granules is based on the number of fines that occur after the friability test by sieving. The granules were

D 3

put into a graded sieve with the top sieve no. 30 mesh and bottom pan, using a sieving machine (50 amplitude, 30 minutes). The water absorption of the granules affects the tablet's disintegration time (disintegration). It was carried out with a series of absorption test equipment by placing 300 mg of granules on a water-saturated sheath paper. The absorbed water was measured using a digital scale for 15 minutes or until it was constant [7].

2.4. Physical properties test parameters of FDT

Requirements and weight uniformity tests are determined based on the regulation of the National Food and Drug Agency of the Republic of Indonesia concerning the quality requirements of traditional medicines. The test was carried out by weighing 20 tablets one by one, from 20 tablets no more than 2 tablets, each of which deviated from the average weight by more than 5% and not one tablet whose weight deviated from the average weight. the average is greater than 10% [12].

Each tablet was measured in diameter in a horizontal position with a caliper, and the thickness of the tablet was measured in a vertical position with a caliper (millimeters). The tablet hardness test was carried out by taking a sample of at least 6 tablets from each formula. The tablet hardness test is carried out using hardness tester tablets. One by one the tablets are tested by placing the tablets in the center perpendicular to the hardness tester. A good FDT hardness is 3-5 kg/cm² [11], [13].

For tablets with a single weight of exactly or less than 650 mg, the total weight of the tested samples was close to 6.5 grams. Prior to testing, the tablets were dusted and weighed. All tablets were put into the friability tester (speed 25 rpm, 4 minutes). The tablets were cleaned of adhering fines and re-weighed. A good friability value should not be more than 1% [11].

The wetting time test was done by placing a sheet of filter paper that has been folded twice into a petri dish with a diameter of 5 cm. The petri dish was filled with 5.0 mL of distilled water containing the strawberry red dye. A tablet is then placed on the filter paper, and simultaneously with the start of the test instrument, the stopwatch is turned on. Wetting time was calculated as the time required for a red color to appear on the entire surface of the tablet [14]. A total of 6 tablets of each formula were placed in each tube and a disk was placed on it, the test was carried out with water medium at a temperature of 37 ± 2 °C or 35-39 °C using a 1000 ml glass beaker. The disintegration time requirement for FDT preparations is at least <1 minute disintegration time [11], [15].

2.5. Data analysis

The results of the physical examination of granules and tablets of FDT bay leaf extract were analyzed using the one-way ANOVA test method with a 95% confidence level.

3. RESULTS AND DISCUSSION

Bay leaf simplicia was tested for macroscopic (organoleptic) specifications on fresh and dried simplicia, as well as microscopic tests for simplicia parts and identification fragments of simplicia powder. Fragments prepared by dissolving in chloralhydrate and water [16]. The simplicia specification test was carried out to ensure the correctness of the identity and quality of the bay leaf simplicia based on the Indonesian Herbal Pharmacopoeia compendia and Indonesian Materia Medika. Based on the microscopic test results on Figure 1 there are identifier fragments namely i) lower epidermis with stomata, ii) upper epidermis, iii) elements with dots, iv) prism-shaped calcium oxalate crystals, and v) sclerenchyma. The results of the specification test for the simplicia of bay leaves showed that the simplicia used met the requirements of organoleptic and microscopic terms. The determination test of bay leaf simplicia at the Biology Laboratory of Universitas Ahmad Dahlan (UAD) was carried out to ensure that the selected bay leaf simplicia was in accordance with its species or identity [3], [17].

3.1. Extraction bay leaf

Fresh simplicia of bay leaves were sorted wet, dried using an oven $(50-60^{\circ}C)$ with a moisture content of not more than 10%, then sorted dry. Simplicia powder for extract preparation is a fine simplicial (No. 60 mesh) in order to optimize the extraction process [18]. The results of the simplicial extraction of bay leaves weighing 3.7 kg obtained a thick extract of 347 grams. The total yield of the thick extract obtained was 9.38%. The morphology of the viscous extract is shown in Figure 2. Screening and determination of the chemical compound flavonoid quercetin contained in the thick extract of bay leaves was carried out at the Integrated Research Laboratory of the Faculty of Pharmacy, UAD. According to the Ministry of Health (2017), bay leaves contain a total flavonoid of not less than 0.40% calculated as quercetin. Based on the results of the test the flavonoid content of 0.8079±0.0045%, so that it meets the specifications for the total flavonoid content of bay leaves[17]. Quercetin is an active substance that has medicinal effects such as antiherlipidemia, antibacterial, antidiabetic, antioxidant [18], [19].



Figure 1. Identification fragments of bay leaf; (a) lower epidermis, (b) upper epidermis, (c) floem, (d) calcium oxalate crystal, (e) sclerenchyma



Figure 2. Bay leaf Extract

3.2. FDT formulation

FDT formulation method with wet granulation is good used for water-resistant active ingredients and heating temperature [10]. The active ingredient in bay leaf is quercetin (3,3',4',5,7-pentahydroxyflavone) is a flavonoid derivative compound that is resistant to heating with a melting point of 316.5 °C [20], while the wet granulation process ranges from at a temperature of 40-60 °C. The wet granulation method can improve the flowability and compatibility of the material and facilitate the FDT formulation process from the active ingredient in the form of a thick extract [21]. Wet granulation has the advantage of good content distribution and uniformity, besides that the hydrophobic surface of the tablet becomes more hydrophilic which affects the water absorption process into the tablet, thereby increasing the speed of tablet disintegration [22]. The work of woven fibers between granules can increase the speed of tablet disintegration[23], it is synergistic with the presence of CCS super disintegrant extra granular which facilitates swelling and absorption of water into granules and their small particles.

The concentration of CCS super disintegrant intra-granular of the three formulas was the same, 2% (10 mg/tablet), and extra granular was formulated with varying levels, F1 2% (10), F2 3.5% (17.5), and F3 5% (25 mg/tablet). The FDT formula for bay leaf extract is shown in Table 1. Based on the Handbook of pharmaceutical excipients the use of CCS in the wet granulation process should be added in the wet or granulation (intra-granular) stage, as well as in the dry (extra-granular) process [21]. Because this method can increase the wicking ability (water absorption) and swelling of the CCS super disintegrant.[8]

Table 1. I official of I DI bay leaf extract				
Ingredient		position	(mg)	
	F1	F2	F3	
Bay leaf extract	70	70	70	
Croscarmellose sodium (intra granular)	10	10	10	
Lactose	380	374	366	
Croscarmellose sodium (extra granular)	10	17.5	25	
Magnesium stearate	5	5	5	
Aspartame	20	20	20	
Menthol	1	1	1	
Corn starch paste 5% w/v	qs	qs	qs	
Total weight (mg)	500	500	500	

Table 1. Formula of FDT bay leaf extract

3.2.1. Granule physical properties test parameters

One of the most crucial elements of wet granulation is the water content, MC and LOD has been was measured and the values of granules from all formulas as shown in Table 2, the results obtained meet the requirements of 10% (p>0.05). Water content will affect granule characteristics, including granule size and compressibility. The flow properties of the granules show the results of the angle of repose of the granules ranging from 30-35 (good), granule flow time of 10-14 grams/second (good), and the determination test/Hausner ratio (very good). The flow properties of granules are very influential on the tablet compression process, granules with good flow properties will cause the granules to enter the die space to be relatively constant so that the tablet weights with small weight variations can be obtained and can increase the uniformity of the resulting tablet dosage [8], [24], [25].

The average diameter of all granules ranged from 650-700 m (p>0.05) and the value of friability percentage of the granules ranged from 0.5-0.9% (p>0.05)[13]. The granule absorption test for all formulas had absorption between 0.120-0.190 mg/minute (p>0.05). The granule absorption shows the speed of the granules in absorbing water to crush the granules into small particles so as to speed up the tablet disintegration time. All in process control (IPC) granule parameters have met the requirements so that the tableting process can be continued with a single punch tablet press with a punch pressure of 13 kg.

Table 2. Results of examination of physical properties of FDT granules of bay leaf extract

Test parameters	Formula				
	F1	F2	F3		
LOD±SD (%)*	7.03±0.76	6.29±1.28	6.57±1.18		
MC±SD (%)*	2.98 ± 0.17	2.35±0.33	2.69±0.31		
Angle of repose±SD (°)*	30.66±1.75	31.18±2.92	30.08±2.30		
Flow time±SD (grams/second)*	12.98 ± 1.28	13.12±0.89	11.85±0.48		
Assignment±SD (%)*	3.17±1.04	2.83±1.04	3.83±0.76		
Hausner ratio Index±SD*	1.03 ± 0.01	1.03 ± 0.01	1.04 ± 0.01		
Average diameter±SD (µm)*	694.64 ± 28.87	657.58±39.65	698.14±26.09		
Fragility±SD (%)*	0.8028 ± 0.20	0.8199 ± 0.05	0.5863 ± 0.04		
Absorption±SD (mg/min)*	0.186 ± 0.05	0.188 ± 0.05	0.127±0.01		

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

3.2.2. Physical properties test parameters of FDT

The uniformity of tablet weight is related to the uniformity of active substance levels and therapeutic effects. The uniformity of tablet weight (Table 3) can be seen from the coefficient of variation F1, F2, and F3 all <5%, (0.0942, 0.1246, and 0.2103) %, respectively. The weight uniformity test based on the National Food and Drug Agency of the Republic of Indonesia requirements, shows the results that meet the requirements, where the tablet meets the specified weight deviation limit value range.

Hardness ranges FDT be good tablet from 3-5 kg/cm² [13], while the conventional tablet, is 4-8 kg/cm². The FDT hardness as shown in Table 3 shows results that meet the requirements. Hardness affects the brittleness and disintegration time of tablets, as well as the dissolution/release of the active drug substance [8]. The results of statistical analysis of tablet hardness with sig. (p<0.05) indicates that the hardness of the three FDT formulas is significantly different. The hardness of FDT tablets is strongly influenced by the CCS super disintegrant excipient. FDT (F3) produced the hardest tablet with the highest extra-granular CCS concentration (5%), causing the particles in the tablet to be tightly bound. Croscarmellose sodium (Ac-Di-Sol) is a super disintegrant with a fibrous cross-linked polymer structure that functions as a disintegrant with a strong swelling mechanism, fiber is able to bind strongly to the particles so as to increase the hardness and reduce the brittleness of the tablet [26]. This causes the tablet to have good compatibility, so there is a tendency

to increase hardness as the amount of CCS in the tablet increases [27]. Another factor that affects the hardness of FDT is the upper punch pressure during the tableting process, the greater the pressure used, the harder the FDT will be. The results of the diameter and thickness measurements of the tablets showed that the size of the entire tablet was uniform (homogeneous).

The friability value of the tablet is based on the requirements, which should not be more than 1% [11]. The fragility value of FDT is shown in Table 3 with a friability value of <1% so that FDT meets the requirements. The results of the tablet friability statistical test have a sig. value (p < 0.05) means that the fragility value between formulas is significantly different. CCS super disintegrant is a factor influencing fragility. Formula 3 with the highest number of CCS has the smallest friability value, where there is a tendency to decrease the friability value as the number of CCS in the tablet increases. The friability value is inversely proportional to the tablet's hardness, where the harder a tablet is, the smaller the friability value will be.

FDT wetting time is shown in Table 3, FDT (F3) with the highest number of CCS has the smallest wetting time. Wetting time is a parameter to determine the speed of FDT in absorbing water, which affects the speed of tablet disintegration. The faster the wetting time, the faster the tablet disintegration. The wetting time of tablets is influenced by the structure of the tablet matrix and the hydrophilicity of the excipients [28]. The result of statistical analysis of FDT wetting time has a sig. value (p<0.05) means that the formulas differ significantly. FDT (F3) with the highest number of CCS had the fastest time for wettable tablets, super disintegrant CCS being the most influential factor. The cross-linked chemical structure of CCS creates excipients that have hydrophilic properties and very easy-to-absorb solvents resulting in outstanding swelling properties in the disintegration process.

Table 3. Results of examination of physical properties of bay leaf extract FDT tablets

Test Demonstern		Note.		
Test Parameters	F1	F2	F3	
Weight uniformity±SD (mg)	50 4.42±0.48	50 4.78±0.63	50 4.43±1.06	Appropriate*
Diameter±SD (mm)	12.00±0.00	12.00±0.00	12.00±0.00	Appropriate*
Thickness±SD (mm)	3.67±0.00	3.67±0.00	3.68±0.01	Appropriate*
Hardness±SD (kg)	4.46±0.25°	4.50±0.27°	5.24±0.32 ^{a,b}	In accordance
Fragility±SD (%)	0.77±0.05°	0.72±0.04°	$0.52 \pm 0.04^{a,b}$	In accordance
Wetting time±SD (seconds)	50.33±3.06°	43.33±4.04°	20.00±2.65 ^{a,b}	In accordance
Disintegration time±SD (seconds)	45.33±5.86°	38.33±3.79°	19.67±3.06 ^{a,b}	In accordance

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

(a) means a formula that is significantly different from the formula 1

(b) means a formula that is significantly different from the formula 2

(c) means a formula that is significantly different from formula 3

F1: CCS intra granular 2%: extra granular 2%

F2: CCS intra granular 2%: extra granular 3.5%

F3: CCS intra granular 2%: extra granular 5%

Disintegration (disintegration) time FDT has at least a disintegration time of less than 1 minute [15]. Another source says that in-vitro about 30 seconds or less [29], whereas according to Ph. euros. is <3 min in the oral cavity before swallowing [29], [30]. The faster disintegration time will increase the speed of drug release from the tablet which affects the effectiveness of therapy. The results of the disintegration time of FDT in table 3 show that it meets the requirements of the disintegration time of FDT (<1 minute). FDT disintegration time test is shown in Table 3 with FDT (F3) having the fastest disintegration time, due to the higher levels of CCS used in F3 than F1 and F2. The results of the statistical test of tablet disintegration time have a sig value (p < 0.05) means that each formula is significantly different. CCS has a dual mechanism, namely water wicking and rapid swelling [8]. The highly porous form of the structure speeds up the disintegration time, because water quickly enters the tablet and increases the tablet wetting rate, through the gaps between the granules and the granule pores [31]. This is related to the factor of the number of CCS super disintegrant excipient used in each formula intra and extra-granular. The use of intra and extra-granular affects the decrease in tablet disintegration time through the swelling mechanism, and along with the increase in the extra-granular CCS concentration, the disintegration time will be faster. The FDT disintegration pattern is also facilitated by intra-granular CCS which causes the tablets to be finely crushed not in the form of granules but into small particles so that the speed of disintegration is optimal. The optimum formula for FDT of bay leaf extract is shown in Table 4 with the results of all evaluations of the physical properties of the granules and FDT (F3) that meet the requirements and show the optimum results of physical parameters.

The results of this study are in line with the research of Puri *et al.* [24], that the combination of intragranular and extra-granular can produce tablets with the most effective disintegration time by providing better tablet physical properties, compared to intra-granular or extra-granular only. Extra-crushing material granules

D 7

have a high tendency to absorb water from the surrounding liquid through the tablet-breaking mechanism [32]. Tablets with disintegrant intra-granular have a higher hardness than tablets with extra granular crushing agents. By combining intra and extra granular super disintegrants, a good level of tablet hardness can be obtained so that the tablet is not brittle and has a faster disintegration time with the mechanism of tablet disintegration into granules (extra granular factor) and granules into small particles (intra granular factor).

1 u 0 0 - 1, $u 1 0 u 0 p u 1 u 1 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0 0$	Ta	ble	4.	Va	arious	parameters	of	01	otimized	tał	olet	form	ıla	(F3)	3)	
--	----	-----	----	----	--------	------------	----	----	----------	-----	------	------	-----	------	----	--

No.	Test parameters	Optimized formula value
1.	Weight uniformity±SD (mg)	504.43±1.06
2.	Diameter±SD (mm)	12.00±0.00
3.	Thickness±SD (mm)	3.68±0.01
4.	Violence±SD (kg)	5.24±0.32
5.	Fragility±SD (%)	0.52±0.04
6.	Wetting time±SD (seconds)	20.00 ±2.65
7.	Disintegration time±SD	19.67±3.06
	(seconds)	

4. CONCLUSION

In the FDT research, to improve the poor flow properties of the extract, the wet granulation method was carried out. The IPC results for the three formulas showed granules that met the requirements. This research provides information that the use of extra granular CCS at higher levels will increase the rate of disintegration. The combination of CCS intra granular 10 mg (2%) and extra granular 25 mg (5%) can produce FDT with the most effective. By combining intra and extra granular super disintegration time. In addition, FDT bay leaves extract can be used as an alternative treatment for hyperlipidemia in patients who experience dysphagia. This research still needs development to determine the stability of FDT bay leaves extract where the active substance is herbal medicine.

ACKNOWLEDGEMENTS

This research was carried out with the assistance of the Higher Education Research Grant No PKPT-013/SKPP/III/2018.

REFERENCES

- L. Hartanti, S. M. K. Yonas, J. J. Mustamu, S. Wijaya, H. K. Setiawan, and L. Soegianto, "Influence of extraction methods of bay leaves (Syzygium polyanthum) on antioxidant and HMG-CoA Reductase inhibitory activity," *Heliyon*, vol. 5, no. 4, p. e01485, Apr. 2019, doi: 10.1016/j.heliyon.2019.e01485.
- [2] E. Kato, R. Nakagomi, M. D. P. T. Gunawan-puteri, and J. Kawabata, "Identification of hydroxychavicol and its dimers, the lipase inhibitors contained in the Indonesian spice, Eugenia polyantha," *Food Chem.*, vol. 136, no. 3–4, pp. 1239–1242, 2013, doi: 10.1016/j.foodchem.2012.09.013.
- [3] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia, 1st ed. Jakarta Indonesia, 2009.
- [4] K. Zeka, K. Ruparelia, R. Arroo, R. Budriesi, and M. Micucci, "Flavonoids and Their Metabolites: Prevention in
- Cardiovascular Diseases and Diabetes," Diseases, vol. 5, no. 3, p. 19, 2017, doi: 10.3390/diseases5030019.
- [5] I. Safitri, S. Sulistiyaningsih, and A. Y. Chaerunisaa, "Review: Superdisintegran dalam Sediaan Oral," *Farmasetika.com* (Online), vol. 4, no. 3, p. 56, 2019, doi: 10.24198/farmasetika.v4i3.22945.
- [6] A. Masih, A. Kumar, S. Singh, and A. K. Tiwari, "Fast Dissolving Tablets: A Review," *Int. J. Curr. Pharm. Res.*, vol. 9, no. 2, p. 8, Mar. 2017, doi: 10.22159/ijcpr.2017v9i2.17382.
 [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam
- [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam leaves ethanol extract with combination of crospovidone and croscarmellose sodium as superdisintegrants," *J. Sains Farm. Klin.*, vol. 8, no. 3, p. 285, 2021, doi: 10.25077/jsfk.8.3.285-295.2021.
- [8] M. E. Rowe, R. C., Sheskey, P. J. dan Quinn, *Handbook of Pharmaceutical Excipients*, Sixth. London: Pharmaceutical press., 2009.
- [9] M. O. Okekunle, O. D. Akin-Ajani, B. L. Ekpukpon, O. A. Odeku, B. I. Olu-Owolabi, and K. O. Adebowale, "Intra and Extragranular Disintegrant Properties of Modified Underutilised Red Lima Bean Starch in Paracetamol Tablet Formulation," *Brazilian J. Pharm. Sci.*, vol. 58, 2022, doi: 10.1590/s2175-97902022e19967.
- [10] L. Lachman and D. Ph, The Theory and Practice of Industrial Pharmacy: Tablets, Gilbert S. Banker, Neil R. Anderson, 3rd ed. Garden City, New York; Livingston, New Jersey; Ridgefield, Connecticut, 1986.
- [11] M. 20852 The United States Pharmacopeial Convention 12601 Twinbrook Parkway, Rockville, Ed., *The United States*
- *Pharmacopeia, USP 41/The National Formulary, NF 36.* Baltimore, MD. Rockville.: United Book Press, Inc., 2018.
 [12] Food and Drug Supervisory Agency of the Republic of Indonesia, *No Title 1200, 2014, BPOM. Obat Tradisional. Mutu.*
- Persyaratan., no. 1200. 2014.
 [13] R. Rahane and P. R. Rachh, "A Review On Fast Dissolving Tablet," J. Drug Deliv. Ther., vol. 8, no. 5, pp. 50–55, Sep. 2018, doi: 10.22270/jddt.v8i5.1888.

8	
0	

- [14] C. P. Jain and P. S. Naruka, "Formulation and Evaluation of Fast Dissolving Tablets of Doxazosin Mesylate," Asian J. Pharm. Res., vol. 6, no. 3, p. 131, 2016, doi: 10.5958/2231-5691.2016.00020.4.
- [15] Allen, L.V., N. G. Popovisch, and H. C. Ansel, *Ansel's Dossage Forms and Drug Delivery Systems, 9th Ed.*, no. 1. 2011.
- [16] T. S. Wahyuni, N. Khoiriyah, L. Tumewu, W. Ekasari, A. Fuad, and A. Widyawaruyanti, "Microscopic and physicochemical evaluation of Ruta angustifolia leaves," *J. Public Health Africa*, vol. 14, no. S1, pp. 123–129, 2023, doi: 10.4081/jphia.2023.2520.
- [17] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia (FHI), II. Jakarta: k, 2017.
- [18] E. Sutrisna, Y. Nuswantoro, and R. F. Said, "Hypolipidemic of ethanolic extract of Salam bark (Syzygium polyanthum (Wight) Walp.) from Indonesia (Preclinical study)," *Drug Invent. Today*, vol. 10, no. 1, pp. 55–58, 2018.
- [19] M. A. Syabana, N. D. Yuliana, I. Batubara, and D. Fardiaz, "α-glucosidase inhibitors from Syzygium polyanthum (Wight) Walp leaves as revealed by metabolomics and in silico approaches," *J. Ethnopharmacol.*, vol. 282, p. 114618, Jan. 2022, doi: 10.1016/j.jep.2021.114618.
- [20] H. Maurya and T. Kumar, "Formulation, standardization, and evaluation of polyherbal dispersible tablet," *Int. J. Appl. Pharm.*, vol. 11, no. 1, pp. 158–167, 2019, doi: 10.22159/ijap.2019v11i1.30113.
- [21] P. Sirame, R. Khairnar, and R. Tasgaonkar, "Granulation," *Int. J. Res. Appl. Sci. Eng. Technol.*, vol. 11, no. 1, pp. 347–355, Jan. 2023, doi: 10.22214/ijraset.2023.48564.
- [22] C. M. Shinko, "Granulation Characterization: Methods and Significance, Dalam: Parikh, D.M.(ed.), Handbook of Pharmaceutical Granulation Technology, Marcel Dekker, New York," p. 422, 1997.
- [23] J. Swarbrick, J. C. Boylan., and 2002, Encyclopaedia of Pharmaceutical Technology, 2nd ed., vol. 3, no. 4. New York: Organic Process Research * Development 2003, 613, 2003.
- [24] D. Puri *et al.*, "Formulation of Herbal Fast Disintegrating Tablets and its ex-vivo Study for Anti-histaminic Activity in Guinea Pig Ileum," *Curr. Clin. Pharmacol.*, vol. 13, no. 2, pp. 128–135, 2018, doi: 10.2174/1574884713666180309153942.
- [25] E. Schlick-Hasper, J. Bethke, N. Vogler, and T. Goedecke, "Flow properties of powdery or granular filling substances of dangerous goods packagings—Comparison of the measurement of the angle of repose and the determination of the Hausner ratio," *Packag. Technol. Sci.*, vol. 35, no. 10, pp. 765–782, Oct. 2022, doi: 10.1002/pts.2678.
- [26] J. Balasubrmaniam, Balasubrmaniam, J. 2009, Influence of Superdisintegrants on The Rate of Drug Dissolution from Oral Solid Dosage Forms, Pharmaceutical Technology, April., April. Pharmaceutical Technology, 2009.
- [27] A. N. Bestari, T. N. S. Sulaiman, and A. Rohman, "Formulasi Orally Disinetgration Tablet (ODT)," vol. 12, no. 2, pp. 453– 465, 2016, doi: 10.22146/farmaseutik.v12i2.26454.
- [28] C. G. Jange, C. R. Wassgren, and K. Ambrose, "The Significance of Tablet Internal Structure on Disintegration and Dissolution of Immediate-Release Formulas: A Review," *Powders*, vol. 2, no. 1, pp. 99–123, Feb. 2023, doi: 10.3390/powders2010008.
- [29] 2010 The Council of Europe: The European Pharmacopoeia, 7th Edition, Strasbourg, France, "Dalam: Brniak, W Jachowicz, R, Pelka, P, The practical approach to the evaluation of methods used to determine the disintegration time of orally disintegrating table," *Saudi Pharm. J.*, vol. 23, no. 4, pp. 437–443, 2015, doi: 10.1016/j.jsps.2015.01.015.
- [30] S. Kakar, R. Singh, and S. Kumar, "Orodispersible tablets: an overview," *MOJ Proteomics Bioinforma.*, vol. 7, no. 3, pp. 180–182, 2018, doi: 10.15406/mojpb.2018.07.00230.
- [31] P. Dey and S. Maiti, "Orodispersible tablets: A new trend in drug delivery," J. Nat. Sci. Biol. Med., vol. 1, no. 1, p. 2, 2010, doi: 10.4103/0976-9668.71663.
- [32] S. U. Pradhan *et al.*, "Characteristics of multi-component formulation granules formed using distributive mixing elements in twin screw granulation," *Drug Dev. Ind. Pharm.*, vol. 44, no. 11, pp. 1826–1837, Nov. 2018, doi: 10.1080/03639045.2018.1503293.

BIOGRAPHIES OF AUTHORS



Widyasari Putranti (D) (S) (S) (S) (C) (C



Desty Restia Rahmawati D S S C was bachelor of faculty of pharmacy, Universitas Ahmad Dahlan, Yogyakarta Indonesia. She is a doctoral Student in Pharmaceutical Sciences Universitas Gadjah Mada. She achieved Best oral Presentation in the 4th International Conference on Pharmacy and Pharmaceutical Science (ICPPS) in Global Front, Surugadai Campus, Meiji University, Tokyo, Japan. She can be contacted by email: desty.restia@gmail.com

https://id.linkedin.com/in/desty-restia-rahmawati-282373193

D 9



Nining Sugihartini 💿 🔀 🖻 🌣 is lecturer in department of technology pharmaceutical Universitas Ahmad Dahlan, Yogyakarta Indonesia. Her research interest in drug and cosmetic development. Currently working on optimization of green tea extract topical dosage formulas as anti-inflammatory. nining.sugihartini@pharm.uad.ac.id

Teuku Nanda Saifullah D S S C is Lecturer in Department of Technology Pharmaceutical Universitas Gadjah Mada, Yogyakarta Indonesia. Medical And Health Sciences/Pharmaceutical Technology, Pharmaceutics Department. His research interest are drug delivery system, technology and formulation of pharmaceutical products. He can be reached by email: <u>in_saifullah@ugm.ac.id</u>

UNIVERSITAS AHMAD DAHLAN

widyasari putranti <widyasari@pharm.uad.ac.id>

[IJPHS] Schedule for Publication for March 2024 Issue

1 pesan

IJPHS IAES <ijphsiaes.core@gmail.com>

21 Desember 2023 pukul 11.30 Kepada: widyasari putranti <widyasari@pharm.uad.ac.id>, desty.restia@gmail.com, nining.sugihartini@pharm.uad.ac.id, tn saifullah@ugm.ac.id

Dear Author(s),

We are glad to inform you that IJPHS Vol 13, No 1: March 2024 has been published. Kindly click the link to check your paper. https://ijphs.iaescore.com/index.php/IJPHS/issue/view/578 The editor give you 1 day in case you want to make a confirmation.

Promote your paper

Share your work online to make it more visible to others and may increase your chances of citation! Promoting your research is now easy thanks to sharing capabilities on social media websites, where you may already have numerous academic and industry connections, like on Twitter, Facebook, Instagram, LinkedIn, Google Scholar, ORCID, Mendeley, ResearchGate, Publons, or Academia.edu.

Kind Regards, Maria Ulfah Editorial Staff on behalf of Editor-in-Chief Layout and Editing Team International Journal of Public Health Science Institute of Advanced Engineering and Science (IAES) ISSN: 2252-8806 https://ijphs.iaescore.com/index.php/IJPHS/index email: ijphsiaes.core@gmail.com, ijphs@iaescore.com

UNIVERSITAS

widyasari putranti <widyasari@pharm.uad.ac.id>

IJPHS March 2024 Authors' certificate of acknowledgement

1 pesan

ijphsiaes.core@gmail.com <ijphsiaes.core@gmail.com> 28 Februari 2024 pukul 09.39 Kepada: widyasari@pharm.uad.ac.id, desty.restia@gmail.com, nining.sugihartini@pharm.uad.ac.id, tn_saifullah@ugm.ac.id

Dear Author,

Herewith we send a certificate for your contribution as author in IJPHS Journal. We are very thankful to publish your prestige paper, we hope you will publish more in the future and spread the journal among your community.

IJPHS, Vol. 13, No. 1, 2024

We encourage authors to share their published articles on social network and repositories, such as SSRN, arXiv academia.edu, ResearchGate, RePEc, Google Scholar, Linkedin, Facebook, etc.

Certificate Author 22666.pdf

Influence of croscarmellose in fast disintegrating tablet of Syzygium polyanthum extract

Widyasari Putranti¹, Desty Restia Rahmawati¹, Nining Sugihartini¹, Teuku Nanda Saifullah²

¹Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan, Yogyakarta, Indonesia ²Department of Pharmaceutical Technology, Faculty of Pharmacy, Gadjah Mada University, Yogyakarta, Indonesia

Article Info ABSTRACT Article history: Bay leaves (Syzygium polyanthum) contain the flavonoid quercetin which can

Received Nov 1, 2022 Revised Oct 13, 2023 Accepted Oct 25, 2023

Keywords:

Croscarmellose Extra granular Intra granular Superdisintegrant *Syzygium polyanthum* extract Fast disintegrating tablet be used as an antihyperlipidemic drug. The development of antihyperlipidemic drug formula in the form of fast disintegrating tablet (FDT) is needed for patients who experience dysphagia. FDT preparations require an optimal super disintegrant concentration to produce a good drug formula. This study aims to develop the FDT formula of bay leaves extract using the super disintegrant croscarmellose sodium (CCS) intra and extragranular. FDT formulation using the wet granulation method with variations of CCS concentrations; F1: 2%, F2: 3.5%, and F3: 5% for extra granular, and 2% for intra granular. The formulation process, in-process control (IPC) granules, weight uniformity tests, and various physical properties tests of tablets were carried out. Data were statistically analyzed using one way ANOVA test (α =95%). The results of statistical tests of IPC granules, uniformity of weight, and tablet size of all FDT formulas were not significantly different (p>0.05). The CCS concentration for extra granular significantly affected the wetting time, disintegration time, hardness, and the value of friability percentage of FDT (p<0.05). The combination of intra and extra-granular CCS (2%:5%) gave the most optimum physical properties of bay leaf extract FDT.

This is an open access article under the <u>CC BY-SA</u> license.



Corresponding Author:

Widyasari Putranti Department of Pharmaceutical Technology, Faculty of Pharmacy, Universitas Ahmad Dahlan Yogyakarta, Indonesia Email: widyasari@pharm.uad.ac.id

1. INTRODUCTION

Plant germplasm resources are raw materials for medicines, one of which is a bay leaf (*Syzygium polyanthum*). Bay leaves are known to contain secondary metabolites, such as saponins, terpenoids, flavonoids (quercetin), polyphenols, alkaloids, steroids, and essential oils (sesquiterpenes)[1]-[2]. Based on the Decree of the Minister of Health of the Republic of Indonesia (2009), bay leaves contain a total flavonoid of not less than 0.40% which is calculated as quercetin [3]. Quercetin in bay leaves the potential as an antihyperlipidemic which can significantly reduce levels of triglycerides, total cholesterol, and low-density lipoprotein (LDL) cholesterol in plasma and tissues in hyperlipidemic rats with a parallel increase in high-density lipoprotein (HDL), as well as inhibiting 3-hidroksi-3-metilglutaril coenzyme A (HMG-CoA), reductase, and LDL oxidation [4].

Conventional drugs for hyperlipidemia are generally available in tablet form. However, the tablet dosage form has several disadvantages. Some of them require a long absorption time in the body, and slow drug action, and elderly patients may experience difficulty swallowing or dysphagia [5]. Seeing the problems that arise from conventional tablets, the herbal fast disintegrating tablet (FDT) of bay leaf extract is the right alternative so that the drug can be used comfortably and has a therapeutic effect more quickly. FDT is a solid

dosage form containing pharmaceutically active substances that can be disintegrated quickly, generally within seconds when placed on the tongue. This properties of FDT can release the active drug substance immediately, speed up the onset, increase the oral bioavailability of the drug in the body so that therapeutic effectiveness can be achieved [6]-[7]. FDT preparations combine two advantages of preparations, namely liquid which can increase the solubility and bioavailability of drugs, as well as the advantages of solid dosage forms that have high physicochemical stability, relatively constant homogeneity of the active ingredients of the preparation, and easier manufacturing.

FDT is formulated with croscarmellose sodium (CCS) super disintegrant which is classified as a strong swelling product with a good water absorption mechanism [7]-[8]. CCS is effectively used in FDT preparations because it rapidly expands to 4-8 times its initial volume when in contact with water, thus accelerating the tablet-crushing process. CCS is effective intra-granular which allows for better capillary and swelling processes, causing the tablet to disintegrate quickly and reduce disintegration time [8]. The addition of intra and extra-granular crushing materials can increase the effectiveness of tablet disintegration through two stages, i) extra granular which breaks the tablet into granules, and ii) intra granular breaks down the granules into smaller particles. The mechanism is crushing material that experiences rapid swelling and water absorption (wicking) through the gaps between the granules and the particles that make up the granules [9]. This study aims to develop herbal FDT with the active ingredient of bay leaf ethanol extract. The concentration of CCS super disintegrant intra and extra-granular needs to be determined to obtain the FDT of bay leaf extract with optimum physical properties.

2. METHOD

The ingredients used were bay leaf extract in the form of fresh bay leaf collected from tradisional market, Beringharjo market, Yogyakarta, Indonesia, ethanol 70%, croscarmellose sodium (PT. Phapros Tbk), lactose, magnesium stearate, aspartame, menthol and corn starch paste (pharmaceutical grade). The tools for extraction used are oven, rotary evaporator calipers, waterbath, vacuum pump. The tools for granule physical properties test used are Ohaus analytical balance, absorption test equipment granules, volumenometer (tapped density tester), flow tester granule, sieving machine, dry granule sieve, wet granule sieve, pan, halogen moisture analyzer. the tools for tablet test used are single punch tablet printing machine, hardness tester, friability tester, and disintegration tester.

2.1. Bay leaf ethanol extract

Bay leaves were dried in an oven at 50-60°C until a moisture content of less than 10% was obtained. Simplicia were extracted by maceration method using 70% ethanol solvent (1:10) (w/v) [3]. Maserati and pulp were separated by the vacuum pump and filter paper. The solvent was evaporated using a rotary evaporator. The viscous extract obtained was calculated for its yield.

2.2. FDT formulation of bay leaf extract

FDT was made by the wet granulation method. At the granulation stage, bay leaves extract was mixed with intra granular CCS, lactose and corn starch. The mixture was sieved through a wet (no. 18 mesh), ovendried (50-60°C), then sifted dry (no. 20 mesh). The granules and excipients consisting of magnesium stearate, aspartame, menthol, and CCS (extra granular) mixed until homogeneous (30 minutes). Mixture of granules and excipients that have been homogenized are compressed using a single punch tablet press.

2.3. Granule physical properties test parameters

The moisture content of granules can be determined based on the value of loss on drying (LOD) and MC. LOD is a test for measuring the difference in heavy total granules when before and after drying or a moisture content statement based on wet weight. Meanwhile, MC is a statement of moisture content based on dry weight. Measurements using the halogen moisture analyzer tool. Good granule moisture content if the LOD and MC values are 10% [10].

The flow properties of the granules can be tested by calculating the angle of repose, the flow time test, and the determination index (Hausner ratio). The granule angle of repose α (degree) is said to be very good (25°-30°); good (30-35°) or 25°><35°. A good granule flow time is not more than 10 seconds for 100 grams of granules [11]. Excellent granule setting index (\leq 10); good (11–15), and the Hausner ratio index that meets the requirements for the physical properties of the granule mass, which is very good (1.00–1.11); good (1.12–1.18).

Sieve no. 14, 16, 20, 30, 50 mesh and pan, installed in stages on the sieving machine (50 amplitude, 15 minutes). The granules left on each sieve were weighed and the percentage was calculated [10]. The fragility of the granules is based on the number of fines that occur after the friability test by sieving. The granules were put into a graded sieve with the top sieve no. 30 mesh and bottom pan, using a sieving machine (50 amplitude,

30 minutes). The water absorption of the granules affects the tablet's disintegration time (disintegration). It was carried out with a series of absorption test equipment by placing 300 mg of granules on a water-saturated sheath paper. The absorbed water was measured using a digital scale for 15 minutes or until it was constant [7].

2.4. Physical properties test parameters of FDT

Requirements and weight uniformity tests are determined based on the regulation of the National Food and Drug Agency of the Republic of Indonesia concerning the quality requirements of traditional medicines. The test was carried out by weighing 20 tablets one by one, from 20 tablets no more than 2 tablets, each of which deviated from the average weight by more than 5% and not one tablet whose weight deviated from the average is greater than 10% [12].

Each tablet was measured in diameter in a horizontal position with a caliper, and the thickness of the tablet was measured in a vertical position with a caliper (millimeters). The tablet hardness test was carried out by taking a sample of at least 6 tablets from each formula. The tablet hardness test is carried out using hardness tester tablets. One by one the tablets are tested by placing the tablets in the center perpendicular to the hardness tester. A good FDT hardness is 3-5 kg/cm² [11], [13].

For tablets with a single weight of exactly or less than 650 mg, the total weight of the tested samples was close to 6.5 grams. Prior to testing, the tablets were dusted and weighed. All tablets were put into the friability tester (speed 25 rpm, 4 minutes). The tablets were cleaned of adhering fines and re-weighed. A good friability value should not be more than 1% [11].

The wetting time test was done by placing a sheet of filter paper that has been folded twice into a petri dish with a diameter of 5 cm. The petri dish was filled with 5.0 mL of distilled water containing the strawberry red dye. A tablet is then placed on the filter paper, and simultaneously with the start of the test instrument, the stopwatch is turned on. Wetting time was calculated as the time required for a red color to appear on the entire surface of the tablet [14]. A total of 6 tablets of each formula were placed in each tube and a disk was placed on it, the test was carried out with water medium at a temperature of 37 ± 2 °C or 35-39 °C using a 1000 ml glass beaker. The disintegration time requirement for FDT preparations is at least <1 minute disintegration time [11], [15].

2.5. Data analysis

The results of the physical examination of granules and tablets of FDT bay leaf extract were analyzed using the one-way ANOVA test method with a 95% confidence level.

3. RESULTS AND DISCUSSION

Bay leaf simplicia was tested for macroscopic (organoleptic) specifications on fresh and dried simplicia, as well as microscopic tests for simplicia parts and identification fragments of simplicia powder. Fragments prepared by dissolving in chloralhydrate and water [16]. The simplicia specification test was carried out to ensure the correctness of the identity and quality of the bay leaf simplicia based on the Indonesian Herbal Pharmacopoeia compendia and Indonesian Materia Medika. Based on the microscopic test results on Figure 1 there are identifier fragments namely i) lower epidermis with stomata, ii) upper epidermis, iii) elements with dots, iv) prism-shaped calcium oxalate crystals, and v) sclerenchyma. The results of the specification test for the simplicia of bay leaves showed that the simplicia used met the requirements of organoleptic and microscopic terms. The determination test of bay leaf simplicia at the Biology Laboratory of Universitas Ahmad Dahlan (UAD) was carried out to ensure that the selected bay leaf simplicia was in accordance with its species or identity [3], [17].

3.1. Extraction bay leaf

Fresh simplicia of bay leaves were sorted wet, dried using an oven $(50-60^{\circ}C)$ with a moisture content of not more than 10%, then sorted dry. Simplicia powder for extract preparation is a fine simplicial (No. 60 mesh) in order to optimize the extraction process [18]. The results of the simplicial extraction of bay leaves weighing 3.7 kg obtained a thick extract of 347 grams. The total yield of the thick extract obtained was 9.38%. The morphology of the viscous extract is shown in Figure 2. Screening and determination of the chemical compound flavonoid quercetin contained in the thick extract of bay leaves was carried out at the Integrated Research Laboratory of the Faculty of Pharmacy, UAD. According to the Ministry of Health (2017), bay leaves contain a total flavonoid of not less than 0.40% calculated as quercetin. Based on the results of the test the flavonoid content of 0.8079±0.0045%, so that it meets the specifications for the total flavonoid content of bay leaves[17]. Quercetin is an active substance that has medicinal effects such as antiherlipidemia, antibacterial, antidiabetic, antioxidant [18], [19].



Figure 1. Identification fragments of bay leaf; (a) lower epidermis, (b) upper epidermis, (c) floem, (d) calcium oxalate crystal, (e) sclerenchyma



Figure 2. Bay leaf extract

3.2. FDT formulation

FDT formulation method with wet granulation is good used for water-resistant active ingredients and heating temperature [10]. The active ingredient in bay leaf is quercetin (3,3',4',5,7-pentahydroxyflavone) is a flavonoid derivative compound that is resistant to heating with a melting point of 316.5 °C [20], while the wet granulation process ranges from at a temperature of 40-60 °C. The wet granulation method can improve the flowability and compatibility of the material and facilitate the FDT formulation process from the active ingredient in the form of a thick extract [21]. Wet granulation has the advantage of good content distribution and uniformity, besides that the hydrophobic surface of the tablet becomes more hydrophilic which affects the water absorption process into the tablet, thereby increasing the speed of tablet disintegration [22]. The work of woven fibers between granules can increase the speed of tablet disintegration[23], it is synergistic with the presence of CCS super disintegrant extra granular which facilitates swelling and absorption of water into granules and their small particles.

The concentration of CCS super disintegrant intra-granular of the three formulas was the same, 2% (10 mg/tablet), and extra granular was formulated with varying levels, F1 2% (10), F2 3.5% (17.5), and F3 5% (25 mg/tablet). The FDT formula for bay leaf extract is shown in Table 1. Based on the Handbook of pharmaceutical excipients the use of CCS in the wet granulation process should be added in the wet or granulation (intra-granular) stage, as well as in the dry (extra-granular) process [21]. Because this method can increase the wicking ability (water absorption) and swelling of the CCS super disintegrant.[8]

Ingredient	Composition (mg)									
	F1	F2	F3							
Bay leaf extract	70	70	70							
Croscarmellose sodium (intra granular)	10	10	10							
Lactose	380	374	366							
Croscarmellose sodium (extra granular)	10	17.5	25							
Magnesium stearate	5	5	5							
Aspartame	20	20	20							
Menthol	1	1	1							
Corn starch paste 5% w/v	qs	qs	qs							
Total weight (mg)	500	500	500							

Table 1. Formula of FDT bay leaf extract

3.2.1. Granule physical properties test parameters

One of the most crucial elements of wet granulation is the water content, MC and LOD has been was measured and the values of granules from all formulas as shown in Table 2, the results obtained meet the requirements of 10% (p>0.05). Water content will affect granule characteristics, including granule size and compressibility. The flow properties of the granules show the results of the angle of repose of the granules ranging from 30-35 (good), granule flow time of 10-14 grams/second (good), and the determination test/Hausner ratio (very good). The flow properties of granules are very influential on the tablet compression process, granules with good flow properties will cause the granules to enter the die space to be relatively constant so that the tablet weights with small weight variations can be obtained and can increase the uniformity of the resulting tablet dosage [8], [24], [25].

The average diameter of all granules ranged from 650-700 m (p>0.05) and the value of friability percentage of the granules ranged from 0.5-0.9% (p>0.05)[13]. The granule absorption test for all formulas had absorption between 0.120-0.190 mg/minute (p>0.05). The granule absorption shows the speed of the granules in absorbing water to crush the granules into small particles so as to speed up the tablet disintegration time. All in process control (IPC) granule parameters have met the requirements so that the tableting process can be continued with a single punch tablet press with a punch pressure of 13 kg.

Ta	bl	e 2	2.1	R	esul	ts	of	examinat	ion of	pł	iysical	proj	perties	of	FDT	granu	les o	f bay	/ leaf e	extract
----	----	-----	-----	---	------	----	----	----------	--------	----	---------	------	---------	----	-----	-------	-------	-------	----------	---------

Formula							
F1	F2	F3					
7.03±0.76	6.29±1.28	6.57±1.18					
2.98 ± 0.17	2.35±0.33	2.69±0.31					
30.66±1.75	31.18±2.92	30.08±2.30					
12.98 ± 1.28	13.12±0.89	11.85 ± 0.48					
3.17±1.04	2.83±1.04	3.83±0.76					
1.03 ± 0.01	1.03 ± 0.01	1.04 ± 0.01					
694.64±28.87	657.58±39.65	698.14±26.09					
0.8028 ± 0.20	0.8199±0.05	0.5863±0.04					
0.186 ± 0.05	0.188 ± 0.05	0.127±0.01					
	F1 7.03±0.76 2.98±0.17 30.66±1.75 12.98±1.28 3.17±1.04 1.03±0.01 694.64±28.87 0.8028±0.20 0.186±0.05	$\begin{array}{c c} F1 & F2 \\ \hline F2 & F2 \\ \hline 7.03 \pm 0.76 & 6.29 \pm 1.28 \\ 2.98 \pm 0.17 & 2.35 \pm 0.33 \\ 30.66 \pm 1.75 & 31.18 \pm 2.92 \\ 12.98 \pm 1.28 & 13.12 \pm 0.89 \\ 3.17 \pm 1.04 & 2.83 \pm 1.04 \\ 1.03 \pm 0.01 & 1.03 \pm 0.01 \\ 694.64 \pm 28.87 & 657.58 \pm 39.65 \\ 0.8028 \pm 0.20 & 0.8199 \pm 0.05 \\ 0.186 \pm 0.05 & 0.188 \pm 0.05 \\ \end{array}$					

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

3.2.2. Physical properties test parameters of FDT

The uniformity of tablet weight is related to the uniformity of active substance levels and therapeutic effects. The uniformity of tablet weight (Table 3) can be seen from the coefficient of variation F1, F2, and F3 all <5%, (0.0942, 0.1246, and 0.2103) %, respectively. The weight uniformity test based on the National Food and Drug Agency of the Republic of Indonesia requirements, shows the results that meet the requirements, where the tablet meets the specified weight deviation limit value range.

Hardness ranges FDT be good tablet from 3-5 kg/cm² [13], while the conventional tablet, is 4-8 kg/cm². The FDT hardness as shown in Table 3 shows results that meet the requirements. Hardness affects the brittleness and disintegration time of tablets, as well as the dissolution/release of the active drug substance [8]. The results of statistical analysis of tablet hardness with sig. (p<0.05) indicates that the hardness of the three FDT formulas is significantly different. The hardness of FDT tablets is strongly influenced by the CCS super disintegrant excipient. FDT (F3) produced the hardest tablet with the highest extra-granular CCS concentration (5%), causing the particles in the tablet to be tightly bound. Croscarmellose sodium (Ac-Di-Sol) is a super disintegrant with a fibrous cross-linked polymer structure that functions as a disintegrant with a strong swelling mechanism, fiber is able to bind strongly to the particles so as to increase the hardness and reduce the brittleness of the tablet [26]. This causes the tablet to have good compatibility, so there is a tendency
to increase hardness as the amount of CCS in the tablet increases [27]. Another factor that affects the hardness of FDT is the upper punch pressure during the tableting process, the greater the pressure used, the harder the FDT will be. The results of the diameter and thickness measurements of the tablets showed that the size of the entire tablet was uniform (homogeneous).

The friability value of the tablet is based on the requirements, which should not be more than 1% [11]. The fragility value of FDT is shown in Table 3 with a friability value of <1% so that FDT meets the requirements. The results of the tablet friability statistical test have a sig. value (p < 0.05) means that the fragility value between formulas is significantly different. CCS super disintegrant is a factor influencing fragility. Formula 3 with the highest number of CCS has the smallest friability value, where there is a tendency to decrease the friability value as the number of CCS in the tablet increases. The friability value is inversely proportional to the tablet's hardness, where the harder a tablet is, the smaller the friability value will be.

FDT wetting time is shown in Table 3, FDT (F3) with the highest number of CCS has the smallest wetting time. Wetting time is a parameter to determine the speed of FDT in absorbing water, which affects the speed of tablet disintegration. The faster the wetting time, the faster the tablet disintegration. The wetting time of tablets is influenced by the structure of the tablet matrix and the hydrophilicity of the excipients [28]. The result of statistical analysis of FDT wetting time has a sig. value (p<0.05) means that the formulas differ significantly. FDT (F3) with the highest number of CCS had the fastest time for wettable tablets, super disintegrant CCS being the most influential factor. The cross-linked chemical structure of CCS creates excipients that have hydrophilic properties and very easy-to-absorb solvents resulting in outstanding swelling properties in the disintegration process.

Table 3. Results of examination of physical properties of bay leaf extract FDT tablets

Test Parameters	Formula			Note.
	F1	F2	F3	
Weight uniformity±SD (mg)	50 4.42±0.48	50 4.78±0.63	50 4.43±1.06	Appropriate*
Diameter±SD (mm)	12.00±0.00	12.00±0.00	12.00±0.00	Appropriate*
Thickness±SD (mm)	3.67±0.00	3.67±0.00	3.68±0.01	Appropriate*
Hardness±SD (kg)	4.46±0.25°	4.50±0.27°	5.24±0.32 ^{a,b}	In accordance
Fragility±SD (%)	$0.77 \pm 0.05^{\circ}$	$0.72 \pm 0.04^{\circ}$	$0.52\pm0.04^{a,b}$	In accordance
Wetting time±SD (seconds)	50.33±3.06°	43.33±4.04°	20.00±2.65 ^{a,b}	In accordance
Disintegration time±SD (seconds)	45.33±5.86°	38.33±3.79°	19.67±3.06 ^{a,b}	In accordance

Note: (*) sig value. (p>0.05) means that the three formulas are not significantly different

(a) means a formula that is significantly different from the formula 1

(b) means a formula that is significantly different from the formula 2

(c) means a formula that is significantly different from formula 3

F1: CCS intra granular 2%: extra granular 2%

F2: CCS intra granular 2%: extra granular 3.5%

F3: CCS intra granular 2%: extra granular 5%

Disintegration (disintegration) time FDT has at least a disintegration time of less than 1 minute [15]. Another source says that in-vitro about 30 seconds or less [29], whereas according to Ph. euros. is <3 min in the oral cavity before swallowing [29], [30]. The faster disintegration time will increase the speed of drug release from the tablet which affects the effectiveness of therapy. The results of the disintegration time of FDT in table 3 show that it meets the requirements of the disintegration time of FDT (<1 minute). FDT disintegration time test is shown in Table 3 with FDT (F3) having the fastest disintegration time, due to the higher levels of CCS used in F3 than F1 and F2. The results of the statistical test of tablet disintegration time have a sig value (p < 0.05) means that each formula is significantly different. CCS has a dual mechanism, namely water wicking and rapid swelling [8]. The highly porous form of the structure speeds up the disintegration time, because water quickly enters the tablet and increases the tablet wetting rate, through the gaps between the granules and the granule pores [31]. This is related to the factor of the number of CCS super disintegrant excipient used in each formula intra and extra-granular. The use of intra and extra-granular affects the decrease in tablet disintegration time through the swelling mechanism, and along with the increase in the extra-granular CCS concentration, the disintegration time will be faster. The FDT disintegration pattern is also facilitated by intra-granular CCS which causes the tablets to be finely crushed not in the form of granules but into small particles so that the speed of disintegration is optimal. The optimum formula for FDT of bay leaf extract is shown in Table 4 with the results of all evaluations of the physical properties of the granules and FDT (F3) that meet the requirements and show the optimum results of physical parameters.

The results of this study are in line with the research of Puri *et al.* [24], that the combination of intragranular and extra-granular can produce tablets with the most effective disintegration time by providing better tablet physical properties, compared to intra-granular or extra-granular only. Extra-crushing material granules have a high tendency to absorb water from the surrounding liquid through the tablet-breaking mechanism [32]. Tablets with disintegrant intra-granular have a higher hardness than tablets with extra granular crushing agents. By combining intra and extra granular super disintegrants, a good level of tablet hardness can be obtained so that the tablet is not brittle and has a faster disintegration time with the mechanism of tablet disintegration into granules (extra granular factor) and granules into small particles (intra granular factor).

Table 4. various parameters of optimized tablet formula (F3)			
No.	Test parameters	Optimized formula value	
1.	Weight uniformity±SD (mg)	504.43±1.06	
2.	Diameter±SD (mm)	12.00±0.00	
3.	Thickness±SD (mm)	3.68±0.01	
4.	Violence±SD (kg)	5.24±0.32	
5.	Fragility±SD (%)	0.52±0.04	
6.	Wetting time±SD (seconds)	20.00 ±2.65	
7.	Disintegration time±SD	19.67±3.06	
	(seconds)		

6 ... 1.11.6

4. CONCLUSION

TT 1 1 4 TZ ·

In the FDT research, to improve the poor flow properties of the extract, the wet granulation method was carried out. The IPC results for the three formulas showed granules that met the requirements. This research provides information that the use of extra granular CCS at higher levels will increase the rate of disintegration. The combination of CCS intra granular 10 mg (2%) and extra granular 25 mg (5%) can produce FDT with the most effective. By combining intra and extra granular super disintegration time. In addition, FDT bay leaves extract can be used as an alternative treatment for hyperlipidemia in patients who experience dysphagia. This research still needs development to determine the stability of FDT bay leaves extract where the active substance is herbal medicine.

ACKNOWLEDGEMENTS

This research was carried out with the assistance of the Higher Education Research Grant No PKPT-013/SKPP/III/2018.

REFERENCES

- L. Hartanti, S. M. K. Yonas, J. J. Mustamu, S. Wijaya, H. K. Setiawan, and L. Soegianto, "Influence of extraction methods of bay leaves (Syzygium polyanthum) on antioxidant and HMG-CoA Reductase inhibitory activity," *Heliyon*, vol. 5, no. 4, p. e01485, Apr. 2019, doi: 10.1016/j.heliyon.2019.e01485.
- [2] E. Kato, R. Nakagomi, M. D. P. T. Gunawan-puteri, and J. Kawabata, "Identification of hydroxychavicol and its dimers, the lipase inhibitors contained in the Indonesian spice, Eugenia polyantha," *Food Chemistry*, vol. 136, no. 3–4, pp. 1239–1242, 2013, doi: 10.1016/j.foodchem.2012.09.013.
- [3] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia, 1st ed. Jakarta Indonesia, 2009.
- [4] K. Zeka, K. Ruparelia, R. Arroo, R. Budriesi, and M. Micucci, "Flavonoids and their metabolites: prevention in cardiovascular diseases and diabetes," *Diseases*, vol. 5, no. 3, p. 19, 2017, doi: 10.3390/diseases5030019.
- [5] I. Safitri, S. Sulistiyaningsih, and A. Y. Chaerunisaa, "Review: superdisintegrants in oral preparations," (in Indonesia), *Farmasetika.com* (Online), vol. 4, no. 3, p. 56, 2019, doi: 10.24198/farmasetika.v4i3.22945.
- [6] A. Masih, A. Kumar, S. Singh, and A. K. Tiwari, "Fast Dissolving Tablets: A Review," *International Journal of Current Pharmaceutical Research*, vol. 9, no. 2, p. 8, Mar. 2017, doi: 10.22159/ijcpr.2017v9i2.17382.
 [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam leaves
- [7] W. Putranti, C. A. Edityaningrum, E. Prastyaningrum, and L. Widiyastuti, "Formulation of fast disintegrating tablet of salam leaves ethanol extract with combination of crospovidone and croscarmellose sodium as superdisintegrants," *Jurnal Sains Farmasi & Klinis*, vol. 8, no. 3, p. 285, 2021, doi: 10.25077/jsfk.8.3.285-295.2021.
- [8] M. E. Rowe, R. C., Sheskey, P. J. dan Quinn, Handbook of Pharmaceutical Excipients, Sixth. London: Pharmaceutical press., 2009.
- [9] M. O. Okekunle, O. D. Akin-Ajani, B. L. Ekpukpon, O. A. Odeku, B. I. Olu-Owolabi, and K. O. Adebowale, "Intra and Extragranular Disintegrant Properties of Modified Underutilised Red Lima Bean Starch in Paracetamol Tablet Formulation," *Brazilian Journal of Pharmaceutical Sciences*, vol. 58, 2022, doi: 10.1590/s2175-97902022e19967.
- [10] L. Lachman and D. Ph, The theory and practice of industrial pharmacy: Tablets, Gilbert S. Banker, Neil R. Anderson, 3rd ed. Garden City, New York; Livingston, New Jersey; Ridgefield, Connecticut, 1986.
- [11] M. 20852 The United States Pharmacopeial Convention 12601 Twinbrook Parkway, Rockville, Ed., The United States Pharmacopeia, USP 41/The National Formulary, NF 36. Baltimore, MD. Rockville.: United Book Press, Inc., 2018.
- [12] Food and Drug Supervisory Agency of the Republic of Indonesia, No Title 1200, 2014, BPOM. Obat Tradisional. Mutu. Persyaratan., no. 1200. 2014.
- [13] R. Rahane and P. R. Rachh, "A review on fast dissolving tablet," *Journal of Drug Delivery and Therapeutics*, vol. 8, no. 5, pp. 50– 55, Sep. 2018, doi: 10.22270/jddt.v8i5.1888.
- [14] C. P. Jain and P. S. Naruka, "Formulation and evaluation of fast dissolving tablets of Doxazosin Mesylate," Asian Journal of Pharmaceutical Research, vol. 6, no. 3, p. 131, 2016, doi: 10.5958/2231-5691.2016.00020.4.
- [15] Allen, L.V., N. G. Popovisch, and H. C. Ansel, Ansel's Dossage Forms and Drug Delivery Systems, 9th Ed., no. 1. 2011. doi: 10.4018/jsds.2010103001.

- [16] T. S. Wahyuni, N. Khoiriyah, L. Tumewu, W. Ekasari, A. Fuad, and A. Widyawaruyanti, "Microscopic and physicochemical evaluation of Ruta angustifolia leaves," *Journal of Public Health in Africa*, vol. 14, no. S1, pp. 123–129, 2023, doi: 10.4081/jphia.2023.2520.
- [17] Ministry of Health of the Republic of Indonesia, Farmakope Herbal Indonesia (FHI), II. Jakarta: k, 2017.
- [18] E. Sutrisna, Y. Nuswantoro, and R. F. Said, "Hypolipidemic of ethanolic extract of Salam bark (Syzygium polyanthum (Wight) Walp.) from Indonesia (Preclinical study)," *Drug Invention Today*, vol. 10, no. 1, pp. 55–58, 2018.
- [19] M. A. Syabana, N. D. Yuliana, I. Batubara, and D. Fardiaz, "α-glucosidase inhibitors from Syzygium polyanthum (Wight) Walp leaves as revealed by metabolomics and in silico approaches," *Journal of Ethnopharmacology*, vol. 282, p. 114618, Jan. 2022, doi: 10.1016/j.jep.2021.114618.
- [20] H. Maurya and T. Kumar, "Formulation, standardization, and evaluation of polyherbal dispersible tablet," *International Journal of Applied Pharmaceutics*, vol. 11, no. 1, pp. 158–167, 2019, doi: 10.22159/ijap.2019v11i1.30113.
- [21] P. Sirame, R. Khairnar, and R. Tasgaonkar, "Granulation," *International Journal for Research in Applied Science and Engineering Technology*, vol. 11, no. 1, pp. 347–355, Jan. 2023, doi: 10.22214/ijraset.2023.48564.
- [22] C. M. Shinko, "Granulation Characterization: Methods and Significance, Dalam: Parikh, D.M.(ed.), Handbook of Pharmaceutical Granulation Technology, Marcel Dekker, New York," p. 422, 1997.
- [23] J. Swarbrick, J. C. Boylan, and 2002, Encyclopaedia of Pharmaceutical Technology, 2nd ed., vol. 3, no. 4. New York: Organic Process Research * Development 2003, 613, 2003.
- [24] D. Puri *et al.*, "Formulation of herbal fast disintegrating tablets and its ex-vivo study for anti-histaminic activity in guinea pig ileum," *Current Clinical Pharmacology*, vol. 13, no. 2, pp. 128–135, 2018, doi: 10.2174/1574884713666180309153942.
- [25] E. Schlick-Hasper, J. Bethke, N. Vogler, and T. Goedecke, "Flow properties of powdery or granular filling substances of dangerous goods packagings—Comparison of the measurement of the angle of repose and the determination of the Hausner ratio," *Packaging Technology and Science*, vol. 35, no. 10, pp. 765–782, Oct. 2022, doi: 10.1002/pts.2678.
- [26] J. Balasubrmaniam, Balasubrmaniam, J. 2009, Influence of Superdisintegrants on The Rate of Drug Dissolution from Oral Solid Dosage Forms, Pharmaceutical Technology, April., April. Pharmaceutical Technology, 2009.
- [27] A. N. Bestari, T. N. S. Sulaiman, and A. Rohman, "Formulasi orally disintegration tablet (Odt)," *Majalah Farmaseutik*, vol. 12, no. 2, pp. 453–465, 2016, doi: 10.22146/farmaseutik.v12i2.26454.
- [28] C. G. Jange, C. R. Wassgren, and K. Ambrose, "The Significance of tablet internal structure on disintegration and dissolution of immediate-release formulas: a review," *Powders*, vol. 2, no. 1, pp. 99–123, Feb. 2023, doi: 10.3390/powders2010008.
- [29] W. Brniak, R. Jachowicz, and P. Pelka, "The practical approach to the evaluation of methods used to determine the disintegration time of orally disintegrating tablets (ODTs)," *Saudi Pharmaceutical Journal*, vol. 23, no. 4, pp. 437–443, 2015, doi: 10.1016/j.jsps.2015.01.015.
- [30] S. Kakar, R. Singh, and S. Kumar, "Orodispersible tablets: an overview," MOJ Proteomics & Bioinformatics, vol. 7, no. 3, pp. 180– 182, 2018, doi: 10.15406/mojpb.2018.07.00230.
- [31] P. Dey and S. Maiti, "Orodispersible tablets: A new trend in drug delivery," *Journal of Natural Science, Biology and Medicine*, vol. 1, no. 1, p. 2, 2010, doi: 10.4103/0976-9668.71663.
- [32] S. U. Pradhan et al., "Characteristics of multi-component formulation granules formed using distributive mixing elements in twin screw granulation," Drug Development and Industrial Pharmacy, vol. 44, no. 11, pp. 1826–1837, Nov. 2018, doi: 10.1080/03639045.2018.1503293.

BIOGRAPHIES OF AUTHORS



Widyasari Putranti (b) S (s) (c) is lecturer in department of technology pharmaceutical, Universitas Ahmad Dahlan, Yogyakarta Indonesia. Her research focuses on drug delivery system, pre-formulation, physical pharmacy, fast disintegrating tablet and herbal formulation. She can be contacted at email: widyasari@pharm.uad.ac.id.



Desty Restia Rahmawati D S S v was bachelor of faculty of pharmacy, Universitas Ahmad Dahlan, Yogyakarta Indonesia. She is a doctoral Student in Pharmaceutical Sciences Universitas Gadjah Mada. She achieved Best oral Presentation in the 4th International Conference on Pharmacy and Pharmaceutical Science (ICPPS) in Global Front, Surugadai Campus, Meiji University, Tokyo, Japan. She can be contacted at email: desty.restia@gmail.com.



Nining Sugihartini (D) 🔀 🔄 C is lecturer in department of technology pharmaceutical Universitas Ahmad Dahlan, Yogyakarta Indonesia. Her research interest in drug and cosmetic development. Currently working on optimization of green tea extract topical dosage formulas as anti-inflammatory. She can be contacted at email: nining.sugihartini@pharm.uad.ac.id.



Teuku Nanda Saifullah (D) (S) (C) is Lecturer in Department of Technology Pharmaceutical Universitas Gadjah Mada, Yogyakarta Indonesia. Medical And Health Sciences/Pharmaceutical Technology, Pharmaceutics Department. His research interest are drug delivery system, technology and formulation of pharmaceutical products. He can be contacted at email: tn_saifullah@ugm.ac.id.