

# Hasil\_ Ichsan Luqmana

*by* Ichsan Luqmana

---

**Submission date:** 26-Jan-2023 01:36PM (UTC+0700)

**Submission ID:** 1999695396

**File name:** 45-IRJIS-7068.doc (230.5K)

**Word count:** 3871

**Character count:** 21703

INSECT PREDATOR OF *SPODOPTERA FRUGIPERDA* (J. E. SMITH, 1797) IN SLEMAN AND GUNUNGKIDUL REGENCY, INDONESIA

Ichsan Luqmana  
Indra Putra<sup>1</sup>  
Novia Saphira  
Prakusya<sup>2</sup>  
Defi Yuliasari<sup>2</sup>

<sup>1</sup>Ecology and Systematics Research Laboratory, Biology Study Program, Faculty of Applied Science and Technology, Ahmad Dahlan University, Indonesia. (+ Corresponding author)

<sup>2</sup>Biology Study Program, Faculty of Applied Science and Technology, Ahmad Dahlan University, Indonesia.

<sup>3</sup>Jl. Ahmad Yani, Kragilan, Tamanan, Bantul Regency, Yogyakarta Province, Indonesia 55191.

\*Corresponding author: [ichsan.luqmana@bio.uad.ac.id](mailto:ichsan.luqmana@bio.uad.ac.id)

ABSTRACT

Article History

Received: xxxxxxxx

Revised: xxxxxxxx

Accepted: xxxxxxxx

Published: xxxxxxxx

Keywords

Abundance

Armyworm

*Menochilus sexmaculatus*

Predator

*Spodoptera frugiperda*

*Sycanus* sp.

This research was conducted to identify and found the abundance of predatory insect species of *S. frugiperda* in Sleman and Gunungkidul Regencies, Yogyakarta Province, Indonesia. This research was conducted in June 2021 - March 2022 on corn plantations in Sleman and Gunungkidul Regencies. Determination of the sampling location was determined by stratified random sampling. Corn land used in sampling was determined by purposive. Insect predator caught by using sweeping net and hand picking. The results showed that there are 3 orders, 8 families, and 16 species of insect predators. 3 orders of the insect predators that found in the Sleman and Gunungkidul Regencies were Coleoptera, Hymenoptera, and Hemiptera. Insect predator families that found in Sleman dan Gunungkidul Regencies were Coccinellidae, Carabidae, Staphylinidae, Formicidae, Vespidae, Pentatomidae and Reduviidae. Insect predators species that was most commonly found was *Menochilus sexmaculatus* with 467 individuals and the least was *Sycanus* sp. with 1 individual. The conclusion of this study is that there were 16 species of predators from *S. frugiperda* in Sleman and Gunungkidul Regencies. Practical implications of this research were the insect predators that found in Sleman and Gunungkidul Regencies can be used to maintain the population of *Spodoptera frugiperda* in two regencies.

**Contribution/Originality:** This research was the first research that study the species predatory insect of *Spodoptera frugiperda* in Indonesia, especially in D.I. Yogyakarta, specifically in Sleman and Gunungkidul Regencies.

1. INTRODUCTION

The corn armyworm (*Spodoptera frugiperda* JE Smith, 1797) is an insect native to the United States that has spread to the Asian continent, including Indonesia. *Spodoptera frugiperda* was originally found in the Pasaman area, West Sumatra [1] and has spread to Bandung, Garut, Sumedang [2] to Sleman, Gunungkidul, Bantul, Yogyakarta [3]. Armyworm caterpillars have several host plants, especially from the Graminae group such as corn, rice [4] cotton [5] and sorghum [6]. *Spodoptera frugiperda* had causes losses to corn crops of 8.3 - 20.6 million tons per year with an economic loss value of US \$ 2.5 - 6.2 billion per year on the African Continent and Europe.

Control of these caterpillars has been carried out to minimize yield losses from corn plants. So far, armyworms have been controlled using synthetic pesticides [7, 8]. Originally there were natural enemies that have been found to be able to control this pest population in nature [9]. Natural enemies found in nature can disrupt species diversity and populations due to the use of insecticides [10]. One of the natural enemies of *S. frugiperda* found in

nature is from the predator class [11]. Several armyworm predators that have been found include *Lycosa* sp., and *Coccinella transversalis* Thunberg [9]. In addition, research in West Lombok conducted by Jannah, et al. [12] found 8 species of predatory insect of *S. frugiperda*, including *Cheilomenes sexmaculata* (Fabr., 1781), *Coelophora inaequalis* (Fabr., 1775), *Verania lineata* (Thunberg, 1781), *Polistes gallicus* (Linnaeus, 1767), *Polyrhachis dives* (Smith, 1857), *Chrysopa* sp., and *Lycosa* sp.

## 2. MATERIAL AND METHODS

This research was conducted in June 2021 - May 2022 on corn (*Z. mays*) plantations in 17 sub-districts of Sleman and 18 sub-district of Gunungkidul Regency. Identification of predators was carried out at the Ecology and Systematics Research Laboratory, Faculty of Applied Science and Technology, Ahmad Dahlan University. The materials used in this study were samples of predatory insects from the field and 70% alcohol.

The methods that used in this research were:

### 2.1. Determination of the Sampling Location

Determination of the location of this sample is determined by a random stratified method. Samples were taken from 17 sub-districts in Sleman and 18 sub-districts in Gunungkidul Regency. Two villages were taken from each sub-district to be used as sampling locations using a random stratified method. Each village is taken one land used in predator sampling. The maize fields used in predator sampling in each village were determined using the purposive. The age of the corn plants used in this study was around 2-3 weeks old and had not yet flowered.

### 2.2. Sampling and Preservation

Sampling was carried out every two weeks, when sampling there is no repetition in the same land (one field for one sampling), this sampling is carried out by exploring corn plantations in Sleman and Gunungkidul Regencies by determining in advance which village and sub-district to go to by looking at maps and then tracing the streets around the village and sub-district. Samples were taken in the form of insect predators of *S. frugiperda* which were carried out mechanically using hands, sweep nets and tweezers. Sampling was carried out by exploring all corn plants affected by *S. frugiperda* with perforated or damaged leaves. The insect predator that obtained were put into a 50 ml plastic bottle filled with 70% alcohol to preserve the sample.

### 2.3. Identification of the Insect Predator

Predator sample which had been preserved in 70% alcohol then observed for its morphological characteristics. Identification was carried out by comparing morphological characters including the shape of the mouthparts, body color, number of *tarsus* the limbs, and patterns on the wings of the predators which were obtained from books and journals Manting, et al. [13]; Ndiaye, et al. [14]; Fallahzadeh, et al. [15]; Rodríguez-Vélez, et al. [16] and Mayadunnage, et al. [17]. The species of predators that are know were then documented using a cellphone and the number of individuals was counted to find out which predators have the most and the least number of individuals.

Table 1. Types of predators of *S. frugiperda* in Sleman and Gunungkidul Regencies.

Ordo	Families	Species
Coleoptera	Coccinellidae	<i>Coccinella transversalis</i>
		<i>Verania lineata</i>
		<i>Menochilus sexmaculatus</i>
		<i>Exochomus nigripennis</i>
		<i>Paederus fuscipes</i>
	Staphylinidae	<i>Ophionea nigrifasciata</i>
Hymenoptera	Formicidae	<i>Polyrhachis dives</i>
		<i>Diacamma rugosum</i>
		<i>Odontomachus chelifer</i>

		Dolichoderus thoracicus
		Paratrechina longicornis
	Vespidae	Vespa sp.
		Polistes sp.
Hemiptera	Pentatomidae	Andralus spinidens
	Reduviidae	Sycanus sp.

12

### 3. RESULT AND DISCUSSION

#### 3.1. Insect Predator of *S. frugiperda* in Sleman and Gunungkidul Regency

This study found 3 orders, 8 families, and 16 species of predators from *S. frugiperda*. Families with species that act as predators of *S. frugiperda* were most commonly found in the Formicidae (5 species) and Coccinellidae (4 species) families. While the families that act as predators at least are Staphylinidae, Carabidae, Reduviidae and Pentatomidae (1 species) Table 1.

The most common species found as predators of *S. frugiperda* in this study came from the Formicidae Family. The Formicidae family is an insect belonging to the ant group (Order Hymenoptera). Ants are insects that are widely distributed throughout the world except in the polar regions [14]. The Formicidae family consists of 14 sub-families and based on their role in nature, 29% of ants act as predators. Predator ants are active and strong and prey on smaller insects [15]. Many members of this family are found as predators of *S. frugiperda* due to the wide range of range of its family members. In addition, ants have pheromone fluids that make it easier for their members to find prey in hidden locations [16].

Apart from the Formicidae family, many members of the Coccinellidae family were also found as predators of *S. frugiperda* in this study. Many members of the Coccinellidae family were found because around the sampling location there were plants other than corn, such as chili and rice. Both of these plants can affect the distribution of members of the Coccinellidae family [17] because these plants contain aphids. These aphids are the main prey of Coccinellidae [18].

The presence of aphids on plants around the corn field will make members of the Coccinellidae family come to prey on these aphids. The more aphids found on plants around the corn field, the more species and abundance of Coccinellidae members found in that location [19]. This is in accordance with the opinion of Rain, et al. [20] that the diversity and abundance of predatory Coccinellidae depends on the presence or absence of aphids at that location. Apart from surrounding plants, aphids can also be found on corn plants [21, 22].

The presence of aphids on corn plants can become an alternative prey if the population of aphids found on surrounding plants has decreased [18]. The advantage of Coccinellidae members who prey on aphids on corn plants, besides getting prey in the form of aphids, they can also prey on the first and second instar larvae of *S. frugiperda* found on these corn plants. This is in accordance with research from Nurkoro, et al. [3] who stated that apart from preying on aphids, predatory Coccinellidae found on corn plants also prey on the early instar larvae of *S. frugiperda*. This opinion is in line with the statement of Malo and Hore [23] which states that the Coccinellidae family is a predator of the eggs and young instar larvae of *S. frugiperda*.

Table 2. The number of individual predators of *S. frugiperda* found on corn plants in Sleman and Gunungkidul Regencies

Species	Number of individuals
<i>Coccinella transversalis</i>	87
<i>Verania lineata</i>	301
<i>Menochilus sexmaculatus</i>	467
<i>Exochomus nigripennis</i>	32
<i>Paederus fuscipes</i>	50
<i>Ophionea nigrifasciata</i>	2
<i>Polyrachis dives</i>	186
<i>Diacamma rugosum</i>	14

<i>Odontomachus chelifer</i>	16
<i>Dolichoderus thoracicus</i>	3
<i>Paratrechina longicornis</i>	5
<i>Vespa</i> sp.	9
<i>Polistes</i> sp.	21
<i>Andralus spinidens</i>	10
<i>Sycanus</i> sp.	1

### 3.2. Abundance of Insect Predator of *S. frugiperda* in Sleman and Gunungkidul Regencies

The results obtained in this study indicate that the predators of *S. frugiperda* that were found mostly on corn, namely *Menochilus sexmaculatus* with a total of 467 individuals and the next is *Verania lineata* with 301 individuals. Meanwhile, the fewest predators of *S. frugiperda* found on corn, namely *Sycanus* sp. with only one individual Table 2.

*Menochilus sexmaculatus* was most commonly found as a predator of *S. frugiperda* in this study because this beetle has a wide range of prey [17]. Apart from aphids, *M. sexmaculatus* is also known to prey on the whitefly *Bemisia tabaci* (Gennadius, 1889) [24] and *Diaphorina citri* (Kuwayama, 1908) [25]. Among the prey that has been mentioned, *M. sexmaculatus* has a preference for *Aphis gossypii*. In addition, the large number of chili plants found around the study site allows for an abundance of individuals of *A. gossypii* as prey for *M. sexmaculatus*. The more prey available, the higher the abundance of predators following the abundance of their prey [26]. *Menochilus sexmaculatus* will certainly prefer prey populations that are always available compared to prey that are only available at certain times [26].

Apart from having a wide range of prey, another factor that causes the high population of *M. sexmaculatus* is because it's a voracious predator [17]. The first to fourth instar larvae of *M. sexmaculatus* were able to prey on 347 *A. gossypii* within 5 days and one adult of *M. sexmaculatus* was able to prey on 270 prey per day with the same type of prey. The predatory insect species of *S. frugiperda* that had the second highest number of individuals in this study was *V. lineata*. The high population of this beetle is because *V. lineata* does not only prey on *S. frugiperda* larvae. *Verania lineata* also known to prey on the brown planthopper (*Nilaparvata lugens*) [27]. These planthoppers are commonly found in rice plants [28]. However, the rice plants around the study site entered the harvest season, so the population of *N. lugens* was not abundant. This is why *V. lineata* also switches to other prey if its main prey is not found. *Verania lineata* also preys on aphids, thrips, and whitefly. Aphids preyed upon by *V. lineata* are *A. gossypii* and *A. craccivora* that in legumes plant. The legume plants found around the research site are not as abundant as chili plants. This is why the population of *V. lineata* is not as large as *M. sexmaculatus*.

Apart from being influenced by other prey preferences, the Coccinellidae family can be found in abundance because they are able to produce a large number of eggs. An adult female of *M. sexmaculatus* can produce as many as 100-200 eggs during her life cycle, while an adult female of female *V. lineata* is able to produce as many as 103 eggs during her life cycle. Meanwhile, for the other two Coccinellidae species, the number of individuals was not abundant because an adult female of *C. transversalis* was only able to lay 100 eggs during its life cycle, and an adult female of *E. nigripennis* was able to lay 69 eggs during its life cycle. These results also support that the ability of female imago to produce eggs is the highest for *M. sexmaculatus* and *V. lineata*, so that in this study both types had the most abundant number of individuals compared to other Coccinellidae species. In addition to finding predators from *S. frugiperda* with the highest abundance of individuals, this study also found predators from *S. frugiperda* with the least abundance of individuals.

Apart from being influenced by food factors and the ability to reproduce, the abundance of predators found was also influenced by abiotic factors. The abiotic factors that influenced the number of individual insect predators of *S. frugiperda* in this study included air temperature and humidity. The air temperature and humidity measured in this study had an average air temperature range of 21.8 – 31.150C and an air humidity range of 52 – 87.78% (Table 3). *Menochilus sexmaculatus* has a minimum temperature range of 200C, optimum temperature of 250C, and maximum temperature of 300C [29] whereas the humidity range of 66 – 85% [30]. So the temperature and humidity at the

study site can support the survival of *M. sexmaculatus*. This is why the individual abundance of *M. sexmaculatus* was found to be high in this study. In contrast to *M. sexmaculatus*, *V. lineata* was also found in the average air temperature range of 21.8 – 31.150C and humidity range of 52 – 87.78%. According to Syahrawati, et al. [27] *V. lineata* can be found in the temperature range of 26.7 – 29.20C and humidity of 70.6 – 80.9% so that the temperature and humidity in the study site are able to support the survival of *V. lineata*.

**Table 3.** Mean abiotic factors measured at sampling locations in Sleman and Gunungkidul Regencies, Indonesia.

Regency	Sub-district	Wind velocity (m/s)	Light intensity (lux)	Air temperature (oC)	Air humidity (%)
Sleman	Berbah	0	9707.5	27.05	87.85
	Kalasan	1.2	66234	28.5	53.5
	Ngemplak	0.65	1310	24.85	58.7
	Prambanan	0	4450	23.15	70.3
	Ngaglik	0.4	7402.5	25.05	61.65
	Pakem	0.2	15588	25.6	63
	Cangkringan	0.7	2290	26.45	52
	Sleman	0.2	5839.5	25.5	70.85
	Turi	0.2	3199	25.05	77
	Mlati	0.2	2079.3	27	63.1
	Tempel	0.9	19327	29.2	67.3
	Depok	0.65	37578.5	30.3	53.15
	Moyudan	0.2	6857	21.8	75.65
	Minggir	0.4	21780	27.2	74
	Seyegan	0.4	23292.5	29.05	61.8
Godean	1.35	11163.5	31.15	52.65	
Gamping	0.05	5108	29.1	53.2	
Gunungkidul	Patuk	0.1	86595.5	30.51	62.67
	Playen	0.12	16879.17	28.27	73
	Paliyan	0.03	52630	29.16	81.34
	Panggung	0.08	9531.83	33.05	63.67
	Gedangsari	0.52	133333.33	31.55	64.67
	Wonosari	0.39	401696.67	32.08	63.56
	Saptosari	0.13	30951.83	27.55	70
	Nglipar	0.03	2033	30.77	64.67
	Semin	0.2	12937.17	26.29	72.67
	Ngawen	0.93	57865	29.56	67.5
	Karangmojo	0.62	46592.11	29.46	66.56
	Semanu	1.23	14831.67	28.40	55.84
	Tepus	0.9	87195	30.18	54
	Tanjungsari	1.71	80916.67	32.05	54.34
	Purwosari	0.62	68396.63	30.36	50
	Ponjong	1.32	4204.5	24.92	94.34
	Girisubo	0.02	11270.33	26.15	91.67
Rongkop	0.12	182316.67	33.87	51.5	

#### 4. CONCLUSION

The insects predator species of *S. frugiperda* found in this research consist of 16 species divided into 3 orders. The predatory type of *S. frugiperda* with the highest number of individuals was *Menochilus sexmaculatus*. While the least number of individuals is *Sycanus* sp.

**Funding:** This study received no specific financial support  
**Competing Interests:** The authors declare that they have no competing interests.  
**Acknowledgement:** The authors are grateful to the Dean of Faculty of the Applied Science and Technology, Ahmad Dahlan University for the opportunity to do this research.

#### REFERENCES

- [1] S. P. Sari, I. Suliansyah, N. Nelly, and H. Hamid. (2021). The occurrence of spodoptera frugiperda attack on maize in West Pasaman District, West Sumatra, Indonesia. Paper presented at the IOP Conference Series: Earth and Environmental Science. IOP Publishing.
- [2] Y. Maharani, V. K. Dewi, L. T. Puspasari, L. Rizkie, Y. Hidayat, and D. Dono, "Cases of fall army worm spodoptera frugiperda JE Smith (Lepidoptera: noctuidae) attack on maize in Bandung, Garut and Sumedang District, West Java," Cropsaver-Journal of Plant Protection, vol. 2, no. 1, pp. 38-46, 2019. <https://doi.org/10.24198/cropsaver.v2i1.23013>
- [3] I. Nurkomar, I. L. I. Putra, D. W. Trisnawati, M. Saman, R. G. Pangestu, and A. Triyono. (2021). The existence and population dynamic of new fall armyworm species spodoptera frugiperda JE Smith (Lepidoptera: Noctuidae) in Yogyakarta, Indonesia. Paper presented at the IOP Conference Series: Earth and Environmental Science. IOP Publishing.
- [4] D. G. Montezano *et al.*, "Host plants of spodoptera frugiperda (Lepidoptera: noctuidae) in the Americas," African Entomology, vol. 26, no. 2, pp. 286-300, 2018. <https://doi.org/10.4001/003.026.0286>
- [5] D. Koffi *et al.*, "Natural enemies of the fall armyworm, spodoptera frugiperda (JE Smith)(Lepidoptera: Noctuidae) in Ghana," Florida Entomologist, vol. 103, no. 1, pp. 85-90, 2020. <https://doi.org/10.1653/024.103.0414>
- [6] R. Baker *et al.*, "Spodoptera frugiperda pest report and datasheet to support ranking of EU candidate priority pests," EFSA Journal, vol. 17, pp. 1-49, 2019. <https://doi.org/10.5281/zenodo.2789778>
- [7] B. Sisay, T. Tefera, M. Wakgari, G. Ayalew, and E. Mendesil, "The efficacy of selected synthetic insecticides and botanicals against fall armyworm, Spodoptera frugiperda, in maize," Insects, vol. 10, no. 2, p. 45, 2019. <https://doi.org/10.3390/insects10020045>
- [8] I. Cruz, M. D. L. C. Figueiredo, R. B. Da Silva, and J. E. Foster, "Efficiency of chemical pesticides to control Spodoptera frugiperda and validation of pheromone trap as a pest management tool in maize crop," Brazilian Magazine of Corn and Sorghum, vol. 9, no. 2, pp. 107-122, 2010. <https://doi.org/10.18512/1980-6477/rbms.v9n2p107-122>
- [9] D. M. Firake and G. T. Behere, "Natural mortality of invasive fall armyworm, Spodoptera frugiperda (JE Smith)(Lepidoptera: Noctuidae) in maize agroecosystems of northeast India," Biological Control, vol. 148, p. 104303, 2020. <https://doi.org/10.1016/j.biocontrol.2020.104303>
- [10] A. De Freitas Bueno, G. A. Carvalho, A. C. dos Santos, D. R. Sosa-Gómez, and D. M. da Silva, "Pesticide selectivity to natural enemies: Challenges and constraints for research and field recommendation," Rural Science, vol. 47, no. 6, pp. 1-10, 2017. <https://doi.org/10.1590/0103-8478cr20160829>
- [11] S. Lamsal, S. Sibi, and S. Yadav, "Fall armyworm in South Asia: Threats and management," Asian Journal of Advances in Agricultural Research, vol. 13, pp. 21-34, 2020. <https://doi.org/10.9734/ajaar/2020/v13i330106>
- [12] M. Jannah, S. Bambang, and W. Mery. (2021). Diversity of fall armyworm (Spodoptera frugiperda) predator during the growth of corn plant (Zea mays L) in Ireng Village, West Lombok. Paper presented at the 45th Celebration National Seminar of UNS.
- [13] M. M. E. Manting, M. A. J. Torres, and C. G. Demayo, "Describing variability in mandibles shapes in selected workers of the ant diacamma rugosum Le guillou (Hymenoptera: Formicidae)," International Research Journal of Biological Science, vol. 2, pp. 8-15, 2013.
- [14] A. Ndiaye, M. Faye, I. Ba, I. Diallo, and P. M. Sembene, "The fall armyworm spodoptera frugiperda J. E. Smith, a new pest of maize in Africa: Monitoring, damage evaluation and identification of natural enemies on production areas of Senegal," International Journal of Biological and Chemical Sciences vol. 15, pp. 2247-2260, 2021. <https://doi.org/10.4314/ijbcs.v15i6.1>
- [15] M. Fallahzadeh, R. Abdmaleki, and N. Saghaei, "Contribution to the knowledge of the ladybird beetles (Coleoptera, Coccinellidae), predators of mealybugs (Hemiptera, Pseudococcidae) in Hormozgan province, Southern Iran," Linz Biological Contributions, vol. 45, no. 1, pp. 673-679, 2013.
- [16] J. M. Rodríguez-Vélez *et al.*, "Identification of twelve species of coccinellidae (Coleoptera) predatory on Melanaphis sacchari (Zehntner)(Hemiptera: Aphididae) in Mexico, and submission of reference COI sequences," The Coleopterists Bulletin, vol. 73, no. 1, pp. 243-251, 2019. <https://doi.org/10.1649/0010-065x-73.1.243>

- [17] S. Mayadunnage, H. Wijayagunasekara, K. Hemachandra, and L. Nugaliyadde, "Predatory coccinellids (Coleoptera: Coccinellidae) of Vegetable Insect Pests: A Survey in Mid Country of Sri Lanka," *Tropical Agricultural Research*, vol. 19, pp. 69-77, 2007.
- [18] I. L. I. Putra, H. Setiawan, and N. Suprihatini, "Species diversity of ants (Hymenoptera: formicidae) around campus 4, Ahmad Dahlan University, Yogyakarta," *Biospecies*, vol. 14, no. 2, pp. 20-30, 2021.
- [19] T. Abdullah, I. D. Daud, and Kartini, "Testing of various species of Ants (*Solenopsis* sp; *Oecophylla* sp; *Dolichoderus* sp.) against fake white pests (*Cnaphalocrocis medinalis*) in rice plant," *Biome*, vol. 5, pp. 176-185, 2020.
- [20] F. F. Rain, A. F. M. Aslam, H. S. Ringki, N. Sultana, N. Akter, and A. J. Howlader, "Coccinellid predators of aphid and their phylogenetic analysis using COI gene sequences," *International Journal of Applied Sciences and Biotechnology*, vol. 4, no. 3, pp. 408-416, 2016. <https://doi.org/10.3126/ijasbt.v4i3.15782>
- [21] J. C. Moraes, M. M. Goussain, G. A. Carvalho, and R. R. Costa, "Feeding non-preference of the corn leaf aphid *rhopalosiphum maidis* (Fitch, 1856)(Hemiptera: Aphididae) to corn plants (*Zea mays* L.) treated with silicon," *Science and Agrotechnology*, vol. 29, no. 4, pp. 761-766, 2005. <https://doi.org/10.1590/s1413-70542005000400007>
- [22] G. J. Michels Jr and J. H. Matis, "Corn leaf aphid, *rhopalosiphum maidis* (Hemiptera: Aphididae), is a key to greenbug, *Schizaphis graminum* (Hemiptera: Aphididae), biological control in grain sorghum, *Sorghum bicolor*," *European Journal of Entomology*, vol. 105, no. 3, pp. 513-520, 2008. <https://doi.org/10.14411/eje.2008.068>
- [23] M. Malo and J. Hore, "The emerging menace of fall armyworm (*Spodoptera frugiperda* JE Smith) in maize: A call for attention and action," *Journal of Entomology and Zoology Studies*, vol. 8, pp. 455-465, 2020.
- [24] W. Setiawati, N. Gunaeni, T. Uhan, and A. Hasyim, "Potency of predator (*Menochilus sexmaculatus*) augmentation for white fly (*Bemisia tabaci*) management and its effect on gemini virus infestation on tomato," *Indonesian Journal of Agricultural Science*, vol. 13, no. 1, pp. 18-26, 2012. <https://doi.org/10.21082/ijas.v13n1.2012.p18-26>
- [25] I. N. Wijaya, W. Adiantayasa, M. Sritamin, and K. A. Yuliadhi, "Population dynamics of diaphorina citri kuwayama (Homoptera: Psyllidae) and CVPD detection with PCR technique," *Indonesian Journal of Entomology* vol. 7, no. 2, pp. 78-87, 2010. <https://doi.org/10.5994/jei.7.2.78>
- [26] P. A. Abrams, "The evolution of predator-prey interactions: Theory and evidence," *Annual Review of Ecology and Systematics*, vol. 31, pp. 79-105, 2000. <https://doi.org/10.1146/annurev.ecolsys.31.1.79>
- [27] M. Syahrawati, Arneti, and S. Desiska, "Controlling brown planthopper (*Nilaparvata lugens* Stal.) by joint predator (*Pardosa pseudoannulata* boesenberg and strand and *verania lineata* thunberg) under competitive conditions," *Agriculture Cri Journal*, vol. 1, pp. 1-13, 2021.
- [28] S. Kumar, H. Singh, A. Patel, J. N. Patel, and C. Kant, "Brown plant hopper, *nilaparvata iugens* (Stal)(Insecta: Delphacidae) a major insect of rice in India: A review," *Journal of Entomological Research*, vol. 46, no. 2, pp. 333-338, 2022. <https://doi.org/10.5958/0974-4576.2022.00061.5>
- [29] N. Singh and G. Mishra, "Does temperature modify slow and fast development in two aphidophagous ladybirds?," *Journal of Thermal Biology*, vol. 39, pp. 24-31, 2014. <https://doi.org/10.1016/j.jtherbio.2013.11.003>
- [30] M. Saleem, D. Hussain, H. Anwar, M. Saleem, G. Ghouse, and M. Abbas, "Predation efficacy of *menochilus sexmaculatus fabricus* (Coleoptera: coccinellidae) against *macrosiphum rosae* under laboratory conditions," *Journal of Entomology and Zoology Studies*, vol. 2, pp. 160-163, 2014.

# Hasil\_ Ichsan Luqmana

---

## ORIGINALITY REPORT

---

9%

SIMILARITY INDEX

7%

INTERNET SOURCES

4%

PUBLICATIONS

2%

STUDENT PAPERS

---

## PRIMARY SOURCES

---

1	<a href="http://jurnal.unej.ac.id">jurnal.unej.ac.id</a> Internet Source	1%
2	<a href="http://repository.lppm.unila.ac.id">repository.lppm.unila.ac.id</a> Internet Source	1%
3	Submitted to Dicle University Student Paper	1%
4	<a href="http://www.managementjournal.usamv.ro">www.managementjournal.usamv.ro</a> Internet Source	1%
5	T. D. C. Priyadarshani, K. S. Hemachandra, U. G. A. I. Sirisena, H. N. P. Wijayagunasekara. "Developmental biology and feeding efficiency of <i>Menochilus sexmaculatus</i> (Coleoptera: Coccinellidae) (Fabricius) reared on <i>Aphis craccivora</i> (Hemiptera:Aphididae) (Koch)", Tropical Agricultural Research, 2016 Publication	1%
6	<a href="http://jurnal.fkip.unram.ac.id">jurnal.fkip.unram.ac.id</a> Internet Source	1%
7	<a href="http://ejurnal.bppt.go.id">ejurnal.bppt.go.id</a> Internet Source	1%

---

8	<a href="http://media.neliti.com">media.neliti.com</a> Internet Source	<1 %
9	<a href="http://www.ajol.info">www.ajol.info</a> Internet Source	<1 %
10	<a href="http://dokumen.pub">dokumen.pub</a> Internet Source	<1 %
11	<a href="http://mdpi-res.com">mdpi-res.com</a> Internet Source	<1 %
12	<a href="http://www.atlantis-press.com">www.atlantis-press.com</a> Internet Source	<1 %
13	S Herlinda, I M P Simbolon, Hasbi, S Suwandi, Suparman. "Host Plant Species Of The New Invasive Pest, Fall Armyworm (Spodoptera Frugiperda) In South Sumatra", IOP Conference Series: Earth and Environmental Science, 2022 Publication	<1 %
14	<a href="http://dspace.nwu.ac.za">dspace.nwu.ac.za</a> Internet Source	<1 %
15	<a href="http://rp2u.unsyiah.ac.id">rp2u.unsyiah.ac.id</a> Internet Source	<1 %
16	"Ecology and Behaviour of the Ladybird Beetles (Coccinellidae)", Wiley, 2012 Publication	<1 %

17

Kanwer Shahzad Ahmed, Atif Idrees, Muhammad Zeeshan Majeed, Muhammad Irfan Majeed et al. "Synergized Toxicity of Promising Plant Extracts and Synthetic Chemicals against Fall Armyworm Spodoptera frugiperda (JE Smith) (Lepidoptera: Noctuidae) in Pakistan", Agronomy, 2022

Publication

<1 %

18

"Environment and Climate-smart Food Production", Springer Science and Business Media LLC, 2022

Publication

<1 %

Exclude quotes On

Exclude matches Off

Exclude bibliography On