

Biogenesis final

by Oktira .

Submission date: 22-Nov-2023 02:07AM (UTC-0600)

Submission ID: 2236002029

File name: 35724-Article_Text-113341-1-15-20230518.docx (807.56K)

Word count: 4707

Character count: 27511

Pancreatic lipase inhibitory activity of butterfly pea flower (*Clitoria ternatea*) kombucha

ABSTRACT. Preventing the action of pancreatic lipase is believed to be an effective method for treating obesity. Pancreatic lipase inhibitor acts by suppressing the activity of pancreatic lipase, leading to decreased lipid absorption. Kombucha is a traditional fermented drink believed to have numerous health benefits, including anti-obese. It can be produced using a range of substrates, such as butterfly pea flowers. The aim of this research was to investigate the inhibitory activity of kombucha made from butterfly pea flowers toward pancreatic lipase. The fermentation parameters monitored in this study included changes in cell density (OD600), the dry weight of the kombucha mushroom, pH, reducing sugar content, and the % titratable acid. The total phenolic and flavonoid content were also analyzed before and after fermentation. The inhibitory effect of butterfly pea flower kombucha on pancreatic lipase was presented as IC_{50} . The findings indicated that as fermentation progressed, the pH level and amount of reducing sugar decreased while the % titratable acid, cell density (OD600), and dry weight of the kombucha mushroom increased. The phenol and flavonoid content of butterfly pea kombucha was greater than butterfly pea infusion, with respective levels of 0.040 mg GAE/g and 0.017 mg QE/g. This study confirms that butterfly pea flower kombucha, with an IC_{50} value of 162.83 $\mu\text{g/mL}$, has the ability to inhibit pancreatic lipase *in vitro* more effectively compared to butterfly pea infusion, which exhibits an IC_{50} value of 239.39 $\mu\text{g/mL}$. Thus, butterfly pea flower kombucha might be a promising candidate to combat obesity.

Keywords: butterfly pea flower; *Clitoria ternatea*; kombucha; pancreatic lipase.

INTRODUCTION

Obesity is a health problem caused by an imbalance between energy intake and output. Obesity, in terms of health, is related to various diseases such as hypertension, cardiovascular, diabetes, and others (Lunagariya *et al.*, 2014). WHO states that 1.9 billion people are overweight and 650 million of them suffer from obesity (WHO, 2016). About 21.8% of adults in Indonesia suffer from obesity (Kemenkes, 2018). A change in lifestyle is the first approach in the treatment of obesity, but maintaining this effort, in the long run, can be a challenge for obesity sufferers (Wadden *et al.*, 2020). Alternatively, obesity treatment is done using antiobesity agents. One antiobesity agent that works by reducing calorie absorption is pancreatic lipase inhibitors. Pancreatic lipase plays a significant role in fat metabolism, where this enzyme is responsible for 60% of fat absorption (Kim *et al.*, 2016). Orlistat is a common commercial drug used to combat obesity by inhibiting pancreatic lipase. However, some unpleasant side effects may occur such as oily stools, stained underwear, bloating, and inhibition of vitamin K absorption (Liu *et al.*, 2020). It's important to note that this medication is intended for adults and teenagers aged 12-18 and above, and should not be used by pregnant women or individuals with malabsorption syndrome. Some individuals may experience difficulty in achieving positive outcomes from medication if they do not adhere to a consistent dosage schedule and have apprehensions about potential long-term side effects (Behl & Misra, 2017). Therefore, it's important to explore other options like the use of functional foods for obesity treatment (Asgary *et al.*, 2018; Payab *et al.*, 2020).

An example of functional food that has become popular is the fermented drink known as kombucha. Made from fermenting black tea with a microbial culture in the form of a symbiotic relationship between yeast, fungus, and acetic acid bacteria, commonly referred to as SCOBY (Symbiotic Culture of Bacteria and Yeast), kombucha is widely consumed as a beverage believed to enhance overall health (de Miranda *et al.*, 2022). Research has indicated that kombucha possesses properties such as antibacterial, anticancer, anti-diabetic, and antioxidant, can prevent cardiovascular disease, has a hepatoprotective effect, improves digestive function, stimulates the immune system and reduces cholesterol levels (Zubaidah *et al.*, 2019; Watawan *et al.*, 2015). *In vivo* studies on rat also suggest that kombucha has better activity in inhibiting pancreatic lipase activity compared to black tea (Aloulou *et al.*, 2012).

In order to achieve a diverse range of flavors, kombucha is commonly created using a variety of substrates, for example, mint leaves, salak, ginger, grapes, and others (de Miranda *et al.*, 2022). An

ingredient that has gained popularity as a healthful beverage is the butterfly pea flower (*Clitoria ternatea*). Based on some research results on animals, butterfly pea flower extract has properties as an anti-asthma, anti-inflammatory, analgesic, antipyretic, anti-diabetic, anti-lipidemic, anti-rheumatic, and antioxidant (Oguis *et al.*, 2019). Like other natural ingredients, butterfly pea flower also has the potential to be used as a substrate for making kombucha. However, at present, information on its potential as a lipase inhibitor for obesity therapy is unknown. Therefore, the purpose of this research is to determine the lipase inhibitor activity of butterfly pea flower kombucha.

MATERIALS AND METHODS

Samples and kombucha culture collection. Butterfly pea (*Clitoria ternatea*) collected from a market in Yogyakarta, Indonesia. Kombucha culture used as a starter of the fermentation process was obtained from the Microbiology Laboratory, Faculty of Applied Science and Technology, Ahmad Dahlan University.

Kombucha preparation and fermentation process. The production of butterfly pea flower kombucha refers to Ahmed *et al.* (2020), with modifications. One gram of butterfly pea flowers was brewed with 200 mL of boiling water for 15 minutes, then poured and filtered into sterilized glass bottles or containers. The SCOBY used is one week old, it has formed a dense, solid nata with a firm texture and a thickness of 0.3 mm. One gram of butterfly pea flowers was soaked with 200 mL of boiling water for 15 minutes, then poured and filtered into five sterilized glass bottles, each containing 40 mL. Twenty grams of sugar (10% w/v) was added to the bottles containing the solution of butterfly pea flower tea. The bottles are sealed and left to cool to around 30°C. Then, 12 mL of liquid culture and 8 grams of solid culture (SCOBY) were added to each bottle. The bottles are then sealed with sterilized cheesecloth, tied with rubber bands, and incubated for 12 days at room temperature (± 28 -30°C).

Biological determination. To measure the optical density (OD) of the fermented culture, a spectrophotometer was used to read the value at 600 nm. The weight of the kombucha mushroom was determined by separating it from the culture, washing it thrice with distilled water, and drying it at 80°C until it reached a constant weight.

Chemical determination. The pH levels were determined by utilizing an electronic pH meter that had been calibrated at pH 4.0 and 7.0. To determine the titratable acidity (TA), a 10 mL sample of the fermentation broth was taken and heated in a water bath at 100°C for 10 minutes to remove the CO₂. The sample was then diluted to 250 mL and 25 mL of the solution was added to a 100 mL Erlenmeyer flask. 3-5 drops of Phenolphthalein (PP) indicator were added and the solution was titrated with a 0.1 N standard NaOH solution until it turns light pink. The total acidity was expressed as acetic acid (BM = 60). To measure the amount of reducing sugars in the kombucha sample, the Nelson-Somogyi method was employed (Somogyi, 1952). A 1 mL sample was diluted to 100 mL and subsequently mixed with 1 mL of Nelson reagent. After being heated for 20 minutes in a boiling water bath, it was cooled to 25°C in running water and combined with 1 mL of Arsenomolybdate reagent. The mixture was thoroughly agitated until the Cuprous Oxide sediment dissolved and then 7 mL of distilled water was added. Based on a glucose standard curve, the spectrophotometer was used to measure the absorbance at a 540 nm wavelength to determine the amount of reducing sugar in kombucha. The total phenol content in Kombucha fermented solutions was determined using the Folin-Ciocalteu method (Singleton *et al.*, 1999). A 0.1 mL sample was transferred into a 100 mL Erlenmeyer flask and diluted with distilled water to a volume of 46 mL. Next, 1.0 mL of Folin-Ciocalteu reactive solution was added and the mixture was allowed to incubate at room temperature for 3 minutes. After that, 3 milliliters of 2% w/v sodium carbonate solution were added and the mixture was left for 30 minutes. Finally, the absorbance was measured at 760 nm. The total phenol content was reported as gallic acid equivalents using a calibration curve. A colorimetric method was utilized to determine the flavonoid content of the sample. The process began by adding 100 microliters of the extract to 4 milliliters of distilled water. Afterward, 0.3 milliliters of 5% sodium

nitrite was added and allowed to sit for 5 minutes. Next, 0.3 milliliters of 10% aluminum chloride was added and left for another 6 minutes. Then, 2 milliliters of 1M sodium hydroxide was added to the mixture, which was immediately diluted by adding 3.3 milliliters of distilled water and mixed thoroughly. The absorbance was then measured at 510 nanometers against a blank. To establish a calibration curve, a standardized measurement of quercetin was utilized and the gathered data was recorded in milligrams of quercetin equivalents per gram of the sample (mg/g).

In vitro assay for lipase inhibitory activity. The in vitro lipase inhibitory activity test was conducted according to the method performed by Aji *et al.*, (2020) with modifications. Lipase activity was determined by evaluating the conversion of p-nitrophenyl palmitate (pNPP) to p-nitrophenol. Five different concentrations, 50, 100, 150, 200, and 250 ug/mL, were made. Then, a 20 mg/mL enzyme stock was made in a 50 mM Tris HCl pH 8 buffer solution, and a 50 mM substrate stock (pNPP) was made in acetone. The activity test was conducted by mixing 0.1 mL of the 20 mg/mL lipase, 0.2 mL of the extract (at different concentrations), and 0.7 mL of 50 mM Tris HCl pH 8. After mixing, the solution was incubated for 15 minutes at 37°C, and then 0.1 mL of 50 mM pNPP was added and incubated for another 30 minutes at 37°C. The absorbance of the test results was measured using a spectrophotometer at a wavelength of 410 nm. In this experiment, the activity of infuse of butterfly pea flower (without fermentation) and the kombucha of butterfly pea flower was measured. Orlistat was used as a positive control, while aquadest was used as a negative control. The activity of the enzyme was determined by measuring the rate of reaction that produced 1 mmol of p-nitrophenol per minute at a temperature of 37°C. The test was conducted three times. The inhibitory activity of the lipase enzyme was calculated as lipase inhibition (I %) = 100 – (A/B x 100), where A is the activity without inhibitor and B is the activity with inhibitor. The lipase inhibitory activity was presented as the IC₅₀ value, which is the concentration required to reduce enzyme activity by 50%. This value is determined by plotting the substrate concentration and percent inhibition results on a regression equation.

Data analysis. All the analyses were conducted in triplicate. Data were represented as mean ± standard deviation and statistically analyzed using SPSS Statistics 26 Software. Identification of any significant differences in the mean comparison between the two samples, an independent t-test was performed on the data for total phenolic and total flavonoids. Meanwhile, the lipase inhibitory results were analyzed using a one-way analysis of variance (ANOVA) followed by Duncan's Multiple Range test (DMRT).

RESULTS AND DISCUSSION

Biological activity of butterfly pea flower kombucha. In this experiment, kombucha was produced using butterfly pea flowers as the substrate. Mushrooms exhibited exponential growth during the initial 2 or 4 days of fermentation, as seen in Fig. 1. The data demonstrates that as the fermentation time increased, the growth of the Kombucha cultures, as measured by OD600, also increased and reached its peak after 9 days. During the fermentation process, the OD600 of Kombucha increased in the initial four days and continued to rise until the end of the fermentation period (Abou-Taleb *et al.*, 2017). The total count of yeast and acetic acid bacteria in Kombucha liquor increases over time and reaches its peak after 10 days of fermentation (Neffe-Skocińska *et al.*, 2017).

The dry weight of the Kombucha mushroom showed a similar pattern, with a peak output of 4.74 g after 9 days of fermentation. The pattern of the dry weight of Kombucha mushrooms increasing as the fermentation period progresses is linked to the expansion of a cellulosic structure made up of acetic acid bacteria (Ahmed, 2020). The kombucha mushroom, also known as the "SCOBY" or "mother", is a cellulose biofilm composed of a symbiotic colony of bacteria and yeast. The composition of microorganisms in kombucha can differ from one culture to another, but certain types of microorganisms are commonly present in all SCOBYs, including acetic acid bacteria (AAB) and yeast (Bishop, 2022). The number of bacteria present in the liquid portion of kombucha was higher than in the solid portion (SCOBY) (Watawana *et al.*, 2016). The partnership between the acetic acid

bacteria and yeast in kombucha effectively prevents the contamination of spoilage and pathogenic microorganisms (Kapp & Sumner., 2019).




Fig. 1. Optical density (OD 600 nm) and mushrooms dry weight patterns during butterfly pea flower kombucha fermentation.

Chemical composition of butterfly pea flower kombucha. Data shown in Fig. 2 indicates that the pH values decreased as the fermentation period prolonged and reached its lowest point at 2.04 on day-12. In contrast, the total acidity, represented as acetic acid concentration, increased throughout fermentation and reached 0.34% on the 12th day. These results align with Ahmed (2020), who observed that kombucha's pH steadily declined as the fermentation progressed. This decline is due to the presence of organic acid, which is produced as a result of sugar metabolism by bacteria and yeast, leading to an increase in the acidity of the kombucha. The drop in pH results from the microorganisms in kombucha producing more organic acids as the fermentation period extends, reducing pH (Amarasinghe *et al.*, 2018).

The data on reducing sugar concentration, as presented in Fig. 3, demonstrates a decrease in the concentration at the start of the fermentation process after two days, and it slightly decreased until day-12. The decline in reducing sugar in kombucha can be attributed to the ability of microorganisms to break down sugar molecules through hydrolysis (Kitwetcharoen *et al.*, 2023). As the fermentation process progresses, the decreasing level of reducing sugar results from the sugar consumed by microorganisms as a source of carbon, which in turn leads to the production of ethanol and organic acids. Therefore, the acid level in kombucha progressively increases with the prolongation of the fermentation period (Amarasinghe *et al.*, 2018). The acidic environment in kombucha favors acid-tolerant microbes and inhibits the growth of potential competitors or invaders. Certain yeast species, like *Dekkera/Brettanomyces*, can survive and flourish in the acidic conditions of kombucha, which would be harmful to other yeast genera. Similarly, acid-tolerant bacteria, such as *Komagataeibacter*, thrive in kombucha, while other bacteria are less tolerant and unable to survive in highly acidic conditions (May *et al.*, 2019).




Fig. 2. Titratable acidity and pH during butterfly pea flower kombucha fermentation.




Fig. 3. Reducing sugar content during butterfly pea flower kombucha fermentation.

Total phenolic and total flavonoid compound of butterfly pea flower kombucha. The total phenolic content was greater after 12 days of fermentation, as indicated in Table 1., than before fermentation. The findings of this study are consistent with prior research that found kombucha fermented drinks to contain a higher amount of total phenolics than the initial fermentation process (Vitas *et al.*, 2018). Another investigation demonstrated a rise in total phenolic content as the duration of fermentation progressed (Ayed and Hamdi, 2015). The enhancement of phenolic compounds in kombucha was observed after three days due to the increased kinetics growth of microorganisms (Antolak *et al.*, 2021). The variety of tea utilized in kombucha production affects the amount of polyphenols present (Bishop, 2022).

Table 1. Total phenolic and flavonoids before and after butterfly pea flower kombucha fermentation.

Fermentation time (days)	Total phenolic (mg GAE/g sample)	Total flavonoids (mg QE/g sample)
0	0,015 ± 0,002 ^a	0,012 ± 0,002 ^a

3	12	0,040 ± 0,001 ^b	0,017 ± 0,001 ^a
---	----	----------------------------	----------------------------

Mean values are significantly different at $p < 0.05$ (Independent t-test). Values are given in means \pm standard deviation.

The total flavonoid content also shows a similar trend, as seen in Figure 3. The increase of phenols and flavonoids in the final stages of Kombucha fermentation is likely due to the growth and metabolic activity of the yeast and bacteria in the mixture. These microorganisms produce various bioactive compounds, including phenols and flavonoids, which contribute to the kombucha's characteristic flavor and health-promoting properties. The fermentation of kombucha involves several chemical reactions, one of which is the oxidation of polyphenolic compounds by some enzymes, resulting in the formation of flavonoids and other healthy compounds through microbial hydrolysis (Antolak *et al.*, 2021). The Kombucha samples exhibit significant antioxidant capacity due to a high concentration of total phenolic and flavonoid substances (Ivanišová *et al.*, 2019).

In vitro assay for lipase inhibitory activity of Butterfly pea flower Kombucha. The results of the in vitro lipase inhibition study of butterfly pea flower Kombucha are displayed in Fig. 4. The study compared the lipase inhibition activity of the butterfly pea flower infusion (without fermentation), butterfly pea flower Kombucha drink, and orlistat. The butterfly pea flower Kombucha drink exhibited lower lipase inhibition activity compared to orlistat, with an IC_{50} of 239.39.9 $\mu\text{g/mL}$, 162.83 $\mu\text{g/mL}$, and 102.00 $\mu\text{g/mL}$ for the butterfly pea flower infusion (without fermentation), butterfly pea flower Kombucha drink, and orlistat, respectively. However, compared to the butterfly pea flower infusion (without fermentation), the butterfly pea flower kombucha exhibits a superior level of inhibition against pancreatic lipase. The results of this study align with the outcomes of previous research on the fermentation of green vegetables (such as spinach, broccoli, and sweetleaf) using kombucha culture as a functional beverage. The research found that kombucha made from these substrates can inhibit pancreatic lipase (Maryati *et al.*, 2020). Additionally, sea grape kombucha (*Caulerpa racemosa*) has been shown to inhibit pancreatic lipase activity and result in weight loss in experimental mice (Permatasari *et al.*, 2022).




Fig. 4. Inhibitory activities (IC_{50}) of butterfly pea flower infusion, kombucha, and orlistat on pancreatic lipase. The data represented by distinct letters exhibit a significant difference ($P < 0.05$). The values are presented as means \pm standard deviation.

During kombucha fermentation, yeast and bacteria produce various bioactive compounds, including organic acids and polyphenols, which are believed to contribute to the inhibition of lipase activity. Several studies have shown that subclasses of polyphenols, including flavonoids, and

phenolic acids, can effectively inhibit pancreatic lipase (Buchholz & Melzig, 2015). Kombucha contains phenol compounds, the main bioactive compound group in kombucha and responsible for the drink's health benefits (Cardoso *et al.*, 2020). These compounds play an important role in inhibiting pancreatic lipase activity (Buchholz & Melzig, 2015). Moreover, the optimal duration of fermentation produces organic acids, such as glucuronic acid, which serve as secondary metabolites and inhibit pancreatic lipase activity, while simultaneously increasing the polyphenol activity during the process (Maryati *et al.*, 2020). The bioavailability of polyphenols is increased by glucuronic acid (GlcUA). The conjugation of phenols with GlcUA leads to greater transportability and bioavailability of the substances.

Numerous studies have demonstrated that the kombucha fermentation process can enhance the health benefits of herbal infusions (de Miranda *et al.*, 2022). The antioxidant and antimicrobial properties of herbal infusions are improved after being fermented with kombucha culture. The antioxidant activity of lemon balm (*Melissa officinalis* L.) kombucha is higher compared to its herbal infusion (Velicanski *et al.*, 2014). Similarly, in the case of winter savory (*Satureja montana* L.), the antioxidant and antimicrobial effects against both gram-positive and gram-negative bacteria were found to be stronger after fermentation compared to its herbal infusion (Cetojevic-Simin *et al.*, 2012). Furthermore, this study demonstrates that, in contrast to butterfly pea flower infusion, kombucha fermentation results in an increase in lipase inhibitor activity. Therefore, butterfly pea flower kombucha has the potential to be developed as an antiobesity agent that may also offer many health benefits, such as antioxidants and antimicrobials.

CONCLUSION

During the fermentation process of butterfly pea flower kombucha, it was observed that the pH and reducing sugar levels showed a decrease, while there was a noticeable increase in cell density (OD600), dry weight of SCOBY, and % titratable acid. The findings from this study indicate that butterfly pea flower kombucha contains various bioactive compounds such as polyphenols and flavonoids, and can inhibit pancreatic lipase in vitro. It is believed that these active compounds play a role in the lipase inhibitory activity of kombucha, thereby presenting potential as an antiobesity agent. However, it is necessary to obtain further evidence to identify bioactive compounds, performance of in-vivo studies, and comprehend the underlying mechanisms of the results obtained.

ACKNOWLEDGEMENTS

This study was funded by the LPPM UAD grant (PD-162/SP3/LPPM-UAD/VII/2022). Authors would like to thank to Laboratory of Microbiology at the Faculty of Applied Science and Technology, Universitas Ahmad Dahlan for research facilities and support.

REFERENCES

- Abou-Taleb, K. A., Ebeed, N. M., Abd El-salam, S. S., Amin, S. A. 2017. Antimicrobial and antiproliferative, pro-apoptotic actions of Kombucha fermented solutions against colon and hepatocellular cancer cell lines. *World Journal of Pharmacy and Life Sciences*. vol 3: 120-132.
- Ahmed, R. F., Hikal, M. S., Abou-Taleb, K. A. 2020. Biological, chemical and antioxidant activities of different types kombucha. *Annals of Agricultural Sciences*. vol 65(1): 35-41. doi: <https://doi.org/10.1016/j.aos.2020.04.001>
- Aji, O. R., Hudaya, R. A., Putri, D. A. 2020. In vitro pancreatic lipase inhibitor activity of *Mangifera foetida* leaves extract. *Jurnal Biodjati*. vol 6(1): 82-92. doi: <https://doi.org/10.15575/biodjati.v6i1.10646>
- Aloulou, A., Hamden, K., Elloumi, D., Ali, M. B., Hargafi, K., Jaouadi, B., Ayadi, F., Elfeki, A., Ammar, E. 2012. Hypoglycemic and anti-lipidemic properties of kombucha tea in alloxan-induced diabetic rats. *BMC Complementary and Alternative Medicine*. vol 12(1): 63. doi: <https://doi.org/10.1186/1472-6882-12-63>
- Amarasinghe, H., Weerakkody, N. S., Waisundara, V. Y. 2018. Evaluation of physicochemical properties and antioxidant activities of kombucha "Tea Fungus" during extended periods of fermentation. *Food science & nutrition*, vol 6(3): 659-665. doi: <https://doi.org/10.1002/fsn3.605>
- Antolak, H., Piechota, D., Kucharska, A. 2021. Kombucha Tea-A Double Power of Bioactive Compounds from Tea and Symbiotic Culture of Bacteria and Yeasts (SCOBY). *Antioxidants*. vol 10(10): 1541. doi: <https://doi.org/10.3390/antiox10101541>

- Asgary, S., Rastqar, A., Keshvari, M. 2018. Functional food and cardiovascular disease prevention and treatment: A review. *Journal of the American College of Nutrition*. vol 37(5): 429-455. doi: <https://doi.org/10.1080/07315724.2017.1410867>
- Ayed, L., Hamdi, M. 2015. Manufacture of a beverage from cactus pear juice using "tea fungus" fermentation. *Annals of Microbiology*. vol 65: 2293-2299. doi: <https://doi.org/10.1007/s13213-015-1071-8>
- Behl, S., Misra, A. 2017. Management of obesity in adult Asian Indians. *Indian Heart Journal*. vol 69(4): 539-544. doi: <https://doi.org/10.1016/j.ihj.2017.04.015>
- Bishop, P., Pitts, E. R., Budner, D., Thompson-Witrick, K. A. 2022. Kombucha: Biochemical and microbiological impacts on the chemical and flavor profile. *Food Chemistry Advances*, 1, 100025. doi: <https://doi.org/10.1016/j.focha.2022.100025>
- Buchholz, T., Melzig, M. F. 2015. Polyphenolic compounds as pancreatic lipase inhibitors. *Planta Medica*, vol 81(10): 771-783. doi: <https://doi.org/10.1055/s-0035-1546173>
- Cardoso, R. R., Neto, R. O., dos Santos D'Almeida, C. T., do Nascimento, T. P., Pressete, C. G., Azevedo, L., Martino, H. S. D. L., Cameron, C., Ferreira, M. S. L., de Barros, F. A. R. 2020. Kombuchas from green and black teas have different phenolic profile, which impacts their antioxidant capacities, antibacterial and antiproliferative activities. *Food Research International*, vol 128. 108782. doi: <https://doi.org/10.1016/j.foodres.2019.108782>
- Cetojevic-Simin, D. D., A. S. Velicanski, D. D. Cvetkovic, S. L. Markov J. Z. Mrdanovic, V. V. Bogdanovic, S. V. Solajic. 2012. Bioactivity of lemon balm kombucha. *Food and Bioprocess Technology*. vol 5:1756-1765. doi: <https://doi.org/10.1007/s11947-010-0458-6>
- de Miranda, J. F., Ruiz, L. F., Silva, C. B., Uekane, T. M., Silva, K. A., Gonzalez, A., Fernandes, F. F., Lima, A. R. 2022. Kombucha: A review of substrates, regulations, composition, and biological properties. *Journal of food science*, vol 87(2): 503-527. doi: <https://doi.org/10.1111/1750-3841.16029>
- Ivanišová, E., Meňhartová, K., Terentjeva, M., Godočiková, L., Árvay, J., Kačániová, M. 2019. Kombucha tea beverage: microbiological characteristic, antioxidant activity, and phytochemical composition. *Acta Alimentaria*, vol 48(3): 324-331. doi: <http://dx.doi.org/10.1556/066.2019.48.3.7>
- Kapp JM, Sumner W. 2019. Kombucha: a systematic review of the empirical evidence of human health benefit. *Ann Epidemiol*. vol 30:66e70. doi: <https://doi.org/10.1016/j.annepidem.2018.11.001>
- Kemenkes. 2018. Hasil Utama Riskesdas 2018. Kementrian Kesehatan Republik Indonesia. <http://www.depkes.go.id/resources/download/hasilriskesdas>
- Kim, G. N., Shin, M. R., Shin, S. H., Lee, A. R., Lee, J. Y., Seo, B. I., Kim, M. Y., Kim, T. H., Noh, J. S., Rhee, M. H., Roh, S. S. 2016. Study of Antiobesity Effect through Inhibition of Pancreatic Lipase Activity of Diospyros kaki Fruit and Citrus unshiu Peel. *BioMed research international*, 1723042. doi: <https://doi.org/10.1155/2016/1723042>
- Kitwetchareon, H., Tu Phung, L., Klanrit, P., Thanonkeo, S., Tippayawat, P., Yamada, M., Thanonkeo, P. 2023. Kombucha Healthy Drink-Recent Advances in Production, Chemical Composition and Health Benefits. *Fermentation*. vol 9(1), 48. doi: <https://doi.org/10.3390/fermentation9010048>
- Liu, T. T., Liu, X. T., Chen, Q. X., Shi, Y. 2020. Lipase Inhibitors for Obesity: A Review. *Biomedicine & pharmacotherapy = Biomedecine & pharmacotherapie*. vol 128, 110314. doi: <https://doi.org/10.1016/j.biopha.2020.110314>
- Lunagariya, N., Patel, N., Sheha, C., Kamesh, J., Bhutani. 2014. (Review Article) Inhibitor of Pancreatic Lipase: State of Art and Clinical Perspective. *EXCLI*. vol 13(1), 897-921
- Maryati, Y. A. Susilowati, Aspiyanto. 2020. Development of fermented vegetables in their ability to inhibit pancreatic lipase as antidyslipidemic agent. IOP Conference Series: Materials Science and Engineering, 722, 012074. doi: <https://doi.org/10.1088/1757-899X/722/1/012074>
- May, A., Narayanan, S., Alcock, J., Varsani, A., Maley, C., Aktipis, A. 2019. Kombucha: a novel model system for cooperation and conflict in a complex multi-species microbial ecosystem. *PeerJ*. vol 7, e7565. doi: <https://doi.org/10.7717/peerj.7565>
- Neffe-Skocińska, K., Sionek, B., Ścibisz, I., Kołożyn-Krajewska, D. 2017. Acid contents and the effect of fermentation condition of Kombucha tea beverages on physicochemical, microbiological and sensory properties. *CYTA-Journal of Food*. vol 15: 601-607. doi: <https://doi.org/10.1080/19476337.2017.1321588>
- Oguis, G. K., Gilding, E. K., Jackson, M. A., Craik, D. J. 2019. Butterfly Pea (Clitoria ternatea), a Cyclotide-Bearing Plant With Applications in Agriculture and Medicine. *Frontiers in plant science*. vol 10, 645. doi: <https://doi.org/10.3389/fpls.2019.00645>
- Payab, M., Hasani-Ranjbar, S., Shahbal, N., Qorbani, M., Aletaha, A., Haghi-Aminjan, H., Soltani, A., Khatami, F., Nikfar, S., Hassani, S., Abdollahi, M., Larijani, B. 2020. Effect of the herbal medicines in obesity and metabolic syndrome: A systematic review and meta-analysis of clinical trials. *Phytotherapy research : PTR*. vol 34(3): 526-545. doi: <https://doi.org/10.1002/ptr.6547>
- Permatasari, H. K., Firani, N. K., Prijadi, B., Irandi, D. F., Riawan, W., Yusuf, M., Amar, N., Chandra, L. A., Yusuf, V. M., Subali, A. D., Nurkolis, F. 2022. Kombucha drink enriched with sea grapes (Caulerpa racemosa) as potential functional beverage to contrast obesity: An in vivo and in vitro approach. *Clinical nutrition ESPEN*. vol 49: 232-240. doi: <https://doi.org/10.1016/j.clnesp.2022.04.015>

- Singleton, V.L., Orthofer, R. Lamuela-Raventos, R.M. 1999. Analysis of Total Phenols and Other Oxidation Substrates and Antioxidants by Means of Folin-Ciocalteu Reagent. *Methods in Enzymology*. vol 299: 152-178. doi: [http://dx.doi.org/10.1016/S0076-6879\(99\)99017-1](http://dx.doi.org/10.1016/S0076-6879(99)99017-1)
- Somogyi, M. 1952. Notes on sugar determination. *Journal of Biological Chemistry*. vol 195: 19-23. doi: [https://doi.org/10.1016/S0021-9258\(19\)50870-5](https://doi.org/10.1016/S0021-9258(19)50870-5)
- Velicanski, A. S., Cvetkovic, D. D., Markov, S. L., Tumbas Saponjac, V. T., Vulic, J. J. 2014. Antioxidant and antibacterial activity of the beverage obtained by fermentation of sweetened lemon balm (*Melissa officinalis* L.) tea with symbiotic consortium of bacteria and yeasts. *Food Technology and Biotechnology*. vol 52(4): 420-429. doi: <https://doi.org/10.17113/ftb.52.04.14.3611>
- Vitas, J. S., Cvetanović, A. D., Mašković, P. Z., Švarc-Gajić, J. V. Malbaša, R. V. 2018. Chemical composition and biological activity of novel types of kombucha beverages with yarrow. *Journal of Functional Foods*. vol 44:95-102. doi: <https://doi.org/10.1016/j.jff.2018.02.019>
- Wadden, T. A., Tronieri, J. S., Butryn, M. L. 2020. Lifestyle modification approaches for the treatment of obesity in adults. *The American psychologist*. vol 75(2): 235-251. doi: <https://doi.org/10.1037/amp0000517>
- Watawana, M. I., Jayawardena, N., Gunawardhana, C. B., Waisundara, V. Y. 2015. Health, wellness, and safety aspects of the consumption of kombucha. *Journal of Chemistry*. vol 2015, 591869. doi: <https://doi.org/10.1155/2015/591869>
- Watawana, M. I., Jayawardena, N., Gunawardhana, C. B., Waisundara, V. Y. 2016. Enhancement of the antioxidant and starch hydrolase inhibitory activities of king coconut water (*Cocos nucifera* var. *aurantiaca*) by fermentation with kombucha' tea fungus'. *International Journal of Food Science and Technology*. vol 51: 490-498. doi: <https://doi.org/10.1111/ijfs.13006>
- World Health Organization. 2016. Obesity and overweight. <https://www.who.int/news-room/fact-sheets/detail/obesity-and-overweight>
- Zubaidah, E., Ifadah, R.A., Kalsum, U., Lyrawati, D., Putri, W.D.R., Srinta, I. Blanc, P.J. 2019. Anti-diabetes activity of kombucha prepared from different snake fruit cultivars. *Nutrition & Food Science*. vol. 49 (2):333-343. doi: <https://doi.org/10.1108/NFS-07-2018-0201>

Biogenesis final

ORIGINALITY REPORT

12%

SIMILARITY INDEX

8%

INTERNET SOURCES

9%

PUBLICATIONS

1%

STUDENT PAPERS

PRIMARY SOURCES

- | | | |
|---|--|---|
| <div style="background-color: red; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">1</div> | <div style="color: red; font-size: 1.2em; margin: 5px;">Rania F. Ahmed, Mohamed S. Hikal, Khadiga A. Abou-Taleb. "Biological, chemical and antioxidant activities of different types Kombucha", Annals of Agricultural Sciences, 2020</div> <div style="font-size: 0.8em; margin: 5px;">Publication</div> | <div style="font-size: 2.5em; color: red; margin: 5px;">3%</div> |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">2</div> | <div style="color: purple; font-size: 1.2em; margin: 5px;">journal.uinsgd.ac.id</div> <div style="font-size: 0.8em; margin: 5px;">Internet Source</div> | <div style="font-size: 2.5em; color: purple; margin: 5px;">3%</div> |
| <div style="background-color: purple; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">3</div> | <div style="color: purple; font-size: 1.2em; margin: 5px;">www.mspp.org.my</div> <div style="font-size: 0.8em; margin: 5px;">Internet Source</div> | <div style="font-size: 2.5em; color: purple; margin: 5px;">1%</div> |
| <div style="background-color: teal; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">4</div> | <div style="color: teal; font-size: 1.2em; margin: 5px;">Raja Noor Zaliha Raja Abd Rahman. "Anti-obesity Potential of Selected Tropical Plants via Pancreatic Lipase Inhibition", Advances in Obesity, Weight Management & Control, 2017</div> <div style="font-size: 0.8em; margin: 5px;">Publication</div> | <div style="font-size: 2.5em; color: teal; margin: 5px;">1%</div> |
| <div style="background-color: green; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">5</div> | <div style="color: green; font-size: 1.2em; margin: 5px;">www.jmbfs.org</div> <div style="font-size: 0.8em; margin: 5px;">Internet Source</div> | <div style="font-size: 2.5em; color: green; margin: 5px;">1%</div> |
| <div style="background-color: brown; color: white; width: 40px; height: 40px; display: flex; align-items: center; justify-content: center; margin: 5px;">6</div> | <div style="color: brown; font-size: 1.2em; margin: 5px;">Li Chen, I Chen, Pei Chen, Ping Huang. "Application of Butterfly Pea Flower Extract in</div> | <div style="font-size: 2.5em; color: brown; margin: 5px;">1%</div> |

Mask Development", Scientia Pharmaceutica, 2018

Publication

7

Rafidah Mohd Ariff, Xin Yi Chai, Lee Sin Chang, Shazrul Fazry, Babul Airianah Othman, Abdul Salam Babji, Seng Joe Lim. "Recent trends in Kombucha: Conventional and alternative fermentation in development of novel beverage", Food Bioscience, 2023

Publication

1 %

8

Happy Kurnia Permatasari, Novi Khila Firani, Bambang Prijadi, Dicky Faizal Irnandi et al. "Kombucha drink enriched with sea grapes (*Caulerpa racemosa*) as potential functional beverage to contrast obesity: An invivo and invitro approach", Clinical Nutrition ESPEN, 2022

Publication

1 %

9

Eun-Hye Lee, Hye-Jung Yeom, Mi-Sun Ha, Dong-Ho Bae. "Development of banana peel jelly and its antioxidant and textural properties", Food Science and Biotechnology, 2010

Publication

1 %

10

Jeniffer Ferreira de Miranda, Larissa Fernandes Ruiz, Cíntia Borges Silva, Thais Matsue Uekane et al. "Kombucha: A review of substrates, regulations, composition, and

1 %

biological properties", Journal of Food Science, 2022

Publication

11

www.isisn.org

Internet Source

1 %

12

www.mdpi.com

Internet Source

1 %

Exclude quotes Off

Exclude bibliography On